

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
9 August 2001 (09.08.2001)

PCT

(10) International Publication Number  
**WO 01/56990 A2**

(51) International Patent Classification<sup>7</sup>: **C07D 213/42**,  
213/40, 405/12, 213/643, 213/89, 213/38, 409/12, A61K  
31/4406, 31/4427, A61P 25/06, 25/18, 25/08

(21) International Application Number: PCT/US01/00643

(22) International Filing Date: 22 January 2001 (22.01.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/180,047 3 February 2000 (03.02.2000) US  
60/180,089 3 February 2000 (03.02.2000) US

(71) Applicant (for all designated States except US): **ELI LILLY AND COMPANY** [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **COLEMAN, Darrell, Stephen** [US/US]; 1036 Waterford Forest Circle, Cary, NC 27513 (US). **JAGDMANN, Gunnar, Erik, Junior** [US/US]; 1027 Oakgate Court, Apex, NC 27502 (US). **JOHNSON, Kirk, Willis** [US/US]; 1177 Helford Lane, Carmel, IN 46032 (US). **JOHNSON, Michael, Parvin** [US/US]; 4308 Powderhorn Court, Carmel, IN 46033 (US). **LARGE, Thomas, Hallett** [US/US]; 1357 James Court, Zionsville, IN 46077 (US). **MONN, James, Allen** [US/US]; 7518 Allenwood Court, Indianapolis, IN 46268 (US). **SCHOEPP, Darryle, Darwin** [US/US]; 638 North Senate Avenue, Indianapolis, IN 46202 (US). **TIZZANO, Joseph, Patrick** [US/US]; 6682 West Sweet Creek Drive, New Palestine, IN 46163 (US). **BARDA, David, Anthony** [US/US]; 6857 Washington Boulevard

West Drive, Indianapolis, IN 46220 (US). **BRITTON, Thomas, Charles** [US/US]; 4700 Royal Oak Lane, Carmel, IN 46033 (US). **DRESSMAN, Bruce, Anthony** [US/US]; 1430 North New Jersey, Indianapolis, IN 46202 (US). **FICHTNER, Michael, William** [US/US]; 6405 Stonecreek Drive, Indianapolis, IN 46268 (US). **HENRY, Steven, Scott** [US/US]; 3362 South Southway Drive, New Palestine, IN 46163 (US). **HORNBACK, William, Joseph** [US/US]; 10063 Bent Tree Lane, Fishers, IN 46038 (US).

(74) Agents: **STEMERICK, David, M.** et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: POTENTIATORS OF GLUTAMATE RECEPTORS



(57) Abstract: The present invention relates to potentiators of metabotropic glutamate receptor function and specifically provides compounds of formula I, compositions thereof and methods of using the same.

-1-

## POTENTIATORS OF GLUTAMATE RECEPTORS

The present invention provides potentiators of glutamate receptors (compounds of formula I), pharmaceutical compositions thereof, and methods of using the same, processes for preparing the same, and intermediates thereof.

### BACKGROUND OF THE INVENTION

The excitatory amino acid L-glutamate (sometimes referred to herein simply as glutamate) through its many receptors mediates most of the excitatory neurotransmission within the mammalian central nervous system (CNS). The excitatory amino acids, including glutamate, are of great physiological importance, playing a role in a variety of physiological processes, such as long-term potentiation (learning and memory), the development of synaptic plasticity, motor control, respiration, cardiovascular regulation, and sensory perception.

Glutamate acts via at least two distinct classes of receptors. One class is composed of the ionotropic glutamate (iGlu) receptors that act as ligand-gated ionic channels. Via activation of the iGlu receptors, glutamate is thought to regulate fast neuronal transmission within the synapse of two connecting neurons in the CNS. The second general type of receptor is the G-protein or second messenger-linked "metabotropic" glutamate (mGlu) receptor. Both types of receptors appear not only to mediate normal synaptic transmission along excitatory pathways, but also participate in the modification of synaptic connections during development and throughout life. Schoepp, Bockaert, and Sladeczek, *Trends in Pharmacol. Sci.*, 11, 508 (1990); McDonald and Johnson, *Brain Research Reviews*, 15, 41 (1990).

The present invention relates to potentiators of mGlu receptors. The mGlu receptors belong to the Type III G-protein coupled receptor (GPCR) superfamily. This superfamily of GPCR's, including the calcium-sensing receptors, GABA<sub>B</sub> receptors and pheromone receptors, which

-2-

are unique in that they are activated by binding of effectors to the amino-terminus portion of the receptor protein. The mGlu receptors are thought to mediate glutamate's demonstrated ability to modulate intracellular signal transduction pathways. Ozawa, Kamiya and Tsuzuski, *Prog. Neurobio.*, **54**, 581 (1998). They have been demonstrated to be localized both pre- and post-synaptically where they can regulate neurotransmitter release, either glutamate or other neurotransmitters, or modify the post-synaptic response of neurotransmitters, respectively.

At present, there are eight distinct mGlu receptors that have been positively identified, cloned, and their sequences reported. These are further subdivided based on their amino acid sequence homology, their ability to effect certain signal transduction mechanisms, and their known pharmacological properties. Ozawa, Kamiya and Tsuzuski, *Prog. Neurobio.*, **54**, 581 (1998). For instance, the Group I mGlu receptors, which include the mGlu<sub>1</sub> and mGlu<sub>5</sub>, are known to activate phospholipase C (PLC) via Gαq-proteins thereby resulting in the increased hydrolysis of phosphoinositides and intracellular calcium mobilization. There are several compounds that are reported to activate the Group I mGlu receptors including DHPG, (R/S)-3,5-dihydroxyphenylglycine. Schoepp, Goldworthy, Johnson, Salhoff and Baker, *J. Neurochem.*, **63**, 769 (1994); Ito, et al., *Neurorep.*, **3**, 1013 (1992). The Group II mGlu receptors consist of the two distinct receptors, mGlu<sub>2</sub> and mGlu<sub>3</sub> receptors. Both have been found to be negatively coupled to adenylate cyclase via activation of Gαi-protein. These receptors can be activated by a group-selective compound such as 1S,2S,5R,6S-2-aminobicyclo[3.1.0]hexane-2,6-dicarboxylate. Monn, et al., *J. Med. Chem.*, **40**, 528 (1997); Schoepp, et al., *Neuropharmacol.*, **36**, 1 (1997). Similarly, the Group III mGlu receptors, including mGlu<sub>4</sub>, mGlu<sub>6</sub>, mGlu<sub>7</sub> and mGlu<sub>8</sub>, are negatively coupled to adenylate cyclase via Gαi and are potently activated by L-AP<sub>4</sub> (L-(+)-2-amino-4-

-3-


phosphonobutyric acid). Schoepp, *Neurochem. Int.*, **24**, 439 (1994).

It should be noted that many of the available pharmacological tools are not ideal in that they cross react not only on the receptors within a Group of mGlu receptors but also often have some activity between Groups of mGlu receptors. For instance, compounds such as 1S,3R-ACPD, (1S,3R)-1-aminocyclopentane-*trans*-1,3-dicarboxylic acid, are believed to activate all of the Group I, II and III mGlu receptors depending upon the dose utilized while others, such as 1S,3S-ACPD, (1S,3S)-1-aminocyclopentane-*trans*-1,3-dicarboxylic acid, are more selective for the Group II receptors (mGlu<sub>2/3</sub>) than the Group I (mGlu<sub>1/5</sub>) or Group III (mGlu<sub>4/6/7/8</sub>). Schoepp, *Neurochem. Int.*, **24**, 439 (1994). To date, there are very few examples of selective agents for the mGlu receptors. Schoepp, Jane, and Monn, *Neuropharmacol.*, **38**, 1431 (1999).

It has become increasingly clear that there is a link between modulation of excitatory amino acid receptors, including the glutamatergic system, through changes in glutamate release or alteration in postsynaptic receptor activation, and a variety of neurological and psychiatric disorders. e.g. Monaghan, Bridges and Cotman, *Ann. Rev. Pharmacol. Toxicol.*, **29**, 365-402 (1989); Schoepp and Sacann, *Neurobio. Aging*, **15**, 261-263 (1994); Meldrum and Garthwaite, *Tr. Pharmacol. Sci.*, **11**, 379-387 (1990). The medical consequences of such glutamate dysfunction makes the abatement of these neurological processes an important therapeutic goal.

#### SUMMARY OF THE INVENTION

This invention provides compounds of formula I:



formula I

wherein



-4-

R<sub>1</sub> is selected from the group consisting of -C(O)R<sub>3</sub>,  
-C(O)OR<sub>4</sub>, and -SO<sub>2</sub>R<sub>5</sub>

wherein R<sub>3</sub> is selected from the group consisting  
of alkyl and cycloalkyl, R<sub>4</sub> is selected from the  
group consisting of alkyl and cycloalkyl, R<sub>5</sub> is  
selected from the group consisting of alkyl,  
cycloalkyl, and fluorinated alkyl;

R<sub>2</sub> is from 1 to 3 substituents independently selected  
from the group consisting of hydrogen, hydroxy,  
trisubstituted silyloxy, alkyl, substituted alkyl,  
alkenyl, substituted alkenyl, alkoxy, substituted  
alkoxy, cycloalkoxy, substituted cycloalkoxy,  
cycloalkyl, substituted cycloalkyl, halogen, cyano,  
nitro, phenyl, substituted phenyl, pyridyloxy,  
thiophenoxy, substituted thiophenoxy, phenylsulfinyl,  
substituted phenylsulfinyl, phenylsulfonyl, substituted  
phenylsulfonyl, benzoyl, substituted benzoyl, phenoxy,  
and substituted phenoxy;

or

two R<sub>2</sub> substituents are taken together, on adjacent  
positions, to form a fused cycloalkyl or a  
methylenedioxy ring, and one R<sub>2</sub> substituent is selected  
from the group consisting of hydrogen, hydroxy,  
trisubstituted silyloxy, alkyl, substituted alkyl,  
alkenyl, substituted alkenyl, alkoxy, substituted  
alkoxy, cycloalkoxy, substituted cycloalkoxy,  
cycloalkyl, substituted cycloalkyl, halogen, cyano,  
nitro, phenyl, substituted phenyl, pyridyloxy,  
thiophenoxy, substituted thiophenoxy, phenylsulfinyl,  
substituted phenylsulfinyl, phenylsulfonyl, substituted  
phenylsulfonyl, benzoyl, substituted benzoyl, phenoxy,  
and substituted phenoxy;

-5-

R<sub>6</sub> is from 1 to 2 substituents independently selected from the group consisting of hydrogen, alkyl, alkoxy, trifluoromethyl, halogen, phenoxy, and substituted phenoxy;

5

X is selected from the group consisting of a bond, -CH<sub>2</sub>-, -CHR<sub>7</sub>-, and -CH<sub>2</sub>CH<sub>2</sub>- wherein R<sub>7</sub> is lower alkyl;

10

Y is selected from the group consisting of a bond, -CH<sub>2</sub>-, -CHR<sub>8</sub>-, -CH<sub>2</sub>CH<sub>2</sub>-, -CHR<sub>9</sub>CH<sub>2</sub>-, and -CH<sub>2</sub>CHR<sub>9</sub> wherein R<sub>8</sub> is lower alkyl and R<sub>9</sub> is lower alkyl;

and the pharmaceutically acceptable salts thereof and the pyridyl N-oxide thereof.

15

The present invention also provides for novel pharmaceutical compositions, comprising: a compound of the formula I and a pharmaceutically acceptable diluent.

Because the compounds of formula I enhance the normal physiological function of the mGlu receptors, the compounds of formula I are useful for the treatment of a variety of neurological and psychiatric disorders associated with glutamate dysfunction, including: acute neurological and psychiatric disorders such as cerebral deficits subsequent to cardiac bypass surgery and grafting, stroke, cerebral ischemia, spinal cord trauma, head trauma, perinatal hypoxia, cardiac arrest, hypoglycemic neuronal damage, dementia (including AIDS-induced dementia), Alzheimer's disease, Huntington's Chorea, amyotrophic lateral sclerosis, ocular damage, retinopathy, cognitive disorders, idiopathic and drug-induced Parkinson's disease, muscular spasms and disorders associated with muscular spasticity including tremors, epilepsy, convulsions, migraine (including migraine headache), urinary incontinence, substance tolerance, substance withdrawal (including, substances such as opiates, nicotine, tobacco products, alcohol, benzodiazepines, cocaine, sedatives, hypnotics, etc.), psychosis,

35

-6-

schizophrenia, anxiety (including generalized anxiety disorder, panic disorder, and obsessive compulsive disorder), mood disorders (including depression, mania, bipolar disorders), trigeminal neuralgia, hearing loss, 5 tinnitus, macular degeneration of the eye, emesis, brain edema, pain (including acute and chronic pain states, severe pain, intractable pain, neuropathic pain, and post-traumatic pain), tardive dyskinesia, sleep disorders (including narcolepsy), attention deficit/hyperactivity disorder, and 10 conduct disorder.

In another embodiment the present invention provides methods of treating neurological and psychiatric disorders associated with glutamate dysfunction, comprising: administering to a patient in need thereof an effective 15 amount of a compound of formula I. That is, the present invention provides for the use of a compound of formula I or pharmaceutical composition thereof for the treatment neurological and psychiatric disorders associated with glutamate dysfunction.

20 Of the disorders above, the treatment of migraine, anxiety, schizophrenia, and epilepsy are of particular importance.

In a preferred embodiment the present invention provides a method for treating migraine, comprising: 25 administering to a patient in need thereof an effective amount of a compound of formula I.

In another preferred embodiment the present invention provides a method for treating anxiety, comprising: administering to a patient in need thereof an effective 30 amount of a compound of formula I. Particularly preferred anxiety disorders are generalized anxiety disorder, panic disorder, and obsessive compulsive disorder.

In another preferred embodiment the present invention provides a method for treating schizophrenia, comprising: 35 administering to a patient in need thereof an effective amount of a compound of formula I.

-7-

In yet another preferred embodiment the present invention provides a method for treating epilepsy, comprising: administering to a patient in need thereof an effective amount of a compound of formula I.

5 In another embodiment of the present invention provides methods of treating neurological and psychiatric disorders associated with glutamate dysfunction, comprising: administering to a patient in need thereof an effective amount of a potentiator of metabotropic glutamate receptors.

10 Specifically, the present invention provides a method of treating neurological and psychiatric disorders associated with glutamate dysfunction, comprising: administering to a patient in need thereof an effective amount of a potentiator of mGlu<sub>2</sub> and/or mGlu<sub>3</sub> receptors.

15 In a preferred embodiment the present invention provides a method for treating migraine, comprising: administering to a patient in need thereof an effective amount of a metabotropic glutamate potentiator, in particular a potentiator of mGlu<sub>2</sub> and/or mGlu<sub>3</sub> receptors.

20 In another preferred embodiment the present invention provides a method for treating anxiety, comprising: administering to a patient in need thereof an effective amount of a metabotropic glutamate potentiator, in particular a potentiator of mGlu<sub>2</sub> and/or mGlu<sub>3</sub> receptors.

25 Particularly preferred anxiety disorders are generalized anxiety disorder, panic disorder, and obsessive compulsive disorder.

In another preferred embodiment the present invention provides a method for treating schizophrenia, comprising:  
30 administering to a patient in need thereof an effective amount of a metabotropic glutamate potentiator, in particular a potentiator of mGlu<sub>2</sub> and/or mGlu<sub>3</sub> receptors.

In yet another preferred embodiment the present invention provides a method for treating epilepsy,  
35 comprising: administering to a patient in need thereof an effective amount of a metabotropic glutamate potentiator, in particular a potentiator of mGlu<sub>2</sub> and/or mGlu<sub>3</sub> receptors.

-8-

Because such potentiators, including the compounds of formula I, positively modulate metabotropic glutamate receptor response to glutamate, it is an advantage that the present methods utilize endogenous glutamate.

5       Because such potentiators positively modulate metabotropic glutamate receptor response to glutamate agonists it is understood that the present invention extends to the treatment of neurological and psychiatric disorders associated with glutamate dysfunction by administering an  
10       effective amount of a metabotropic glutamate potentiator, including the compounds of formula I, in combination with a potentiated amount of a metabotropic glutamate receptor agonist. Such a combination may be advantageous in that it may augment the activity and selectivity of an agonist of  
15       metabotropic glutamate receptors, in particular a potentiator of mGlu<sub>2</sub> and/or mGlu<sub>3</sub> receptors.

#### DETAILED DESCRIPTION OF THE INVENTION

This invention provides methods of potentiating metabotropic glutamate receptors, in particular a mGlu<sub>2</sub>  
20       and/or mGlu<sub>3</sub> receptors. In the present methods an effective amount of a potentiator of metabotropic glutamate receptors, including a compound of formula I, is administered which positively modulates the effect of glutamate or glutamate agonists on the subject receptors.

25       Before describing the present invention in greater detail, it is understood that the invention in its broadest sense is not limited to particular embodiments described herein, as variations of the particular embodiments described herein are within the scope of the claimed  
30       invention.

Thus, compounds useful in the present invention are those which are potentiators of metabotropic glutamate receptors, particularly, those that potentiate the effects of glutamate and glutamate agonists at Group II metabotropic  
35       glutamate receptors, and even more particularly, those that potentiate the effects of glutamate and glutamate agonists at mGlu<sub>2</sub> receptors and/or mGlu<sub>3</sub> receptors. Useful compounds

-9-

are varied in structure, and so long as they embrace the above properties, they are suitable for use in the present invention. Preferred compounds include, but are not limited to, those described herein.

5       The compounds of formula I potentiate the function of glutamate receptors. Specifically, the compounds of formula I are potentiators of the Group II metabotropic glutamate receptors, and in particular, the compounds of formula I are potentiators of the mGlu<sub>2</sub> receptor and/or the mGlu<sub>3</sub>  
10 receptor.

As used herein, the following terms have the meanings indicated:

The term "alkyl" refers to a straight or branched alkyl chain having from one to ten carbon atoms, and includes  
15 methyl, ethyl, propyl, iso-propyl, butyl, iso-butyl, sec-butyl, t-butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, and the like.

The term "substituted alkyl" refers to a straight or branched alkyl chain having from one to ten carbon atoms,  
20 and includes methyl, ethyl, propyl, iso-propyl, butyl, iso-butyl, sec-butyl, t-butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, and the like having from 1 to 3 substituents selected from the group consisting of hydroxy, halogen, alkoxy, acyloxy, carboxy, alkoxycarbonyl, amido, substituted  
25 amido, amino, acylamino, sulfonylamido, sulfonamide, phenyl, substituted phenyl, phenoxy, substituted phenoxy, benzyloxy, substituted benzyloxy, pyridyl, substituted pyridyl, thienyl, and substituted thienyl.

The term "lower alkyl" refers to a straight or branched  
30 alkyl chain having from one to four carbon atoms, and includes methyl, ethyl, propyl, iso-propyl, butyl, iso-butyl, sec-butyl, and t-butyl.

The term "alkenyl" refers to a straight or branched alkyl chain having from one to ten carbon atoms and one or  
35 more carbon-carbon double bonds, and includes ethylene, propylene, iso-propylene, butylene, iso-butylene, sec-

-10-


butylene, pentylene, hexylene, heptylene, octylene, nonylene, decylene, and the like.

The term "substituted alkenyl" refers to a straight or branched alkyl chain having from one to ten carbon atoms and one or more carbon-carbon double bonds, and includes ethylene, propylene, iso-propylene, butylene, iso-butylene, sec-butylene, pentylene, hexylene, heptylene, octylene, nonylene, decylene, and the like having from 1 to 3 substituents selected from the group consisting of carboxy, alkoxy, carbonyl, amido, substituted amido, phenyl, and substituted phenyl.

The term "cycloalkyl" refers to saturated cyclic alkyl group having from three to seven carbon atoms and includes, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl.

The term "substituted cycloalkyl" refers to saturated cyclic alkyl group having from three to seven carbon atoms and includes, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl having from 1 to 3 substituents selected from the group consisting of hydroxy, halogen, alkoxy, acyloxy, carboxy, alkoxy, carbonyl, amido, substituted amido, amino, acylamino, sulfonylamido, sulfonamide, phenyl, substituted phenyl, phenoxy, substituted phenoxy, benzyloxy, and substituted benzyloxy.

The terms "benzyloxy and substituted benzyloxy" refer to a radical of the formula



wherein R<sub>b</sub> is from 1 to 3 groups independently selected from the group consisting of hydrogen, alkyl, alkoxy, cyano, nitro, trifluoromethyl, halogen, phenyl, and substituted phenyl. The substituent is benzyloxy where each R<sub>b</sub> is hydrogen and is a substituted benzyloxy where at least one R<sub>b</sub> is not hydrogen.

-11-

The term "halogen" refers to a chloro, fluoro, bromo or iodo atom.

The term "fluorinated alkyl" refers to a straight or branched alkyl chain having from one to ten carbon atoms substituted with one or more fluorine atoms, and includes fluoromethyl, difluoromethyl, trifluoromethyl, 2,2,2-trifluoroethyl, 1,1,2,2,2-pentafluoroethyl 3,3,3-trifluoropropyl, 4,4,4-trifluorobutyl, and the like.

The term "alkoxy" refers to straight or branched alkyl chain having from one to ten carbon atoms attached to an oxygen atom, and includes methoxy, ethoxy, propoxy, isopropoxy, butoxy, iso-butoxy, sec-butoxy, t-butoxy, pentoxy, hexoxy, heptoxy, octoxy, nonoxy, decoxy, and the like.

The term "substituted alkoxy" refers to a straight or branched alkyl chain having from one to ten carbon atoms attached to an oxygen atom, and includes methoxy, ethoxy, propoxy, iso-propoxy, butoxy, iso-butoxy, sec-butoxy, t-butoxy, pentoxy, hexoxy, heptoxy, octoxy, nonoxy, decoxy, and the like having from 1 to 3 substituents selected from the group consisting of hydroxy, halogen, alkoxy, acyloxy, carboxy, alkoxycarbonyl, amido, substituted amido, amino, acylamino, sulfonylamido, sulfonamide, phenyl, substituted phenyl, phenoxy, substituted phenoxy, benzyloxy, and substituted benzyloxy; and when one or more of the substituents is hydroxy, halogen, alkoxy, acyloxy, amino, acylamino, sulfonamide, phenoxy, substituted phenoxy, benzyloxy, or substituted benzyloxy then those substituents are not attached to the same carbon as the alkoxy oxygen atom.

The term "cycloalkoxy" refers to saturated cyclic alkyl group having from three to seven carbon atoms attached to an oxygen atom, and includes, cyclopropyloxy, cyclobutyloxy, cyclopentyloxy, cyclohexyloxy, and cycloheptyloxy.


The term "substituted cycloalkoxy" refers to saturated cyclic alkyl group having from three to seven carbon atoms attached to an oxygen atom, and includes, cyclopropyloxy, cyclobutyloxy, cyclopentyloxy, cyclohexyloxy, and



-12-


cycloheptyloxy having from 1 to 3 substituents selected from the group consisting of hydroxy, halogen, alkoxy, acyloxy, carboxy, alkoxycarbonyl, amido, substituted amido, amino, acylamino, sulfonylamido, sulfonamide, phenyl, substituted phenyl, phenoxy, substituted phenoxy, benzyloxy, and substituted benzyloxy; and when one or more of the substituents is hydroxy, halogen, alkoxy, acyloxy, amino, acylamino, sulfonamide, phenoxy, substituted phenoxy, benzyloxy, or substituted benzyloxy then those substituents are not attached to the same carbon as the cycloalkoxy oxygen atom.

The term "pyridyloxy" refers to a radical of the formula



wherein G is selected from the group consisting of pyrid-2-yl, pyrid-3-yl, and pyrid-4-yl.


The terms "phenoxy and substituted phenoxy" refer to a radical of the formula



wherein  $R_c$  is from 1 to 3 groups independently selected from the group consisting of hydrogen, hydroxy, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, alkoxy, substituted alkoxy, halogen, acyloxy, carboxy, alkoxycarbonyl, amido, substituted amido, amino, acylamino, sulfonylamido, sulfonamide, cyano, nitro, phenyl, substituted phenyl, and  $-S(O)_p$ alkyl (wherein  $p$  is 0, 1, or 2). The substituent is phenoxy where each  $R_c$  is hydrogen and is a substituted phenoxy where at least one  $R_c$  is not hydrogen.


The terms "thiophenoxy and substituted thiophenoxy" refer to a radical of the formula

-13-




wherein  $R_d$  is from 1 to 3 groups independently selected from the group consisting of hydrogen, hydroxy, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, alkoxy, substituted alkoxy, halogen, acyloxy, carboxy, alkoxycarbonyl, amido, substituted amido, amino, acylamino, sulfonylamido, sulfonamide, cyano, nitro, phenyl, substituted phenyl, and  $-S(O)_p$ alkyl (wherein  $p$  is 0, 1, or 2). The substituent is thiophenoxy where each  $R_d$  is hydrogen and is a substituted thiophenoxy where at least one  $R_d$  is not hydrogen.

The term "trisubstituted silyloxy" refer to a radical of the formula



wherein  $R_f$  is independently selected from the group consisting of alkyl, phenyl, and benzyl.

The terms "phenylsulfinyl and substituted phenylsulfinyl" refer to a radical of the formula




wherein  $R_g$  is from 1 to 3 groups independently selected from the group consisting of hydrogen, hydroxy, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, alkoxy, substituted alkoxy, halogen, acyloxy, carboxy, alkoxycarbonyl, amido, substituted amido, amino, acylamino, sulfonylamido, sulfonamide, cyano, nitro, phenyl,

-14-


substituted phenyl, and  $-S(O)_p\text{alkyl}$  (wherein  $p$  is 0, 1, or 2). The substituent is phenylsulfinyl where each  $R_g$  is hydrogen and is a substituted phenylsulfinyl where at least one  $R_g$  is not hydrogen.

5        The term "phenylsulfonyl and substituted phenylsulfonyl, refer to a radical of the formula



10        wherein  $R_h$  is from 1 to 3 groups independently selected from the group consisting of hydrogen, hydroxy, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, alkoxy, substituted alkoxy, halogen, acyloxy, carboxy, alkoxycarbonyl, amido, substituted amido, amino, acylamino, sulfonylamido, sulfonamide, cyano, nitro, phenyl, substituted phenyl, and  $-S(O)_p\text{alkyl}$  (wherein  $p$  is 0, 1, or 2). The substituent is phenylsulfonyl is where each  $R_h$  is hydrogen and is a substituted phenylsulfonyl where at least one  $R_h$  is not hydrogen.

20        The term "benzoyl and substituted benzoyl refer to a radical of the formula




25        wherein  $R_i$  is from 1 to 3 groups independently selected from the group consisting of hydrogen, hydroxy, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, alkoxy, substituted alkoxy, halogen, acyloxy, carboxy, alkoxycarbonyl, amido, substituted amido, amino, acylamino,

-15-


sulfonylamido, sulfonamide, cyano, nitro, phenyl, substituted phenyl, and  $-S(O)_p$ alkyl (wherein  $p$  is 0, 1, or 2). The substituent is benzoyl where each  $R_i$  is hydrogen and is a substituted benzoyl where at least one  $R_i$  is not hydrogen.

The term "acyloxy" refers to a radical of the formula




wherein  $R_j$  is selected from the group consisting of alkyl, phenyl, and benzyl.

The term "carboxy" refers to a radical of the formula




The term "alkoxycarbonyl" refers to a radical of the formula




wherein  $R_k$  is selected from the group consisting of alkyl, benzyl, and substituted benzyl.

The term "amido" refers to a radical of the formula



The term "substituted amido" refers to a radical of the formula




wherein  $R_m$  is selected from the group consisting of alkyl and benzyl and  $R_n$  is selected from the group

-16-


consisting of hydrogen, alkyl, benzyl, substituted benzyl, phenyl and substituted phenyl.

The term "acylamino" refers to a radical of the formula



5        wherein  $R_o$  is selected from the group consisting of alkyl, phenyl, substituted phenyl, benzyl, and substituted benzyl.


The term "sulfonylamido" refers to a radical of the formula



10

wherein  $R_p$  is selected from the group consisting of alkyl, phenyl, and substituted phenyl; and  $R_{p'}$  is selected from the group consisting of hydrogen and alkyl.

The term "sulfonamide" refers to a radical of the formula




15

wherein  $R_q$  is selected from the group consisting of alkyl, phenyl, and substituted phenyl.

The terms "phenyl and substituted phenyl" refer to a radical of the formula

20




wherein  $R_r$  is from 1 to 3 groups independently selected from the group consisting of hydrogen, hydroxy, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl,

-17-

alkoxy, substituted alkoxy, halogen, acyloxy, carboxy, alkoxy carbonyl, amido, substituted amido, amino, acylamino, sulfonylamido, sulfonamide, cyano, nitro, phenyl, substituted phenyl, and  $-S(O)_p$ alkyl (wherein  $p$  is 0, 1, or 2). The substituent is phenyl where each  $R_x$  is hydrogen and is a substituted phenyl where at least one  $R_x$  is not hydrogen.

The terms "pyridyl and substituted pyridyl" refer to a radical of the formula




10

wherein  $R_s$  is from 1 to 2 groups independently selected from the group consisting of hydrogen, alkyl, halogen, and trifluoromethyl. The substituent is pyridyl where each  $R_s$  is hydrogen and is a substituted pyridyl where at least one  $R_s$  is not hydrogen. It is understood that the radical may be attached by the 2-, 3-, or 4-positions.

15

The terms "thienyl and substituted thienyl" refer to a radical of the formula




20

wherein  $R_t$  is from 1 to 2 groups independently selected from the group consisting of hydrogen and alkyl. The substituent is thienyl where each  $R_t$  is hydrogen and is a substituted thienyl where at least one  $R_t$  is not hydrogen. It is understood that the radical may be attached by the 2- or 3-positions.

25

The terms "benzyl or substituted benzyl" refer to a radical of the formula


-18-



wherein  $R_e$  is from selected from the group consisting of hydrogen, hydroxy, lower alkyl, and fluoro; and


wherein  $R_a$  is from 1 to 3 groups independently selected  
 5 from the group consisting of hydrogen, alkyl, alkoxy, cyano, nitro, trifluoromethyl, and halogen. The substituent is benzyl where each  $R_a$  is hydrogen (which term includes compounds having all the allowed values of  $R_e$ ) and is a substituted benzyl where at least one  $R_a$  is not hydrogen,  
 10 including those compounds having all the allowed values of  $R_e$ .

Two  $R_2$  substituents taken together, on adjacent positions, to form a "fused cycloalkyl" having from 5 to 7 carbon atoms refers to compounds in which a cycloalkyl ring  
 15 having a total of 5 to 7 ring atoms is formed on adjacent positions of the phenyl to which they are attached. For example, the compound of formula I in which two  $R_2$  substituents are taken together to form a fused cycloalkyl having 5 carbon atoms at the adjacent 3- and 4-positions and  
 20 having hydrogen as the other  $R_2$  is depicted by the formula below:




It is understood that formula I encompasses the compounds in which the X is attached to the pyridyl moiety  
 25 in the 2-position, the 3-position, and the 4-position. For example, the compound of formula I in which X is attached to the pyridyl moiety in the 3-position is depicted by the formula below:

-19-



The term "pyridyl N-oxide" refers to the compounds of the formula



5       The term "pharmaceutically-acceptable addition salt" refers to an acid addition salt.

      The compound of formula I and the intermediates described herein form pharmaceutically acceptable acid addition salts with a wide variety of organic and inorganic acids and include the physiologically acceptable salts which are often used in pharmaceutical chemistry. Such salts are also part of this invention. A pharmaceutically-acceptable addition salt is formed from a pharmaceutically-acceptable acid as is well known in the art. Such salts include the pharmaceutically acceptable salts listed in Journal of Pharmaceutical Science, 66, 2-19 (1977) which are known to the skilled artisan. Typical inorganic acids used to form such salts include hydrochloric, hydrobromic, hydriodic, nitric, sulfuric, phosphoric, hypophosphoric, metaphosphoric, pyrophosphoric, and the like. Salts derived from organic acids, such as aliphatic mono and dicarboxylic acids, phenyl substituted alkanolic acids, hydroxyalkanoic and hydroxyalkandioic acids, aromatic acids, aliphatic and aromatic sulfonic acids, may also be used. Such pharmaceutically acceptable salts thus include acetate, phenylacetate, trifluoroacetate, acrylate, ascorbate, benzoate, chlorobenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, methylbenzoate, o-acetoxybenzoate,



-20-

naphthalene-2-benzoate, isobutyrate, phenylbutyrate,  $\alpha$ -hydroxybutyrate, butyne-1,4-dicarboxylate, hexyne-1,4-dicarboxylate, caprate, caprylate, cinnamate, citrate, formate, fumarate, glycollate, heptanoate, hippurate, 5 lactate, malate, maleate, hydroxymaleate, malonate, mandelate, mesylate, nicotinate, isonicotinate, nitrate, oxalate, phthalate, teraphthalate, propiolate, propionate, phenylpropionate, salicylate, sebacate, succinate, suberate, benzenesulfonate, p-bromobenzenesulfonate, 10 chlorobenzenesulfonate, ethylsulfonate, 2-hydroxyethylsulfonate, methylsulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, naphthalene-1,5-sulfonate, p-toluenesulfonate, xylenesulfonate, tartarate, and the like.

15 As with any group of pharmaceutically active compounds, some groups are preferred in their end use application. Preferred embodiments of the present invention are given below:

Compounds in which  $R_1$  is selected from the group 20 consisting of  $-C(O)OR_4$ , and  $-SO_2R_5$  are preferred.

When  $R_1$  is  $-SO_2R_5$ , compounds in which  $R_5$  is selected from the group consisting of alkyl and fluorinated alkyl are preferred.

When  $R_1$  is  $-C(O)OR_4$ , compounds in which  $R_4$  is alkyl are 25 preferred.

As potentiators of  $mGluR_2$ , the compounds in which the pyridyl moiety is attached to X at the 3-position and  $R_1$  is  $-SO_2R_5$  are preferred.

As potentiators of  $mGluR_3$ , compounds in which the 30 pyridyl moiety is attached to X at the 4-position and  $R_1$  is  $-C(O)OR_4$  are preferred.


Compounds in which  $R_2$  is from 1 to 3 substituents independently selected from the group consisting of hydrogen, hydroxy, trisubstituted silyloxy, alkyl, 35 substituted alkyl, alkenyl, substituted alkenyl, alkoxy, substituted alkoxy, cycloalkoxy, substituted cycloalkoxy,

-21-

cycloalkyl, substituted cycloalkyl, halogen, cyano, nitro, phenyl, substituted phenyl, pyridyloxy, thiophenoxy, substituted thiophenoxy, phenylsulfinyl, substituted phenylsulfinyl, phenylsulfonyl, substituted phenylsulfonyl, benzoyl, substituted benzoyl, phenoxy, and substituted phenoxy are preferred, with compound in which one  $R_2$  is selected from the group consisting of substituted alkyl, phenyl, substituted phenyl, thiophenoxy, substituted thiophenoxy, phenoxy, and substituted phenoxy being more preferred, and with benzyl, substituted benzyl, phenyl, substituted phenyl, phenoxy and substituted phenoxy being more preferred, and with benzyl, substituted benzyl, phenyl, substituted phenyl, phenoxy and substituted phenoxy being even more preferred.

When one  $R_2$  is substituted alkyl preferred substituents are halogen, alkoxy, phenyl, substituted phenyl having from 1 to 3 substituents selected from the group consisting of hydroxy, halogen, alkoxy, carboxy, alkoxy carbonyl, amido, amino, and pyridyl, thienyl, and phenoxy.

Particularly preferred substituted alkyls are those depicted as the radical of the formula



wherein  $Ru'$  is from selected from the group consisting of hydrogen, hydroxy, lower alkyl, and fluoro;  $Ru''$  is from selected from the group consisting of hydrogen, lower alkyl, alkoxy, and fluoro; provided that when  $Ru'$  is hydroxy then  $Ru''$  is not alkoxy or fluoro, and when  $Ru'$  is fluoro then  $Ru''$  is not alkoxy; and  $Rv$  is from 1 to 3 groups independently selected from the group consisting of hydrogen, alkyl, alkoxy, cyano, nitro, trifluoromethyl, and halogen.

-22-

When one of  $R_2$  is substituted phenyl preferred substituents are from 1 to 3 groups independently selected from the group consisting of hydrogen, trifluoromethyl, trifluoromethoxy, hydroxy, alkyl, alkoxy, halogen, amido, substituted amido, sulfonylamido, cyano, and nitro.

When one of  $R_2$  is substituted thiophenoxy preferred substituents are from 1 to 3 groups independently selected from the group consisting of hydrogen, trifluoromethyl, trifluoromethoxy, alkyl, alkoxy, halogen, amido, cyano, and nitro.

When one of  $R_2$  is substituted phenoxy preferred substituents are from 1 to 3 groups independently selected from the group consisting of hydrogen, hydroxy, alkyl, substituted alkyl having from 1 to 3 groups independently selected from the group consisting of hydroxy, alkyl, cycloalkyl, alkoxy, halogen, amido, substituted amido, sulfonylamido, cyano, nitro, phenyl, and  $-S(O)_p$ alkyl (wherein  $p$  is 0, 1, or 2), cycloalkyl, alkoxy, trifluoromethoxy, halogen, acyloxy, carboxy, amido, substituted amido, amino, acylamino, sulfonylamido, sulfonamide, cyano, nitro, phenyl, substituted phenyl having from 1 to 3 groups independently selected from the group consisting of trifluoromethyl, hydroxy, alkyl, alkoxy, halogen, amido, cyano, and nitro.

Compounds in which one  $R_2$  is sulfonylamido wherein  $R_p$  is selected from the group consisting of alkyl, phenyl, and substituted phenyl having from 1 to 3 groups independently selected from the group consisting of hydrogen, alkyl, alkoxy, halogen, cyano, nitro, and trifluoromethoxy, and  $R_p$  is selected from the group consisting of hydrogen and alkyl are preferred.

Compounds in which at least one of the  $R_2$  substituents is attached in the 3-position are preferred.

Compounds in which at least one of the  $R_2$  substituents is attached in the 4-position are preferred.

Compounds in which  $R_6$  is from 1 to 2 substituents independently selected from the group consisting of

-23-

hydrogen, alkyl, alkoxy, trifluoromethyl, and halogen are preferred.


Compounds in which X is selected from the group consisting of a  $-\text{CH}_2-$ ,  $-\text{CHR}_7-$ , and  $-\text{CH}_2\text{CH}_2-$  are preferred, with compounds in which X is  $-\text{CH}_2-$  being more preferred.

Compounds in which Y is selected from the group consisting of a bond,  $-\text{CH}_2-$ ,  $-\text{CHR}_8-$ ,  $-\text{CH}_2\text{CH}_2-$  are preferred, with compounds in which Y is a bond being more preferred.

Compounds in which X is a  $-\text{CH}_2-$  and Y is bond are even most preferred.

The compounds of formula I are prepared as described in Reaction Scheme A. In Reaction Schemes below all substituents, unless otherwise indicated, are as previously defined, and all reagents are well known and appreciated in the art.

Reaction Scheme A



formula I

Reaction Scheme A, step a, depicts the reductive amination of an appropriate compound of formula (1) and an appropriate amine of formula (2) to give a compound of

-24-

formula (3). An appropriate compound of formula (1) is one in which  $R_6$  is as desired in the final compound of formula I,  $X'$  is a bond or  $-CH_2-$  and  $R$  is hydrogen, giving  $X$  to be  $-CH_2-$  or  $-CH_2CH_2-$ , respectively, as desired for the compound of formula I, or  $X'$  is a bond and  $R$  is  $R_7$  as desired in the final compound of formula I, and the point of attachment of  $X'$  is the point of attachment of  $X$  as desired in the final compound of formula I. Compounds of formula (1) are well known in the art. An appropriate compound of formula (2) is one in which  $R_2$  is as desired in the final product of formula I or gives rise after modification or deprotection to  $R_2$  as desired in the final compound of formula I and  $Y$  is as desired in the final product of formula I. Compounds of formula (2) are known in the art and can be obtained by procedures well known in the art and those described herein. As described herein, compounds of formula (2) in which  $Y$  is a bond can be obtained by reduction of nitro groups. Compounds of formula (2) in which  $Y$  is  $-CH_2-$ ,  $-CH_2CH_2-$ ,  $-CHR_8-$ ,  $-CHR_9CH_2-$ , or  $-CH_2CHR_9-$  can be obtained by reduction of a nitrile, transforming a carboxylic acid to an amine as is well known in the art, for example such transformations can be carried out by reduction of an amide, transformation to the aldehyde followed by reductive amination, or reduction to the alcohol followed by elaboration to the amine (such as by Mitsunobu reaction using phthalimide followed by deprotection). All of these transformations may involve the introduction and removal of appropriate protecting groups as is well known and appreciated in the art.

The reductive amination depicted in Reaction Scheme A, step a, can be carried out under a variety of conditions, such as by hydrogenation using a suitable catalyst or using a suitable reducing agent. For example, an appropriate aldehyde of formula (1) is contacted with an appropriate amine of formula (2) and a suitable reducing agent to give a compound of formula (3). The reaction is carried out in a suitable solvent, such as methanol, ethanol,

-25-

tetrahydrofuran, or mixtures of methanol or ethanol and tetrahydrofuran. The reaction may be carried out in the presence of a drying agent, such as sodium sulfate, cupric sulfate, or molecular sieves. The reaction is carried out in the presence of from 1.0 to 6.0 molar equivalents of a suitable reducing agent, such as, sodium borohydride or sodium cyanoborohydride. It may be advantageous to allow Schiff base formation to proceed before addition of the suitable reducing agent. When sodium cyanoborohydride is used it may be advantageous to monitor and adjust the pH during the course of the reaction as is known in the art. The reaction is generally carried out at temperatures of from 0°C to the refluxing temperature of the solvent. Generally, the reactions require 1 to 72 hours. The product can be isolated and purified by techniques well known in the art, such as filtration, extraction, evaporation, trituration, chromatography, and recrystallization.

Reaction Scheme A, step b, depicts the acylation or sulfonation, using appropriate reagents, of a compound of formula (3) to give a compound of formula (I) or a protected compound of formula I. An appropriate reagent is one that transfers a  $-C(O)R_3$ ,  $-C(O)OR_4$ , or  $-SO_2R_5$  group as required to give the desired  $R_1$  group. In the case of  $-C(O)R_3$  such reagents include acids, acid halides, anhydrides (including mixed anhydrides), activated amides (such as imidazolides), and activated esters. In the case of  $-C(O)OR_4$  such reagents include esters of chloroformate and pyrocarbonates (such as BOC anhydride). In the case of  $-SO_2R_5$  such reagents includes halosulfonates and sulfonate anhydrides with the use of sulfonyl chlorides are preferred. Acylation and sulfonation reactions using such reagents are well known and appreciated in the art.

For example, acylation reactions to transfer  $-C(O)R_3$  involve an amide formation reaction similar to those which are conventionally conducted for peptide synthesis and synthetic methods used therein can also be employed. For example, well known coupling reagents such as carbodiimides

-26-

with or without the use of well known additives such as N-hydroxysuccinimide, 1-hydroxybenzotriazole, etc. can be used to facilitate amide formation. The reaction is conventionally conducted in an inert aprotic solvent such as  
5 pyridine, dimethylformamide, dichloromethane, chloroform, 1,2-dichloroethane, acetonitrile, tetrahydrofuran and the like.

Alternatively, for example an acid halide can be employed in the reaction. It may be advantageous to use a  
10 suitable base to scavenge the acid generated during the reaction. Suitable bases include, by way of example, triethylamine, N,N-diisopropylethylamine, N-methylmorpholine, pyridine, sodium carbonate, potassium carbonate, sodium bicarbonate, and the like. The reaction is  
15 preferably conducted at from about 0°C to about 60°C until reaction completion which typically occurs within 1 to about 24 hours. Upon reaction completion, the product of formula I or protected compound of formula I can be isolated and purified by techniques well known in the art, such as  
20 filtration, extraction, evaporation, trituration, chromatography, and recrystallization.

For example, acylation reactions to transfer  $-C(O)OR_4$  involve a reaction of a chloroformate ester with a compound of formula (3). For example, the reaction is conventionally  
25 conducted in an inert aprotic polar diluent such as dimethylformamide, dichloromethane, chloroform, acetonitrile, 1,2-dichloroethane, pyridine, tetrahydrofuran and the like. The reaction may be carried out in the presence of a suitable base to scavenge the acid generated  
30 during the reaction. Suitable bases include, by way of example, pyridine, triethylamine, N,N-diisopropylethylamine, N-methylmorpholine, and the like. The reaction is preferably conducted at from about 0°C to about 80°C until reaction  
35 completion which typically occurs within 1 to about 24 hours. Upon reaction completion, the product of formula I or protected product of formula I can be isolated and purified by techniques well known in the art, such as

-27-

filtration, extraction, evaporation, trituration, chromatography, and recrystallization.

For example, sulfonation reactions to transfer  $-SO_2R_5$  involve the reaction of halosulfonates or sulfonate anhydrides with compounds of formula (3). Such reactions are carried out in a suitable solvent, such as pyridine, tetrahydrofuran, dioxane, diethyl ether or dimethylformamide, dichloromethane, chloroform, acetonitrile, 1,2-dichloroethane, tetrahydrofuran, and the like, at a temperature from about  $0^\circ\text{C}$  to about ambient temperature, and are carried out in the presence of a suitable base such as pyridine, triethylamine, N,N-diisopropylethylamine, sodium carbonate, potassium carbonate, sodium bicarbonate, potassium bicarbonate, and the like. The resultant sulfonamide is isolated by methods well known in the art, such as filtration, extraction, evaporation, trituration, chromatography, and recrystallization.

Reaction Scheme A, optional step c, not shown, a protected compound of formula I is deprotected to give a compound of formula I. Such deprotection reactions are well known and appreciated in the art. Protecting Groups in Organic Synthesis, Theodora Greene (Wiley-Interscience).

Reaction Scheme A, optional step d, not shown, an acid addition salt is formed using a pharmaceutically acceptable acid. The formation of acid addition salts is well known and appreciated in the art.


A variety of compounds of formula I and related compounds (such as compounds of formula (3) above) can be used as starting materials which upon further elaboration give rise to compounds of formula I. Some of these transformations are trivial, such as hydrolysis of a nitrile to an amide or acid, hydrolysis of an ester to an acid, formation of an amide from an acid or an ester, alkylation and acylation of hydroxy groups, oxidation, reduction, deprotection, displacements, and the like. A number other transformations, while readily apparent to the skilled



-28-

artisan, are described in detail in the Reaction Schemes below.

## Reaction Scheme B



Reaction Scheme B, step a, depicts a coupling reaction of an appropriate compound of formula (4) and an appropriate aryl compound of formula (5). An appropriate compound of formula (4) is one in which X, Y, R<sub>6</sub>, R<sub>2</sub> is as defined for the compound of formula I, except one of R<sub>2</sub> is a bromo or iodo or triflate or -CH<sub>2</sub>Br, and R<sub>1</sub> is a hydrogen or R<sub>1</sub> as desired in the final compound of formula I. Compounds of formula (4) are readily prepared as described in Reaction Scheme A. An appropriate aryl compound of formula (5) is one in which Ar is one of the R<sub>2</sub> groups as desired in the final product of formula I, such as phenyl, substituted phenyl, pyridyl, substituted pyridyl, thienyl, or substituted thienyl. It is appreciated in the art that boronic esters are also suitable for use in such coupling reactions. The preparation and use of such arylboronic acids (and boronic esters) is well known and appreciated in the art. W. J. Thompson and J Gaudino, J. Org. Chem., 49, 5237-5243 (1984). Arylboronic acids are frequently contaminated with their corresponding anhydrides which do not perform well in the Suzuki coupling. Material contaminated by detrimental amounts of anhydride can be converted to the corresponding acid by hydrolysis. The hydrolysis is performed, if required, by briefly boiling in water.

As will be appreciated by the skilled person, Reaction Scheme B, step a, depicts a Suzuki coupling. See for

-29-

example, A. Suzuki, Pure Appl. Chem., 66, 213-222 (1994); A. Suzuki, Pure Appl. Chem., 63, 419-422 (1991), N. Miyaura et al., J. Org. Chem., 51, 5467-5471 (1986); Y. Hoshino et al., Bull. Chem. Soc. Japan, 61, 3008-3010 (1988); N. Miyaura et al., J. Am. Chem. Soc., 111, 314-321 (1989); W. J. Thompson et al., J. Org. Chem., 53, 2052-2055 (1988); and T. I. Wallow and B. M. Novak, J. Org. Chem., 59, 5034-5037 (1994). Other coupling reactions, such as the Stille reaction, can be used.


10 It is appreciated that the coupling gives a compound of formula I when, the compound of formula (4),  $R_1$ , is  $R_1$  as desired in the final product and when  $R_1$  is hydrogen the product will required acylation or sulfonation, as depicted in Reaction Scheme A, step b, to give a compound of formula  
15 I.

For example, an appropriate compound of formula (4) and an appropriate aryl compound of formula (5) can be coupled under Suzuki conditions to give a compound of formula I or a compound that gives rise to a compound of formula I. The  
20 Suzuki coupling reaction is performed in a suitable solvent, such as toluene, dichloromethane, or tetrahydrofuran. The reaction is performed using from about 1.1 to about 3 molar equivalents of an appropriate arylboronic acid or ester. The reaction is carried out in the presence of from about 1 to  
25 about 3 molar equivalents of a suitable base, such as potassium carbonate or sodium carbonate. The coupling is performed using a suitable palladium catalyst, such as tetrakis(triphenylphosphine) palladium (0), bis(acetonitrile)palladium (II) chloride, palladium (II) chloride, palladium (II) acetoacetate, and  
30 tris(dibenzylideneacetone)dipalladium(0). The suitable palladium catalyst chosen may be modified by the use of ligands, such as tri(fur-2-yl)phosphine and tri(o-tolyl)phosphine: V. Farina and B. Krishnan, J. Am. Chem. Soc., 113, 9586-9595 (1991). The coupling is performed at a  
35 temperature ranging from 0°C to the refluxing temperature of the solvent. The Suzuki coupling reactions depicted in

-30-

Reaction Scheme B generally require from 6 hours to 14 days. The product of the coupling reaction can be isolated and purified using techniques well known in the art. These techniques include extraction, evaporation, chromatography and recrystallization.

## Reaction Scheme C.1



Reaction Scheme C.1, step a, depicts a coupling reaction of an appropriate compound of formula (4) and an appropriate phenol compound of formula (6). An appropriate compound of formula (4) is as described above in Reaction Scheme B, except that R<sub>2</sub> is not -CH<sub>2</sub>Br. An appropriate phenol compound of formula (6) is one in which R<sub>c</sub> is as desired in the final product of formula I or gives rise upon modification to the desired final product of formula I.


As will be appreciated by the skilled person, Reaction Scheme C, step a, depicts an Ullmann coupling. See for example, F. D. Rychnozsky and K. Hwang, J. Org. Chem., 59, 5414 (1994). It is appreciated that the coupling gives a compound of formula I when R<sub>1'</sub> is R<sub>1</sub> as desired in the final product and when R<sub>1'</sub> is hydrogen the product will require acylation or sulfonation, as depicted in Reaction Scheme A, step b, to give a compound of formula I.

For example, an appropriate compound of formula (4) and an appropriate phenol compound of formula (6) are coupled give a compound of formula I or a compound that gives rise to a compound of formula I. The Ullmann coupling reaction is performed in a suitable solvent, such as pyridine. The

-31-

reaction is performed using from about 1.1 to about 3 molar equivalents of an appropriate phenol. The reaction is carried out in the presence of from about 1 to about 3 molar equivalents of a suitable base, such as potassium carbonate or sodium carbonate. The coupling is performed using a suitable catalyst, such as copper salts. These Ullmann coupling reactions depicted in Reaction Scheme C generally require from 6 hours to 24 hours. The product of the coupling reaction can be isolated and purified using techniques well known in the art. These techniques include extraction, evaporation, chromatography and recrystallization.

Reaction Scheme C.2



Reaction Scheme C.2, step a, depicts a coupling reaction of an appropriate compound of formula (4') and an appropriate compound of formula (6'). An appropriate compound of formula (4') is one in which X, Y,  $R_6$ ,  $R_2$  is as defined for the compound of formula I, except one of  $R_2$  is a hydroxy, and  $R_{1'}$  is a hydrogen or  $R_1$  as desired in the final compound of formula I. An appropriate compound of formula (6') is one in which  $R_c$  is one or more electron withdrawing group as desired in the final product of formula I or gives rise upon modification to the desired final product of formula I.


It is appreciated that the coupling gives a compound of formula I when  $R_{1'}$  is  $R_1$  as desired in the final product and when  $R_{1'}$  is hydrogen the product will required acylation or sulfonation, as depicted in Reaction Scheme A, step b, to

-32-

give a compound of formula I. For the reaction depicted in Scheme C.2 it is preferred that the compound of formula (4') is one in which R<sub>1</sub>' is hydrogen.

For example, an appropriate compound of formula (4') and an appropriate compound of formula (6') are coupled give a compound of formula I or a compound that gives rise to a compound of formula I. The reaction is performed in a suitable solvent, such as dimethylformamide, dimethylacetamide, and dimethyl sulfoxide. The reaction is performed using from about 1.1 to about 3 molar equivalents of an appropriate phenol. The reaction is carried out in the presence of from about 1 to about 3 molar equivalents of a suitable base, such as potassium hydride, sodium hydroxide, or sodium hydride. The coupling is performed using a suitable catalyst, such as copper salts. The reaction generally require from 6 hours to 48 hours. The product of the coupling reaction can be isolated and purified using techniques well known in the art. These techniques include extraction, evaporation, chromatography and recrystallization.

Reaction Scheme D



Reaction Scheme D, step a, depicts a reaction of an appropriate compound of formula (7) with appropriate organophosphorous ylid in a Wittig-type reaction to give a compound of formula I. An appropriate compound of formula (7) is one in which X, Y, R<sub>1</sub>, R<sub>2</sub>, and R<sub>6</sub>, are as defined for the compound of formula I. As will be appreciated by the skilled person, in addition to the aldehyde depicted in Scheme D, a variety of carbonyl compounds can be used in the reaction of Reaction Scheme D, step a, depending of the


-33-

substitution desired in the final compound of formula I. An appropriate organophosphorous ylid is one which converts the carbonyl of a compound of formula (7) to an alkenyl or substituted alkenyl as desired in the final product of formula I. Appropriate organophosphorous reagents, such as Wittig, Wittig-Horner, Horner-Emmons-Wadsworth reagents, and the use of appropriate organophosphorous reagents is well known and appreciated in the art.

For example, an appropriate compound of formula (7) is reacted with appropriate organophosphorous ylid to give a compound of formula I. An appropriate organophosphorous ylid is formed by contacting an appropriate organophosphorous reagent with a suitable base, such as lithium diisopropylamide, sodium hydride, lithium bis(trimethylsilyl)amide or potassium t-butoxide. The ylid formation is carried out in a suitable solvent, such as tetrahydrofuran, benzene, or diethyl ether. The ylid formation is generally carried out at a temperature of from -78°C to ambient temperature. An appropriate organophosphorous ylid is contacted with an appropriate compound of formula (7). The reaction is carried out in a suitable solvent, such as tetrahydrofuran, benzene, or diethyl ether. Generally, the reaction is carried out in the same solvent and temperatures used to form the appropriate organophosphorous ylid. The reaction generally requires from 1 hour to 48 hours. The product can be isolated by techniques well known in the art, such as extraction and evaporation. The product can then be purified by techniques well known in the art, such as distillation, chromatography, or recrystallization.

-34-

## Reaction Scheme E




Reaction Scheme E, step a, depicts a reaction of an appropriate compound of formula (7) with appropriate organometallic reagent, such as a Grignard reagent to give a compound of formula I. An appropriate compound of formula (7) is as defined in Reaction Scheme D. An appropriate organometallic reagent is one which converts the carbonyl of a compound of formula (7) to a hydroxy substituted group as desired in the final product of formula I. Appropriate organometallic reagents and their use is well known and appreciated in the art.

For example, an appropriate compound of formula (7) is reacted with appropriate Grignard reagent to give a compound of formula I. Generally, 1 to 2 equivalents of Grignard reagent are used. The reaction is carried out in a suitable solvent, such as tetrahydrofuran, diethyl ether. The reaction is generally carried out at temperatures of from -40°C to ambient temperature. The reaction generally requires from 1 hour to 48 hours. The product can be isolated by techniques well known in the art, such as extraction and evaporation. The product can then be purified by techniques well known in the art, such as distillation, chromatography, or recrystallization.

As will be appreciated by the skilled person, the hydroxy compound thus obtained can undergo a variety of further reactions, such as halogenation, acylation, alkylation, reduction, oxidation, elimination, and the like to give further compounds of formula I.

-35-

In addition, the compounds of formula I in which  $R_1$  is  $-S(O)_2R_5$ ,  $X$  is  $-CH_2-$ , and  $Y$  is a bond can be prepared as described in Reaction Scheme F.



Reaction Scheme F, step a, depicts coupling of an appropriate compound of formula (8) with an appropriate compound of formula (9) to give a compound of formula I. An appropriate compound of formula (8) is one in which  $R_2$  is as defined for the compound of formula I and  $R_1$  is  $-S(O)_2R_5$ . An appropriate compound of formula (9) is one in which  $R_6$  is as defined for the compound of formula I. Compounds of formula (8) and (9) are readily prepared by methods known in the art and as described herein.

For example, where the appropriate compound of formula (9) is one substituted with  $-CH_2Br$  or, preferably  $-CH_2Cl$ , a compound of formula (8) is treated with a suitable base, such as sodium hydride, potassium hydride, potassium *t*-butoxide, potassium carbonate, and cesium carbonate.

Generally, from 2 to about 5 molar equivalents of base are used and additional base may be added if the compound of formula (9) is used in the form of a salt. Generally, the reaction is carried out in a suitable solvent, such as tetrahydrofuran, diethyl ether, and dimethylformamide. The reaction is generally carried out at temperatures of from  $-10^\circ\text{C}$  to  $80^\circ\text{C}$ . The reaction generally requires from 1 hour to 48 hours. The product can be isolated by techniques well known in the art, such as extraction and evaporation. The product can then be purified by techniques well known in the



-36-

art, such as distillation, chromatography, or recrystallization.

For example, where the appropriate compound of formula (9) is substituted with  $-\text{CH}_2\text{OH}$  the coupling is carried out under Mitsunobu conditions which are well known in the art. The reaction is carried out in a suitable solvent, such as tetrahydrofuran and diethyl ether using a phosphine, such as triphenylphosphine or a resin bound phosphine and a dialkyl azodicarboxylate, such as diethyl azodicarboxylate, di-iso-propyl azodicarboxylate or di-t-butyl azodicarboxylate. The reaction is generally carried out at temperatures of from ambient temperatures to  $60^\circ\text{C}$ . The reaction generally requires from 1 hour to 12 hours. The product can be isolated by techniques well known in the art, such as extraction and evaporation. The product can then be purified by techniques well known in the art, such as distillation, chromatography, or recrystallization.

The present invention is further illustrated by the following examples and preparations. These examples and preparations are illustrative only and are not intended to limit the invention in any way.

The terms used in the examples and preparations have their normal meanings unless otherwise designated. For example, " $^\circ\text{C}$ " refers to degrees Celsius; "N" refers to normal or normality; "M" refers to molar or molarity; "mol" refers to mole or moles; "mmol" refers to millimole or millimoles; " $\mu\text{mol}$ " refers to micromole or micromoles; "kg" refers to kilogram or kilograms; "g" refers to gram or grams; " $\mu\text{g}$ " refers to microgram or micrograms; "mg" refers to milligram or milligrams; " $\mu\text{L}$ " refers to microliter or microliters; "mL" refers milliliter or milliliters; "L" refers to liter or liters; "bp" refers to boiling point; "mp" refers to melting point; "brine" refers to a saturated aqueous sodium chloride solution; "h" refers to hour or hours; "min" refers to minute or minutes; "MS" refers to mass spectrometry; "NMR" refers to nuclear magnetic resonance spectroscopy; "TFA" refers to trifluoroacetic

-37-

acid; "CH<sub>2</sub>Cl<sub>2</sub>" or "DCM" refers to dichloromethane; "DCE" refers to dichloroethane; "MeOH" refers to methanol; "NH<sub>4</sub>OH" refers to a concentrated aqueous ammonia solution; "KOH" refers to potassium hydroxide; "H<sub>2</sub>O" refers to water; "HCl" refers to hydrogen chloride; etc. Chemical shifts are given in  $\delta$  and NMR spectra were obtained in CDCl<sub>3</sub>, unless otherwise indicated.

#### Preparation 1

##### 10 3-(2-Methoxyphenoxy)aniline

A solution of 3-fluoronitrobenzene (3.53g, 25mmol) and 2-methoxyphenol (3.72g, 30mmol) in anhydrous N,N-dimethylformamide (25mL) under nitrogen was treated with anhydrous potassium carbonate (7.6g, 55mmol) and heated to 15 110°C for 65h, then cooled to room temperature. The solution was diluted with dichloromethane (50mL) and treated with DOWEX<sup>®</sup> 550A OH anion exchange resin (10g) which had been washed dry with methanol. Stirring was continued for one hour, the mixture was filtered, and the filtrate 20 concentrated *in vacuo* (exhaustively at end to remove DMF) to give 3-(2-methoxyphenoxy)nitrobenzene as a residue.

The residue was dissolved in 2:1 hexane/dichloromethane and passed through a short column of silica gel (eluted with 1:1 hexane/dichloromethane). The concentrated product was 25 recrystallized from ethyl acetate/hexane (two crops) then placed in a Parr<sup>®</sup> bottle under nitrogen with methanol (50mL), carefully treated with Raney<sup>®</sup> nickel (one tsp.), placed on a Parr<sup>®</sup> apparatus, and hydrogenated at 40-50 psi pressure for 18h. The mixture was carefully filtered 30 through celite<sup>®</sup> under nitrogen without allowing the filter cake to dry, and the filtrate was concentrated *in vacuo* to afford the title compound (2.79g, 52%) as a colorless viscous oil. <sup>1</sup>H NMR: 7.05-7.15 (m,1H), 7.00-7.05 (m,1H), 6.95-7.00 (m,2H), 6.85-6.95 (m,1H), 6.30-6.35 (m,2H), 6.25 35 (m,1H), 3.81 (s,3H), 3.60 (br s,2H).

#### Preparation 2

-38-

4-(2-Benzyloxyphenoxy)aniline


A solution of 2-benzyloxyphenol (6.01g, 30mmol) in anhydrous N,N-dimethylformamide (25mL) was treated with anhydrous potassium carbonate (7.6g, 55mmol), then with 4-fluoronitrobenzene (3.53g, 25mmol), and heated to 90°C for 5h under nitrogen. The solution was cooled to room temperature, diluted with dichloromethane (50mL), and treated with DOWEX® 550A OH anion exchange resin (10g) which had been washed dry with methanol. The mixture was stirred for 3h, filtered, and the filtrate concentrated *in vacuo* (exhaustively at end to remove DMF). The residue was chromatographed on silica gel (eluted with 2:1 hexane/ethyl acetate) to afford 4-(2-benzyloxyphenoxy)nitrobenzene (6.40g, 66%) as a white solid.

4-(2-Benzyloxyphenoxy)nitrobenzene (2.57g, 8mmol) was dissolved in reagent grade ethanol (20mL) and treated with concentrated aqueous hydrochloric acid (2.5mL). The solution was heated to 55°C under nitrogen, then cautiously treated with iron powder (3.1g, 55mmol) in portions with cooling needed after initial exotherm to 75°C. After addition was complete, stirring was continued at 50-60°C for 45 minutes, then the mixture was cooled to room temperature and filtered through celite® (rinse filter cake with dichloromethane). The filtrate was concentrated *in vacuo*, the residue dissolved in dichloromethane and the solution washed with saturated aqueous sodium carbonate. The organic layer was separated and the aqueous layer extracted with dichloromethane. The combined organic solution was dried (sodium sulfate), concentrated *in vacuo*, and chromatographed on silica gel (eluted with dichloromethane, then 10% ethyl acetate/dichloromethane, then 20% ethyl acetate/dichloromethane) to afford the title compound (2.22g, 95%) as a tan solid; mp 111-112.5°C. <sup>1</sup>H NMR: 7.30 (m,4H), 7.20-7.30 (m,1H), 6.97 (m,2H), 6.80-6.90 (m,4H), 6.60-6.65 (m,2H), 5.12 (s,2H), 3.50 (br s, 2H).

-39-

Example 1

N-(3-(2-Methoxyphenoxy)phenyl)-N-(2,2,2-  
trifluoroethylsulfonyl)pyrid-3-ylmethylamine



5        A solution of 3-(2-methoxyphenoxy)aniline (269mg, 1.25mmol) and pyridine-3-carboxaldehyde (107mg, 1.00mmol) was refluxed in methanol (2.5mL) for 2h, then cooled on an ice bath to 5°C. Sodium borohydride (95mg, 2.5mmol) was added and the mixture was warmed to room temperature,  
10       stirred for 2h, concentrated *in vacuo*, and taken up in dichloromethane. The resultant suspension was filtered, the filtrate concentrated *in vacuo*, and the residue chromatographed on silica gel (eluted with 2% methanol/dichloromethane) to give N-(3-(2-  
15       methoxyphenoxy)phenyl)pyrid-3-ylmethylamine.


      The resultant N-(3-(2-methoxyphenoxy)phenyl)pyrid-3-ylmethylamine was dissolved in 2:1 1,2-dichloroethane/pyridine (6mL) under nitrogen, cooled (5°C), and treated with 2,2,2-trifluoroethylsulfonyl chloride  
20       (300mg, 1.64mmol). After one hour at 5°C, the mixture was allowed to warm to room temperature and stirred for 18h. The product mixture was treated with anhydrous potassium carbonate (500mg), stirred 30 min, filtered, and the filtrate concentrated *in vacuo*. The residue was  
25       chromatographed on silica gel (eluted with 2:1 dichloromethane/ethyl acetate) to afford the title compound (226mg, 50%) as a viscous pale amber oil. The hydrochloride salt was obtained by treatment of this material in methanol with ethereal HCl. <sup>1</sup>H NMR: 8.49 (d, 1H, J=4Hz), 8.29 (s, 1H),  
30       7.60-7.65 (m, 1H), 7.20-7.25 (m, 2H), 7.10-7.20 (m, 1H), 6.97 (d, 1H, J=8Hz), 6.90 (m, 2H), 6.80-6.85 (m, 2H), 6.76 (m, 1H),

-40-

4.82 (s,2H), 3.70-3.85 (m,2H), 3.75 (s,3H). MS calcd. 452.4; MS (M+1) 453.3

### Example 2

5 N-(3-(2-Benzoyloxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylamine




The title compound was prepared using a method similar to the method of Example 1 using 3-(2-benzoyloxyphenoxy)aniline. <sup>1</sup>H NMR: 8.45 (m,1H), 8.28 (s,1H), 7.77 (d,1H,J=8Hz), 7.20-7.25 (m,5H), 7.10-7.15 (m,2H), 7.00-7.10 (m,3H), 6.97 (t,1H,J=8Hz), 6.85-6.90 (m,2H), 6.77 (m,1H), 4.99 (s,2H), 4.79 (s,2H), 3.55-3.65 (m,2H). MS calcd. 528.6; MS (M+1) 529.4

15

### Example 3

N-(4-Phenoxy)phenyl)-N-(ethylsulfonyl)-2-(pyrid-3-yl)ethylamine



20 The title compound was prepared using a method similar to the method of Example 1 using (4-phenoxy)aniline, 2-(pyrid-3-yl)acetaldehyde, and ethylsulfonyl chloride. <sup>1</sup>H NMR: 8.45 (m,1H), 8.36 (s,1H), 7.50 (m,1H), 7.35-7.40 (m,2H), 7.15-7.25 (m,3H), 7.10-7.15 (m,1H), 7.03 (d,2H,J=8Hz), 6.97 (d,2H,J=8Hz), 3.91 (t,2H,J=7Hz), 2.95


25

-41-

(q, 2H, J=7Hz), 2.83 (t, 2H, J=7Hz), 1.31 (t, 3H, J=7Hz). MS calcd. 382.5; MS (M+1) 383.4

#### Example 4

5 N-((4-Phenoxy)phenylethyl)-N-(ethylsulfonyl)pyrid-3-ylmethylamine



A solution of pyridine-3-carboxaldehyde (322mg, 3.0mmol) and 4-phenoxyphenethylamine (747mg, 3.5mmol) in reagent grade methanol (5mL) under nitrogen was refluxed for 2h, then cooled to 5°C (ice bath), and diluted with more methanol (5mL). Sodium borohydride (0.23g, 6mmol) was added, and the mixture was slowly allowed to reach room temperature and stirred at room temperature overnight (18h). The mixture was concentrated *in vacuo*, taken up in dichloromethane, and filtered. The filtrate was concentrated *in vacuo* and chromatographed on silica gel (eluted with 7% methanol/dichloromethane) to afford N-((4-phenoxy)phenylethyl)pyridine-3-methylamine (863mg, 94%).

A cooled (5°C) solution of N-((4-phenoxy)phenylethyl)pyridine-3-methylamine (304mg, 1.0mmol) in anhydrous 1,2-dichloroethane (4mL) with anhydrous potassium carbonate (0.50g, 3.6mmol) under nitrogen was treated dropwise with ethylsulfonyl chloride (257mg, 2.0mmol), then slowly warmed to room temperature and stirred at room temperature for 44h. The mixture was treated with tris(2-aminoethyl)amine resin (4 meq/g, 1.0g), stirred 1h, treated with DOWEX<sup>®</sup> 550A OH anion exchange resin (1.5g), stirred 30 min, then filtered and the filter cake was rinsed with dichloromethane. The filtrate was concentrated *in vacuo* and the residue chromatographed on silica gel (eluted


-42-

with 4:1 ethyl acetate/dichloromethane) to afford the title compound (100mg, 25%) as a pale yellow oil. The title compound was converted to the hydrochloride salt by treatment of this material in methanol with ethereal HCl. <sup>1</sup>H NMR: 8.55 (m, 2H), 7.78 (m, 1H), 7.25-7.35 (m, 3H), 7.00-7.10 (m, 3H), 6.94 (d, 2H, J=8Hz), 6.89 (d, 2H, J=9Hz), 4.43 (s, 2H), 3.38 (t, 2H, J=8Hz), 2.88 (q, 2H, J=7Hz), 2.74 (t, 2H, J=8Hz), 1.31 (t, 3H, J=7Hz). MS calcd. 396.5; MS(M+1) 397.5

10

Example 5

N-(4-Phenoxy)phenylmethyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline




The title compound was prepared using a method similar to the method of Example 4 using 4-phenoxyphenylmethylaniline. <sup>1</sup>H NMR: 8.50 (m, 1H), 8.40 (s, 1H), 7.66 (d, 1H, J=8Hz), 7.30-7.35 (m, 2H), 7.20-7.25 (m, 1H), 7.17 (d, 2H, J=8Hz), 7.10 (m, 1H), 6.95 (d, 2H, J=8Hz), 6.90 (d, 2H, J=8Hz), 4.36 (s, 2H), 4.30 (s, 2H), 2.99 (q, 2H, J=7Hz), 1.35 (t, 3H, J=7Hz). MS calcd. 382.5; MS (M+1) 383.4

20

Example 6

N-(4-(Phenoxy)phenyl)-N-(iso-propylsulfonyl)pyrid-3-ylmethylaniline



25

The title compound was prepared using a method similar to the method of Example 1 using (4-phenoxy)aniline and isopropylsulfonyl chloride. <sup>1</sup>H NMR: 8.47 (m, 1H), 8.32 (m, 1H),


-43-

7.70 (m,1H), 7.30-7.35 (m,2H), 7.20-7.25 (m,1H), 7.10-7.15 (m,3H), 6.97 (d,2H,J=8Hz), 6.85 (d,2H,J=8Hz), 4.84 (s,2H), 3.20-3.30 (m,1H), 1.42 (d,6H,J=7Hz). MS calcd. 382.5; MS (M+1) 383.4.

5

Example 7


N-(2-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethanamine



10 The title compound was prepared using a method similar to the method of Example 1 using (2-phenoxy)aniline. <sup>1</sup>H NMR: 8.50 (m,1H), 8.31 (m,1H), 7.75 (m,1H), 7.40 (t, 2H,, J=8Hz), 7.15-7.25 (m,3H), 6.95-7.05 (m,3H), 6.95 (t,1H,J=8Hz), 6.83 (d,2H,J=8Hz), 4.84 (s,2H), 3.90-4.00 (m,2H). MS calcd. 15 422.4; MS (M+1) 423.4.

Example 8

N-(4-(Phenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethanamine



20

The title compound was prepared using a method similar to the method of Example 1 using (4-phenoxy)aniline and ethylsulfonyl chloride. <sup>1</sup>H NMR: 8.50 (m,1H), 8.37 (s,1H), 7.70 (d,1H,J=8Hz), 7.30-7.35 (m,2H), 7.20-7.25 (m,1H), 7.10- 25 7.15 (m,3H), 6.98 (d,2H,J=8Hz), 6.87 (d,2H,J=9Hz), 4.83




-44-

(s,2H), 3.09 (q,2H,J=7Hz), 1.42 (t,3H,J=7Hz). MS calcd. 368.5; MS (M+1) 369.3

Example 9


5 N-(4-(2-Methoxyphenoxy)-3-chlorophenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylamine



The title compound was prepared using a method similar to the method of Example 1 using 4-(2-methoxyphenoxy)-3-chloroaniline. The hydrochloride salt was obtained by treatment of this material in methanol with ethereal HCl. <sup>1</sup>H NMR: 8.51 (m,1H), 8.32 (s,1H), 7.66 (m,1H), 7.25-7.30 (m,2H), 7.15-7.20 (m,1H), 6.85-7.00 (m,4H), 6.55 (d,1H,J=8.5Hz), 4.82 (s,2H), 3.75-3.85 (m,2H), 3.76 (s,3H). MS calcd. 486.9; MS (M+1) 487.6

Example 10

20 N-(4-(2,6-Dimethoxyphenoxy)-3-chlorophenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylamine



The title compound was prepared using a method similar to the method of Example 1 using 4-(2,6-dimethoxyphenoxy)-3-


-45-

chloroaniline.  $^1\text{H}$  NMR: 8.51 (m,1H), 8.32 (s,1H), 7.66 (m,1H), 7.25-7.30 (m,2H), 7.15-7.20 (m,1H), 6.85-7.00 (m,4H), 6.55 (d,1H, $J=8.5\text{Hz}$ ), 4.82 (s,2H), 3.75-3.85 (m,2H), 3.76 (s,3H). MS calcd. 486.9; MS (M+1) 487.6

5

Example 11

N-(4-(2-Benzyloxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylamine




10

The title compound was prepared using a method similar to the method of Example 1 using 4-(2-benzyloxyphenoxy)aniline.  $^1\text{H}$  NMR: 8.48 (m,1H), 8.33 (s,1H), 7.60-7.65 (m,1H), 7.15-7.25 (m,5H), 7.05-7.15 (m,5H), 7.03 (t,1H, $J=8\text{Hz}$ ), 6.94 (t,1H, $J=8\text{Hz}$ ), 6.84 (d,2H, $J=9\text{Hz}$ ), 5.02 (s,2H), 4.82 (s,2H), 3.70-3.80 (m,2H). MS calcd. 528.6; MS (M+1) 529.4

15

Example 12

N-(4-(2-Methoxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylamine



20


The title compound was prepared using a method similar to the method of Example 1 using 4-(2-methoxyphenoxy)aniline

-46-

and ethylsulfonyl chloride.  $^1\text{H}$  NMR:  $\delta$  8.46 (d, 1H,  $J=4.8\text{Hz}$ ), 8.33 (s, 1H), 7.67-7.69 (d, 1H,  $J=7.7\text{Hz}$ ), 7.19-7.24 (m, 1H), 7.10-7.16 (m, 1H), 7.06-7.10 (d, 2H,  $J=7.0$ ), 6.94-6.99 (d, 2H,  $J=8.1\text{Hz}$ ), 6.89-6.92 (d, 1H,  $J=7.7\text{Hz}$ ), 6.78-6.82 (d, 2H,  $J=8.8\text{Hz}$ ), 4.81 (s, 2H), 3.76 (s, 3H), 3.02-3.10 (dd, 2H,  $J=14.7$ , 7.3Hz), 1.4 (t, 3H,  $J=7.7\text{Hz}$ ). MS calcd. 398.5; MS (M+1) 399.2.

Example 13

10 N-(4-(2-Methoxyphenoxy)phenyl)-N-(cyclopropylsulfonyl)pyrid-3-ylmethanamine




The title compound was prepared using a method similar to the method of Example 1 using 4-(2-methoxyphenoxy)aniline and cyclopropylsulfonyl chloride.  $^1\text{H}$  NMR: 8.49 (d, 1H), 8.34 (s, 1H), 7.62-7.68 (m, 1H), 7.17-7.23 (m, 2H), 7.10-7.18 (m, 3H), 6.85-6.96 (m, 3H), 6.78 (m, 2H), 4.81 (s, 2H), 3.76 (s, 3H), 2.42 (m, 1H), 1.10 (m, 2H), 0.96 (m, 2H). MS calcd. 410.5; MS (M+1) 411.4.

-47-

Example 14

N-(4-(2-Methoxyphenoxy)-3-chlorophenyl)-N-  
ethylsulfonyl)pyrid-3-ylmethylamine




5

The title compound was prepared using a method similar to the method of Example 1 using 4-(2-methoxyphenoxy)-3-chloroaniline and ethylsulfonyl chloride. <sup>1</sup>H NMR: 8.48 (d, 1H, J=4.8Hz), 8.35 (s, 1H), 7.64-7.73 (d, 1H, J=8.0Hz), 7.28-7.33 (d, 1H, J=2.5Hz), 7.21-7.27 (m, 2H), 7.10-7.20 (m, 1H), 6.87-7.00 (m, 3H), 6.51-6.62 (d, 1H, J=8.8Hz), 4.81 (s, 2H), 3.76 (s, 3H), 3.01-3.15 (dd, 2H, J=14.7, 7.3Hz), 1.41 (t, 3H, J=7.3Hz). MS calcd. 432.9; MS (M+1) 433.6

15

Example 15

N-(4-(2-Methoxyphenoxy)-3-chlorophenyl)-N-  
cyclopropylsulfonyl)pyrid-3-ylmethylamine



20

The title compound was prepared using a method similar to the method of Example 1 using 4-(2-methoxyphenoxy)-3-chloroaniline and cyclopropylsulfonyl chloride. <sup>1</sup>H NMR: 8.55 (d, 1H, J=8.1Hz), 8.35 (s, 1H), 8.03 (s, 1H), 7.88 (s, 1H), 7.65 (s, 1H), 7.12-7.29 (m, 3H), 6.89-7.02 (m, 2H), 6.49-6.58


-48-

(d,1H, J=19.4Hz), 5.06 (s, 2H), 3.67 (s,3H), 2.87 (m,1H), 1.01 (d, 2H, J=5.5Hz), 0.84 (m, 2H). MS calcd. 444.9; MS (M+1) 445.5

5

Example 16

N-(4-(2-Methoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylamine



10


The title compound was prepared using a method similar to the method of Example 1 using 4-(2-methoxyphenoxy) aniline. The hydrochloride salt was obtained by treatment of this material in methanol with ethereal HCl. <sup>1</sup>H NMR: 8.49 (d,1H), 8.29 (s,1H), 7.62-7.68 (m,1H), 7.20-7.28 (m,2H), 7.10-7.20 (m,1H), 7.03 (d,1H), 6.96 (m,2H), 6.85-6.90 (m,1H), 6.80 (m,2H), 4.82 (s,2H), 3.75 (s,3H), 1.6 (s,2H). MS calcd. 452.5; MS (M+1) 453.3

15

Example 17

20

N-(4-(2-Ethoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylamine



25

The title compound was prepared using a method similar to the method of Example 1 using 4-(2-ethoxyphenoxy) aniline. <sup>1</sup>H NMR: 8.54 (d,2H, J=22Hz), 8.03 (d,1H, J=7.7Hz), 7.63 (m,1H), 7.20-7.25 (d,2H, J=6.8Hz), 7.10-7.20 (m,2H),


-49-

7.00 (d,1H, J=6.6Hz), 6.92 (t,1H, J=7.7Hz), 6.70-6.80 (d,2H, J=8.8Hz), 4.94 (s,2H), 4.64-4.81 (dd,2H, J=20.2,9.9Hz), 3.84-3.97 (dd,2H, J=13.9,7.0Hz), 0.99 (t,3H, J=7.0Hz). MS calcd. 466.5; MS (M+1) 467.8

5

Example 18

N-(4-(2-Ethoxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline




10 The title compound was prepared using a method similar to the method of Example 1 using 4-(2-ethoxyphenoxy) aniline and ethylsulfonyl chloride. <sup>1</sup>H NMR: 8.61 (d,2H, J=13.2Hz), 8.13 (d,1H, J=8.1Hz), 7.70-7.75 (m,1H), 7.27 (d,2H, J=8.8Hz), 7.08-7.20 (m,2H), 7.00 (d,1H, J=6.2Hz), 6.92 (t,1H, J=7.7Hz), 6.72-6.76 (d,2H, J=8.8Hz), 4.97 (s,2H), 3.85-3.95 (dd,2H, J=13.9,6.7), 3.16-3.30 (dd,2H, J=14.7,7.3Hz), 1.25 (t,3H, J=7.3Hz), 0.98 (t,3H, J=7.0Hz). MS calcd. 412.4; MS (M+1) 413.6

20

Example 19

N-(4-(2-Ethoxyphenoxy)phenyl)-N-(cyclopropylsulfonyl)pyrid-3-ylmethylaniline




25 The title compound was prepared using a method similar to the method of Example 1 using 4-(2-ethoxyphenoxy) aniline

-50-

and cyclopropylsulfonyl chloride.  $^1\text{H}$  NMR: 7.88 (d, 1H,  $J=7.7\text{Hz}$ ), 7.26 (d, 2H,  $J=8.8\text{Hz}$ ), 7.5-7.18 (m, 2H), 6.97-7.05 (d, 1H,  $J=7.7\text{Hz}$ ), 6.87-6.96 (m, 1H), 6.67-6.77 (d, 2H,  $J=8.8\text{Hz}$ ), 4.92 (s, 2H), 3.84-3.94 (dd, 2H,  $J=7.0\text{Hz}$ ), 2.71-2.83 (m, 1H), 0.98 (m, 5H), 0.84 (m, 2H). MS calcd. 424.4; MS (M+1) 425.2

#### Example 20

10 N-(3-(2-Methoxyphenoxy)phenyl)-N-(cyclopropylsulfonyl)pyrid-3-ylmethylamine




The title compound was prepared using a method similar to the method of Example 1 using 3-(2-methoxyphenoxy)aniline and cyclopropylsulfonyl chloride.  $^1\text{H}$  NMR: 8.49 (d, 2H), 7.60 (d, 1H), 7.10-7.35 (m, 4H), 7.03-7.10 (d, 2H), 6.85-6.99 (m, 2H), 6.89 (s, 1H), 6.76 (m, 1H), 4.86 (s, 2H), 3.65 (s, 3H), 2.77 (m, 1H), 0.93 (m, 2H), 0.84 (m, 2H). MS calcd. 410.4; MS (M+1) 411.5

20

#### Example 21

N-(3-(2-Methoxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylamine




-51-

The title compound was prepared using a method similar to the method of Example 1 using 3-(2-methoxyphenoxy)aniline and ethylsulfonyl chloride. <sup>1</sup>H NMR: 8.64 (d, 2H, J=4.3Hz), 8.00 (t, 2H, J=6.6Hz), 7.72-7.76 (t, 1H, J=5.5Hz), 7.07-7.27 (m, 4H), 6.84-6.93 (m, 3H), 6.66 (d, 1H, J=8.1Hz), 5.00 (s, 2H), 3.64 (s, 3H), 3.25 (dd, 2H, J=15.0, 7.3Hz), 1.23 (t, 3H, J=7.7Hz). MS calcd. 398.3; MS (M+1) 399.2.

#### Example 22

10 N-(4-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethanamine




The title compound was prepared using a method similar to the method of Example 1 using (4-phenoxy)aniline. The hydrochloride salt was obtained by treatment of this material in methanol with ethereal HCl. <sup>1</sup>H NMR: 8.50 (m, 1H), 8.30 (s, 1H), 7.67 (m, 1H), 7.30-7.35 (m, 2H), 7.20-7.25 (m, 1H), 7.10-7.15 (m, 1H), 7.11 (d, 2H, J=9Hz), 6.99 (d, 2H, J=8Hz), 6.89 (d, 2H, J=9Hz), 4.83 (s, 2H), 3.80 (q, 2H, J=9Hz). MS calcd. 422.4; MS (M+1) 423.4.



-52-

Example 23


N-(3-(2-Hydroxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylamine



- 5           A solution of N-(3-(2-benzyloxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylamine (529mg, 1.00mmol) in methanol (50mL) was carefully treated with Raney<sup>®</sup> nickel (one tsp.), placed on a Parr<sup>®</sup> apparatus, and hydrogenated at 45-50 psi pressure for 3h. The mixture was  
10 carefully filtered through celite<sup>®</sup> under nitrogen without allowing the filter cake to dry, and the filtrate was concentrated in vacuo. The residual solid was triturated from petroleum ethers (30°-60°) to afford the title compound (394mg, 90%) as a white solid; mp 57.5-60.0°C. The title  
15 compound was converted to the hydrochloride salt by treatment in methanol with ethereal HCl. <sup>1</sup>H NMR: 8.50 (br s, 1H), 8.25 (br s, 1H), 7.68 (d, 1H, J=8Hz), 7.25-7.30 (m, 2H), 6.95-7.10 (m, 2H), 6.90-6.95 (m, 2H), 6.70-6.85 (m, 3H), 6.05 (br s, 1H), 4.82 (s, 2H), 3.75-3.85 (m, 2H). MS calcd. 438.4;  
20 MS (M+1) 439.3

Example 24

N-(4-(2-Hydroxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylamine



-53-

The title compound was prepared using a method similar to the method of Example 23 using N-(4-(2-benzyloxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline <sup>1</sup>H NMR: 8.48 (m,1H), 8.28 (s,1H), 7.70 (d,1H,J=8Hz), 7.20-7.25 (m,2H), 7.11 (d,2H,J=9Hz), 7.00-7.10 (m,2H), 6.90 (d,2H,J=9Hz), 6.80-6.85 (m,1H), 6.00 (br s,1H), 4.83 (s,2H), 3.75-3.85 (m,2H). MS calcd. 438.4; MS (M+1) 439.4

10

General Example AGeneral synthesis of compound of formula I

Compounds in the table below were prepared using substantially the procedure described as follows, using the corresponding reagents to provide the desired compound of formula I: Each well of a multi-well, 1.2 mL, glass, reaction plate was charged with an appropriate aryl aldehyde solution in methanol (1.0N, 100μL, 100μmol) followed by an appropriate aniline solution in methanol (0.65N, 200μL, 130μmol), and an additional 100μL of methanol was added to each well. The plate was capped and heated with shaking to 60°C for 18h, then cooled and uncapped. Sodium borohydride on alumina (10%, 60mg) was added via solid addition plate, and the reaction plate was shaken for 2h without caps and for 4h with caps. Plate contents were filtered through a filter plate into another multi-well, 1.2 mL, glass, reaction plate, and the empty plate was rinsed twice with 9:1 dichloromethane/methanol (200μL each) into the filter plate. The receiving plate was concentrated to dryness. A slurry made from polystyrene carboxaldehyde resin (1.8meq/g) and 4:1 1,2-dichloroethane/methanol (1g resin for each 10mL solvents) was added to each well (600μL/well), followed by 15% acetic acid/1,2-dichloroethane (100μL/well), and the plate was capped and shaken for 18h. The plate was uncapped, and DOWEX<sup>®</sup> 550A OH anion exchange resin (which had been washed dry with methanol) was added by solid addition plate (100mg per well). The plate was recapped and shaken for 2h, then uncapped and plate contents were filtered

-54-

through a filter plate into a multi-well, 2mL square well, plastic plate. The empty plate was rinsed twice with 9:1 dichloromethane/methanol (200 $\mu$ L each) into the filter plate, and the receiving plate was concentrated to dryness. The

5 plate contents were dissolved in 9:1 dichloromethane/methanol (400 $\mu$ L/well), and the solutions were passed through a filter plate containing a half volume of silica gel into a 96-well 1.2mL vial glass plate. The empty plate was rinsed twice with 9:1

10 dichloromethane/methanol (200 $\mu$ L each) into the filter plate, and the receiving plate was concentrated to dryness. In the case of sulfonation, each well is dissolved with 1:1 anhydrous 1,2-dichloroethane/pyridine (400 $\mu$ L), then treated with the appropriate sulfonyl chloride in anhydrous 1,2-

15 dichloroethane (1.0N, 200 $\mu$ L, 200 $\mu$ M). In the case of acylation, each well is dissolved with anhydrous 1,2-dichloroethane (400 $\mu$ L), then treated with the appropriate acid chloride or chloroformate ester in anhydrous 1,2-dichloroethane (1.0N, 200 $\mu$ L, 200 $\mu$ M). In either case, the

20 plate was capped and shaken for 48h, then uncapped, and each well was treated with tris(2-aminoethyl)amine resin (4meq/g, 70mg/well) by solid addition plate. Shaking with caps was continued for 3-4h, then the plate was opened, treated with dry DOWEX<sup>®</sup> 550A OH anion exchange resin (120mg/well) by

25 solid addition plate, and shaken with caps for 2-3h more. The plate contents were filtered through a filter plate into a 96-well receiving plate, the empty plate was rinsed twice with 9:1 dichloromethane/methanol (200 $\mu$ L each) into the filter plate, and the receiving plate was concentrated to

30 dryness to give a compound of formula I. The product plate was analyzed by whole plate mass spec and HPLC to give the compounds described in the table below. The HPLC conditions were as follows: a 4.6 x 50 mm C18 ODS<sub>am</sub> column (YMC, Inc. CombiScreen<sup>™</sup>) having a 3 micron particle size and 120

35 angstrom pore size was eluted at ambient temperature at a flow rate of 3 mL/min. The fluent was a gradient from 5-100% acetonitrile/water (containing 0.1% trifluoroacetic


-55-

acid) over 4 min. The methods of detection employed included a Gilson 170 diode array detector monitoring 254 nm and a SEDEX Model 75 evaporative light scattering detector held at a 40° C. Injection volumes were held between 5-10

5  $\mu$ L per injection. The product can optionally be obtained as the hydrochloride salt by dissolving the free base in minimum methanol and adding 1.3eq of 2.0N HCl in ether, concentrating *in vacuo* to obtain a residue, and the residue triturated from ether. In the table below the point of

10 attachment of the pyridyl moiety is designated with the substituent X. In the table below "M+1" refers to the mass ion, where observed, from mass spectral analysis using P.E. Sciex API 100 using was Positive APCI (Atmospheric Pressure Chemical Ionization) and heated nebulizer technique by (flow


15 injection analysis) with a run time of 30 sec per sample. All results yielded M+1 from positive APCI. The mobile phase used was 100% MeOH with 10 mM ammonium acetate using a Gilson 215 liquid handler injector and 1 $\mu$ L injections.


















20

No.	X	Y	R <sub>1</sub>	R <sub>2</sub>	R <sub>6</sub>	M+1
25	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	4-phenoxy	H	355.28
26	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-phenoxy	H	369.34
27	2-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>3</sub> CH <sub>3</sub>	4-phenoxy	H	383.39
28	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-phenoxy	H	423.39
29	2-CH <sub>2</sub>	bond	C(O)OCH <sub>3</sub>	4-phenoxy	H	335.83
30	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	4-phenoxy	H	355.43
31	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>3</sub> CH <sub>3</sub>	4-phenoxy	H	369.35
32	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-phenoxy	H	423.39
33	3-CH <sub>2</sub>	bond	C(O)OCH <sub>3</sub>	4-phenoxy	H	335.85
34	4-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	4-phenoxy	H	355.23
35	4-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-phenoxy	H	369.34
36	4-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>3</sub> CH <sub>3</sub>	4-phenoxy	H	397.51
37	4-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-phenoxy	H	423.41

38	4-CH <sub>2</sub>	bond	C(O)OCH <sub>3</sub>	4-phenoxy	H	335.84
39	2-CH <sub>2</sub>	bond	C(O)CH <sub>2</sub> CH <sub>3</sub>	4-phenoxy	H	333.57
40	2-CH <sub>2</sub>	bond	C(O)-cyclopentyl	4-phenoxy	H	373.32
41	3-CH <sub>2</sub>	bond	C(O)CH <sub>2</sub> CH <sub>3</sub>	4-phenoxy	H	333.58
42	3-CH <sub>2</sub>	bond	C(O)-cyclopentyl	4-phenoxy	H	373.53
43	4-CH <sub>2</sub>	bond	C(O)CH <sub>2</sub> CH <sub>3</sub>	4-phenoxy	H	333.59
44	4-CH <sub>2</sub>	bond	C(O)-cyclopentyl	4-phenoxy	H	373.32
45	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	4-phenoxy	H	423.34
46	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	4-phenoxy	H	423.4
47	4-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	4-phenoxy	H	423.4
48	2-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	4-phenoxy	H	383.38
49	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	4-phenoxy	H	383.4
50	4-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	4-phenoxy	H	383.4
51	2-CH <sub>2</sub>	bond	C(O)CH <sub>2</sub> CH(CH <sub>3</sub> ) 2	4-phenoxy	H	361.93
52	3-CH <sub>2</sub>	bond	C(O)CH <sub>2</sub> CH(CH <sub>3</sub> ) 2	4-phenoxy	H	361.93
53	4-CH <sub>2</sub>	bond	C(O)CH <sub>2</sub> CH(CH <sub>3</sub> ) 2	4-phenoxy	H	361.9
54	2-CH <sub>2</sub>	bond	C(O)CH <sub>3</sub>	4-phenoxy	H	319.18
55	3-CH <sub>2</sub>	bond	C(O)CH <sub>3</sub>	4-phenoxy	H	319.23
56	4-CH <sub>2</sub>	bond	C(O)CH <sub>3</sub>	4-phenoxy	H	319.17
57	2-CH <sub>2</sub>	bond	C(O)-cyclopropyl	4-phenoxy	H	345.16
58	3-CH <sub>2</sub>	bond	C(O)-cyclopropyl	4-phenoxy	H	345.15
59	4-CH <sub>2</sub>	bond	C(O)-cyclopropyl	4-phenoxy	H	345.12
60	3-CH <sub>2</sub>	bond	C(O)-cyclopropyl	4-(4-benzyloxy- phenoxy)	H	359.57
61	3-CH <sub>2</sub>	bond	C(O)-cyclopropyl	4-(4-cyclohexyl- phenoxy)	H	335.93
62	3-CH <sub>2</sub>	bond	C(O)-cyclopropyl	4-(4-benzyl- phenoxy)	H	343.41
63	3-CH <sub>2</sub>	bond	C(O)-cyclopropyl	4-(4-thiophenoxy- phenoxy)	H	361.88
64	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	4-(4-benzyloxy- phenoxy)	H	369.32
65	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	4-(4-cyclohexyl- phenoxy)	H	345.18
66	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	4-(4-benzyl- phenoxy)	H	353.62
67	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	4-(4-thiophenoxy- phenoxy)	H	371.08
68	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(4-benzyloxy- phenoxy)	H	383.42
69	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(4-cyclohexyl- phenoxy)	H	359.5
70	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(4-benzyl- phenoxy)	H	367.55

71	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(4-thiophenoxy- phenoxy)	H	385.5
72	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(4-benzyloxy- phenoxy)	H	437.78
73	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(4-cyclohexyl- phenoxy)	H	413.65
74	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(4-benzyl- phenoxy)	H	421.19
75	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(4-thiophenoxy- phenoxy)	H	439.52
76	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	4-(4-benzyloxy- phenoxy)	H	423.4
77	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	4-(4-cyclohexyl- phenoxy)	H	399.4
78	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	4-(4-benzyl- phenoxy)	H	407.4
79	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	4-(4-thiophenoxy- phenoxy)	H	425.5
80	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	4-(4-benzyloxy- phenoxy)	H	397.5
81	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	4-(4-cyclohexyl- phenoxy)	H	373.5
82	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	4-(4-benzyl- phenoxy)	H	381.63
83	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	4-(4-thiophenoxy- phenoxy)	H	399.5
84	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(4-benzyloxy- phenoxy)	H	383.57
85	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(4-cyclohexyl- phenoxy)	H	359.55
86	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(4-benzyl- phenoxy)	H	367.5
87	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(4-thiophenoxy- phenoxy)	H	385.5
88	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(4-benzyloxy- phenoxy)	H	437.78
89	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(4-cyclohexyl- phenoxy)	H	413.65
90	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(4-benzyl- phenoxy)	H	421.16
91	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(4-thiophenoxy- phenoxy)	H	439.49
92	3-CH <sub>2</sub>	bond	C(O)-cyclopropyl	4-pentoxy	H	339.31
93	3-CH <sub>2</sub>	bond	C(O)-cyclopropyl	4- 	H	297.63
94	3-CH <sub>2</sub>	bond	C(O)-cyclopropyl	4-methoxy	H	283.69

95	3-CH <sub>2</sub>	bond	C(O)-cyclopropyl	4- 	H	399.33
96	3-CH <sub>2</sub>	bond	C(O)-cyclopropyl	4-(4-cyclohexoxy-phenoxy)	H	351.86
97	3-CH <sub>2</sub>	bond	C(O)-cyclopropyl	4-(4-pyrid-2-yloxy-phenoxy)	H	346.31
98	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	4-pentoxy	H	349.82
99	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	4- 	H	307.71
100	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	4-methoxy	H	293.21
101	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	4- 	H	409.41
102	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	4-(4-cyclohexoxy-phenoxy)	H	361.9
103	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	4-(4-pyrid-2-yloxy-phenoxy)	H	356.37
104	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-pentoxy	H	363.83
105	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4- 	H	321.41
106	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-methoxy	H	307.71
107	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4- 	H	423.49
108	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(4-cyclohexoxy-phenoxy)	H	375.65
109	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(4-pyrid-2-yloxy-phenoxy)	H	370.05
110	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-pentoxy	H	417.65
111	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4- 	H	375.53
112	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-methoxy	H	361.81
113	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4- 	H	477.29
114	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(4-cyclohexoxy-phenoxy)	H	429.83
115	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(4-pyrid-2-yloxy-phenoxy)	H	424.41
116	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	4-pentoxy	H	403.4

117	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	4- 	H	361.3
118	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	4-methoxy	H	347.3
119	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	4- 	H	463.5
120	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	4-(4-cyclohexoxy-phenoxy)	H	415.4
121	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	4-(4-pyrid-2-yloxy-phenoxy)	H	401.4
122	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	4-pentoxy	H	377.76
123	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	4- 	H	335.85
124	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	4-methoxy	H	321.41
125	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	4- 	H	437.87
126	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	4-(4-cyclohexoxy-phenoxy)	H	389.88
127	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	4-(4-pyrid-2-yloxy-phenoxy)	H	384.61
128	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-pentoxy	H	363.81
129	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4- 	H	321.4
130	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-methoxy	H	307.71
131	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4- 	H	423.47
132	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(4-cyclohexoxy-phenoxy)	H	375.67
133	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(4-pyrid-2-yloxy-phenoxy)	H	361.5
134	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-pentoxy	H	417.66
135	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4- 	H	375.52
136	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-methoxy	H	361.79
137	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4- 	H	477.5
138	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(4-cyclohexoxy-phenoxy)	H	429.62
139	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(4-pyrid-2-	H	424.36



				xyloxy-phenoxy)		
140	3-CH <sub>2</sub>	bond	C(O)-cyclopropyl	3-phenoxy	H	345.16
141	3-CH <sub>2</sub>	bond	C(O)-cyclopropyl	2-phenoxy	H	345.18
142	3-CH <sub>2</sub>	bond	C(O)-cyclopropyl	4-(4-chloro-phenoxy)	H	379.4
143	3-CH <sub>2</sub>	bond	C(O)-cyclopropyl	4-(3-trifluoromethyl-phenoxy)	H	413.68
144	3-CH <sub>2</sub>	bond	C(O)-cyclopropyl	4-(2-methoxy-phenoxy)	H	375.65
145	3-CH <sub>2</sub>	bond	C(O)-cyclopropyl	4-(4-cyano-phenoxy)	H	370.14
146	3-CH <sub>2</sub>	bond	C(O)-cyclopropyl	4-(2,4-difluoro-phenoxy)	H	381.62
147	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	3-phenoxy	H	355.28
148	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	2-phenoxy	H	355.4
149	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	4-(4-chloro-phenoxy)	H	389.78
150	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	4-(3-trifluoromethyl-phenoxy)	H	423.4
151	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	4-(2-methoxy-phenoxy)	H	385.44
152	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	4-(4-cyano-phenoxy)	H	380.52
153	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	4-(2,4-difluoro-phenoxy)	H	391.63
154	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-phenoxy	H	369.3
155	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	2-phenoxy	H	369.3
156	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(4-chloro-phenoxy)	H	403.9
157	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(3-trifluoromethyl-phenoxy)	H	437.5
158	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-methoxy-phenoxy)	H	399.26
159	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(4-cyano-phenoxy)	H	448.44
160	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2,4-difluoro-phenoxy)	H	459.37
161	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-phenoxy	H	423.43
162	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	2-phenoxy	H	423.39
163	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(4-chloro-phenoxy)	H	457.68
164	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(3-trifluoromethyl-phenoxy)	H	491.56

165	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-methoxy- phenoxy)	H	453.36
166	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(4-cyano- phenoxy)	H	448.44
167	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2,4-difluoro- phenoxy)	H	459.37
168	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-phenoxy	H	369.5
169	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	2-phenoxy	H	369.5
170	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(4-chloro- phenoxy)	H	403.75
171	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(3- trifluoromethyl- phenoxy)	H	437.78
172	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-methoxy- phenoxy)	H	399.28
173	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(4-cyano- phenoxy)	H	394.5
174	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2,4-difluoro- phenoxy)	H	405.5
175	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-phenoxy	H	423.41
176	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	2-phenoxy	H	423.4
177	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(4-chloro- phenoxy)	H	457.7
178	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(3- trifluoromethyl- phenoxy)	H	491.57
179	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-methoxy- phenoxy)	H	453.63
180	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(4-cyano- phenoxy)	H	448.45
181	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2,4-difluoro- phenoxy)	H	459.37
182	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	3-phenoxy	H	409.4
183	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	2-phenoxy	H	409.4
184	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	4-(4-chloro- phenoxy)	H	443.8
185	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	4-(3- trifluoromethyl- phenoxy)	H	477.4
186	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	4-(2-methoxy- phenoxy)	H	439.4
187	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	4-(4-cyano- phenoxy)	H	434.4
188	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	4-(2,4-difluoro- phenoxy)	H	445.4
189	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	3-phenoxy	H	383.5
190	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	2-phenoxy	H	383.5

191	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	4-(4-chloro-phenoxy)	H	417.9
192	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	4-(3-trifluoromethyl-phenoxy)	H	451.5
193	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	4-(2-methoxy-phenoxy)	H	413.5
194	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	4-(4-cyano-phenoxy)	H	408.5
195	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	4-(2,4-difluoro-phenoxy)	H	419.35
196	3-CH <sub>2</sub>	bond	C(O)-cyclopropyl	2-chloro-5-methyl-4-phenoxy	H	393.9
197	3-CH <sub>2</sub>	bond	C(O)-cyclopropyl	3-chloro-4-phenoxy	H	379.33
198	3-CH <sub>2</sub>	bond	C(O)-cyclopropyl	2-methyl-4-phenoxy	H	359.55
199	3-CH <sub>2</sub>	bond	C(O)-cyclopropyl	3-trifluoromethyl-4-phenoxy	H	413.63
200	3-CH <sub>2</sub>	bond	C(O)-cyclopropyl	4-(4-nitro-phenoxy)	H	390.75
201	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	2-chloro-5-methyl-4-phenoxy	H	403.9
202	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	3-chloro-4-phenoxy	H	389.71
203	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	2-methyl-4-phenoxy	H	369.3
204	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	3-trifluoromethyl-4-phenoxy	H	423.4
205	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>3</sub>	4-(4-nitro-phenoxy)	H	400.4
206	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	2-chloro-5-methyl-4-phenoxy	H	417.9
207	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-chloro-4-phenoxy	H	403.9
208	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	2-methyl-4-phenoxy	H	383.5
209	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-trifluoromethyl-4-phenoxy	H	437.5
210	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(4-nitro-phenoxy)	H	414.5
211	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	2-chloro-5-methyl-4-phenoxy	H	471.9
212	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-chloro-4-phenoxy	H	457.9
213	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	2-methyl-4-phenoxy	H	437.72

214	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-trifluoromethyl-4-phenoxy	H	491.51
215	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(4-nitro-phenoxy)	H	468.89
216	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	2-chloro-5-methyl-4-phenoxy	H	457.9
217	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	3-chloro-4-phenoxy	H	443.8
218	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	2-methyl-4-phenoxy	H	423.4
219	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	3-trifluoromethyl-4-phenoxy	H	477.4
220	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CF <sub>3</sub>	4-(4-nitro-phenoxy)	H	454.4
221	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	2-chloro-5-methyl-4-phenoxy	H	432.0
222	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	3-chloro-4-phenoxy	H	417.9
223	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	2-methyl-4-phenoxy	H	397.5
224	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	3-trifluoromethyl-4-phenoxy	H	451.5
225	3-CH <sub>2</sub>	bond	SO <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	4-(4-nitro-phenoxy)	H	428.5
226	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	2-chloro-5-methyl-4-phenoxy	H	417.9
227	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-chloro-4-phenoxy	H	403.9
228	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	2-methyl-4-phenoxy	H	383.5
229	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-trifluoromethyl-4-phenoxy	H	437.5
230	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(4-nitro-phenoxy)	H	414.5
231	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	2-chloro-5-methyl-4-phenoxy	H	471.6
232	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-chloro-4-phenoxy	H	457.66
233	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	2-methyl-4-phenoxy	H	437.73
234	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-trifluoromethyl-4-phenoxy	H	491.52
235	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(4-nitro-phenoxy)	H	466.87
236	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-phenoxy	6-methoxy	453
237	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-phenoxy	6-	453

					methoxy	
238	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-methoxy-phenoxy)	6-methoxy	484
239	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-phenoxy	6-methoxy	399
240	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-phenoxy	6-methoxy	400
241	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-methoxy-phenoxy)	6-methoxy	430
242	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-phenoxy	6-methyl	438
243	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-phenoxy	6-methyl	438
244	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-methoxy-phenoxy)	6-methyl	468
245	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-phenoxy	6-methyl	384
246	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-phenoxy	6-methyl	384
247	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-methoxy-phenoxy)	6-methyl	414
248	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-phenoxy	6-(4-chloro-phenoxy)	550
249	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-phenoxy	6-(4-chloro-phenoxy)	550
250	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-methoxy-phenoxy)	6-(4-chloro-phenoxy)	580
251	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-phenoxy	6-(4-chloro-phenoxy)	496
252	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-phenoxy	6-(4-chloro-phenoxy)	496
253	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-methoxy-phenoxy)	6-(4-chloro-phenoxy)	526
254	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-phenoxy	2-chloro	458
255	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-phenoxy	2-chloro	458
256	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-methoxy-phenoxy)	2-chloro	488
257	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-phenoxy	2-chloro	404
258	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-phenoxy	2-chloro	404
259	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-methoxy-phenoxy)	2-chloro	434
260	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-phenoxy	2-phenoxy	516
261	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-phenoxy	2-phenoxy	516

262	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-methoxy-phenoxy)	2-phenoxy	546
263	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-phenoxy	2-phenoxy	462
264	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-phenoxy	2-phenoxy	462
265	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-methoxy-phenoxy)	2-phenoxy	492
266	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-phenoxy	6-chloro	458
267	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-phenoxy	6-chloro	458
268	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-methoxy-phenoxy)	6-chloro	488
269	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-phenoxy	6-chloro	404
270	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-phenoxy	6-chloro	404
271	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-methoxy-phenoxy)	6-chloro	434
272	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-phenoxy	2-chloro-6-methyl	472
273	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-phenoxy	2-chloro-6-methyl	471
274	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-methoxy-phenoxy)	2-chloro-6-methyl	501
275	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-phenoxy	2-chloro-6-methyl	418
276	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-phenoxy	2-chloro-6-methyl	418
277	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-methoxy-phenoxy)	2-chloro-6-methyl	448
278	3-CH <sub>2</sub>	bond	SO <sub>2</sub> -cyclopropyl	4-phenoxy	H	382
279						
280	4-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-methoxy-phenoxy)	H	453
281	4-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-methoxy-phenoxy)	H	399
282	3-CH <sub>2</sub>	bond	SO <sub>2</sub> -cyclopropyl	4-(2,6-dimethoxy-phenoxy)	H	442
283	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2,6-dimethoxy-phenoxy)	H	484
284	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2,6-dimethoxy-phenoxy)	H	430
285	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2,6-dimethoxy-phenoxy)	H	399
286	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2,6-dimethoxy-phenoxy)	H	453
287	4-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2,6-dimethoxy-phenoxy)	H	399
288	4-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2,6-dimethoxy-	H	453

				phenoxy)		
289	3-CH <sub>2</sub>	CH <sub>2</sub>	SO <sub>2</sub> -cyclopropyl	2-phenoxy-6-chloro	H	430
290	3-CH <sub>2</sub>	CH <sub>2</sub>	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	2-phenoxy-6-chloro	H	418
291	3-CH <sub>2</sub>	CH <sub>2</sub>	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	2-phenoxy-6-chloro	H	472
292	2-CH <sub>2</sub>	CH <sub>2</sub>	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	2-phenoxy-6-chloro	H	418
293	2-CH <sub>2</sub>	CH <sub>2</sub>	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	2-phenoxy-6-chloro	H	472
294	4-CH <sub>2</sub>	CH <sub>2</sub>	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	2-phenoxy-6-chloro	H	418
295	4-CH <sub>2</sub>	CH <sub>2</sub>	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	2-phenoxy-6-chloro	H	472
296	3-CH <sub>2</sub>	bond	SO <sub>2</sub> -cyclopropyl	4-(2,6-dichloro-phenoxy)	H	449
297	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2,6-dichloro-phenoxy)	H	493
298	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2,6-dichloro-phenoxy)	H	439
299	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2,6-dichloro-phenoxy)	H	439
300	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2,6-dichloro-phenoxy)	H	494
301	4-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2,6-dichloro-phenoxy)	H	439
302	4-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2,6-dichloro-phenoxy)	H	493
303	3-CH <sub>2</sub>	bond	SO <sub>2</sub> -cyclopropyl	4-(2-isopropoxy-phenoxy)	H	440
304	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-isopropoxy-phenoxy)	H	482
305	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-isopropoxy-phenoxy)	H	427
306	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-isopropoxy-phenoxy)	H	427
307	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-isopropoxy-phenoxy)	H	481
308	4-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-isopropoxy-phenoxy)	H	427
309	4-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-isopropoxy-phenoxy)	H	481
310	3-CH <sub>2</sub>	bond	SO <sub>2</sub> -cyclopropyl	4-(2-fluoro-6-methoxy-phenoxy)	H	430
311	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-fluoro-6-	H	472

				methoxy- phenoxy)		
312	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-fluoro-6-methoxy- phenoxy)	H	418
313	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-fluoro-6-methoxy- phenoxy)	H	418
314	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-fluoro-6-methoxy- phenoxy)	H	472
315	4-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-fluoro-6-methoxy- phenoxy)	H	418
316	4-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-fluoro-6-methoxy- phenoxy)	H	471
317	3-CH <sub>2</sub>	bond	SO <sub>2</sub> -cyclopropyl	4-(2-trifluoromethyl- phenoxy)	H	449
318	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-trifluoromethyl- phenoxy)	H	492
319	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-trifluoromethyl- phenoxy)	H	438
320	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-trifluoromethyl- phenoxy)	H	437
321	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-trifluoromethyl- phenoxy)	H	492
322	4-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-trifluoromethyl- phenoxy)	H	437
323	4-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-trifluoromethyl- phenoxy)	H	492
324	3-CH <sub>2</sub>	bond	SO <sub>2</sub> -cyclopropyl	4-(2-isopropyl-5-methyl- phenoxy)	H	438
325	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-isopropyl-5-methyl- phenoxy)	H	479
326	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-isopropyl-5-methyl- phenoxy)	H	425
327	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-isopropyl-5-methyl- phenoxy)	H	426
328	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-isopropyl-5-	H	479



-68-

				methyl-phenoxy)		
329	4-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-isopropyl-5-methyl-phenoxy)	H	425
330	4-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-isopropyl-5-methyl-phenoxy)	H	479
331	3-CH <sub>2</sub>	bond	SO <sub>2</sub> -cyclopropyl	4-(2-cyclopentyl-phenoxy)	H	450
332	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-cyclopentyl-phenoxy)	H	492
333	3-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-cyclopentyl-phenoxy)	H	438
334	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-cyclopentyl-phenoxy)	H	438
335	2-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-cyclopentyl-phenoxy)	H	492
336	4-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-(2-cyclopentyl-phenoxy)	H	438
337	4-CH <sub>2</sub>	bond	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-cyclopentyl-phenoxy)	H	492

### Preparation 3

#### 4-(4-Benzyloxyphenoxy) aniline

A solution of 4-benzyloxyphenol (6.01g, 30mmol) in  
 5 anhydrous dimethylformamide (50ml) under a nitrogen cover  
 was treated with sodium hydride (1.2g of 60% oil dispersion,  
 30mmol) in one portion at room temperature. A moderate gas  
 evolution ensued, and the reaction temperature rose to 41°C.  
 After stirring an additional 35 minutes,  
 10 4-Fluoronitrobenzene was added rapidly dropwise, neat via  
 syringe. A mild exotherm brought the reaction temperature  
 to 32°C, after which the reaction was stirred for 18 hours  
 at room temperature. The reaction mixture was poured over  
 ice/water (250ml), stirred for thirty minutes and the yellow  
 15 solid collected by suction filtration. Silica gel  
 chromatography (Waters® LC2000; two Pre-Pak® columns,  
 eluting with 3/1 hexanes/methylene chloride) afforded 4-(4-  
 benzyloxyphenoxy)-1-nitrobenzene (5.9g, 73%) as a white  
 solid. <sup>1</sup>H NMR(400MHz): 5.08(2H,s), 6.94-6.99(2H,m),

-69-


7.02 (4H, s), 7.32-7.47 (5H, m), 8.15-8.20 (2H, m); MS (ESI) m/z 320.2 ([M<sup>+</sup>H] -, 100).

A mechanically-stirred solution of 4-(4-benzyloxyphenoxy)-1-nitrobenzene (5.6g, 17mmol) in absolute ethanol (50ml) was treated with concentrated hydrochloric acid (6ml). The solution was warmed to 55-60°C under a nitrogen cover. Iron powder (6.5g, 115mmol) was added in several portions. A mild exotherm (while still being heated) brought the reaction temperature to 68°C. After completing the iron addition the reaction temperature was brought to 70°C and maintained at that temperature for 1.5 hours. The mixture was then cooled to room temperature and filtered through a pad of Celite®. The filter cake was washed with ethanol (100ml) and methylene chloride (50ml), and the filtrate partitioned between methylene chloride and 1M potassium carbonate. The green aqueous layer was extracted a second time with methylene chloride and then dried over anhydrous magnesium sulfate. Filtration and concentration in vacuo afforded crude product (5.4g), contaminated with unreacted starting material. Silica gel chromatography (Waters® LC2000 with 2 PrePak® columns, eluting with methylene chloride) provided the title compound (3.75g, 76%), as well as unreacted starting material (1.2g). <sup>1</sup>H NMR (400MHz): 3.53 (2H, br s), 5.02 (2H, s), 6.62-6.67 (2H, m), 6.79-6.84 (2H, m), 6.86-6.93 (4H, m), 7.29-7.44 (5H, m); MS (ESI) m/z 291.9 ([M+H]<sup>+</sup>, 100).

-70-

Example 338

N-(4-(4-Benzyloxyphenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylamine



5           4-(4-Benzyloxyphenoxy)aniline (3.4g, 11.67mmol) was suspended in methanol (100ml) and treated with pyridine-3-carboxaldehyde (1.25g, 11.67mmol) in one portion via syringe. This solution was heated to reflux for two hours, and then cooled to -5°C. Solid sodium borohydride (0.88g, 10 23mmol) was added in one portion. A mild gas evolution ensued, and the reaction temperature rose to -2°C. The mixture was then stirred at room temperature for two hours, after which TLC analysis of an aliquot showed incomplete reduction. An additional quantity of sodium borohydride was 15 added (0.45g, 12mmol) was added and the mixture stirred another two hours. The solvent was removed in vacuo and the residue slurried in methylene chloride. After filtration to remove insolubles and concentration in vacuo, the crude oil thereby obtained was purified via silica gel chromatography 20 (Waters® LC2000 using two PrePak® columns, eluting with 99/1 methylene chloride/methanol). N-4-(4-Benzyloxyphenoxy)phenylpyrid-3-ylmethylamine was obtained as a white solid (3.6g, 81%). <sup>1</sup>H NMR(400MHz): 4.33(2H,s), 5.02(2H,s), 6.56-6.62(2H,m), 6.83-6.92(6H,m), 7.24- 7.44(6H,m), 7.70(1H,ddd,J=7.8,~2.0,~2.0), 8.53(1H,dd, J=4.9,1.5), 8.63(1H,d,J=1.5); MS(ESI)m/z 382.9([M+H]<sup>+</sup>,100).


N-4-(4-Benzyloxyphenoxy)phenylpyrid-3-ylmethylamine (3.55g, 9.3mmol) was dissolved in 1,2-dichloroethane (46ml) in a flame-dried, serum-capped round bottom flask under a

-71-

nitrogen cover. Anhydrous pyridine (24ml) was added via syringe, the solution cooled to  $-5^{\circ}\text{C}$ , and trifluoroethanesulfonyl chloride was added neat, dropwise via syringe. After stirring thirty minutes at  $0^{\circ}\text{C}$  the solution was stirred at room temperature for 18 hours. Potassium carbonate (6.0g) was added to the mixture and it was stirred another hour. The mixture was then filtered and the filtrate concentrated in vacuo to a dark oil. Silica gel chromatography (Waters® LC2000 using two PrePak® columns, eluting with 99/1 methylene chloride/methanol). Product-containing fractions were combined and concentrated, and the resulting mixture chromatographed a second time using a 90/10 to 84/16 methylene chloride/ethyl acetate gradient. The title compound was obtained (4.36g, 89%) as an off-white solid.  $^1\text{H}$  NMR(400MHz): 3.81(2H,q,J=9.3), 4.84(2H,s), 5.05(2H,s), 6.83-6.88(2H,m), 6.96(4H,s), 7.07-7.12(2H,m), 7.26(1H,dd,J=4.9,3.9), 7.30-7.46(4H,m), 7.71(1H,ddd,J=7.8,2.0,2.0), 8.32(1H,d,J=1.9), 8.52(1H,dd,J=4.9,1.4); MS(ESI)m/z 529.0([M+H]<sup>+</sup>,100); Anal. Calcd. for  $\text{C}_{27}\text{H}_{23}\text{F}_3\text{N}_2\text{O}_4\text{S}$  C,61.36; H,4.39; N,5.30. Found: C,61.32; H,4.44; N,5.31.

#### Example 339

N-(4-(4-Hydroxyphenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethanamine



N-4-(4-Benzoyloxyphenoxy)phenyl-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethanamine (3.0g, 5.68mmol) was dissolved in absolute ethanol/ethyl acetate

-72-

(100ml, 1/1) in a Parr® bottle. Raney nickel (approximately 1-1.5 tsp.) was added and the mixture hydrogenated at 40psi on a shaker for twenty-two hours. After filtering to remove the catalyst, the filtrate was concentrated in vacuo to provide crude product as a white solid. Recrystallization from hot ethyl acetate/methanol (6/1) provided the title compound (1.9g, 76%). <sup>1</sup>H NMR(400MHz, DMSO-d<sub>6</sub>): 4.73(2H,q,J=10.3), 4.88(2H,s), 6.75-6.84(4H,m), 6.85-6.91(2H,m), 7.23-7.29(2H,m), 7.34(1H,dd,J=7.8,4.9), 7.67(1H,m), 8.38(1H,d,J=2.0), 8.45(1H,dd,J=4.9,1.5), 9.4(1H,s); MS(ESI)m/z 439.1([M+H]<sup>+</sup>,100); Anal. Calcd. for C<sub>20</sub>H<sub>17</sub>F<sub>3</sub>N<sub>2</sub>O<sub>4</sub>s C,54.79; H,3.91; N,6.39. Found: C,54.72; H,3.85; N,6.32.

#### Preparation 4

##### 3-(2-Methoxyphenylsulfinyl)aniline

A solution of 3-fluoronitrobenzene (9.50g, 67.3mmol) and 2-methoxythiophenol (7.50g, 53.5mmol) in anhydrous DMF (40mL) under nitrogen was treated with anhydrous potassium carbonate (14g, 101mmol), and heated to 115°C for 18h. The solution was cooled, filtered (solid rinsed with methylene chloride), and the filtrate concentrated exhaustively in vacuo under high vacuum to remove all DMF. The residue was taken up in water and extracted with ether (2X75mL). The combined ether solution was dried (MgSO<sub>4</sub>), filtered, the filtrate concentrated in vacuo and the residue chromatographed on silica gel (eluted with 2:1 hexane/methylene chloride) to afford 4.01g (29%) of 2-methoxy-3'-nitrodiphenylsulfide as a viscous orange oil. <sup>1</sup>H NMR: 3.82(3H,s), 6.95-7.00(2H,m), 7.35-7.45(4H,m), 7.95-8.00(2H,m).

2-Methoxy-3'-nitrodiphenylsulfide (332mg, 1.27mmol) was taken up in chloroform (1.5ml) and treated with trichloroacetone (250μl, 2.8mmol), followed immediately by

-73-


the addition of aqueous hydrogen peroxide (250 $\mu$ l of 30%, 2.4mmol). The reaction was stirred at room temperature for twenty-four hours and then partitioned between methylene chloride and aqueous sodium bisulfite. The organic layer was washed once with brine, dried over magnesium sulfate, filtered and concentrated to an oil. Purification by radial silica gel chromatography (4mm Chromatotron<sup>®</sup> plate, eluting with 2/1 hexanes/ethyl acetate) afforded 1-methoxy-2-(3-nitrophenylsulfinyl)benzene (272mg, 77%) as an oil which crystallized on standing. <sup>1</sup>H NMR(400MHz): 3.89(3H,s), 6.90(1H,d,J=7.3), 7.18(1H,dt,J=7.8,1.0), 7.41-7.47(1H,m), 7.63(1H,t, J=7.8), 7.89(1H,dd,J=7.8,1.4), 8.07-8.12(1H,m), 8.22-8.28(1H,m), 8.61(1H,m,); MS(ESI)m/z 276.8([M-H]<sup>-</sup>,70); Anal. Calcd. for C<sub>13</sub>H<sub>11</sub>NO<sub>4</sub>S C,56.31; H,4.00; N,5.05. Found: C,56.34; H,4.01; N,5.05.

1-Methoxy-2-(3-nitrophenylsulfinyl)benzene (217mg, 0.78mmol) was dissolved in ethyl alcohol (5.0ml) and water (1.75ml), to which was added iron powder (488mg, 8.7mmol) and ammonium chloride (77mg, 1.4mmol). This mixture was heated to reflux for four hours, cooled to room temperature and the supernatant decanted. The insoluble material was rinsed with methylene chloride, followed by aqueous 1M potassium carbonate. All decanted solutions were combined and the organic layer separated. The aqueous layer was extracted again with methylene chloride, and the combined organics washed once with brine. After drying over magnesium sulfate and concentration in vacuo, the title compound was obtained as a white solid (145mg, 75%). <sup>1</sup>H NMR(400MHz): 3.89(3H,s), 6.69-6.74(1H,m), 6.86(1H,d,J=8.4), 7.03-7.09(2H,m), 7.10-7.20(2H,m), 7.38(1H,dt,J=8.3,1.5), 7.86(1H,dd,J=7.8,1.5).

-74-

Example 340

N-(3-(2-Methoxyphenylsulfinyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine



5           3-(1-Methoxyphenylsulfinyl)aniline (145mg, 0.59mmol)  
and pyridine-3-carboxaldehyde (63mg, 0.59mmol) were reacted  
in anhydrous methanol (3.0ml), followed by treatment with  
sodium borohydride (44mg, 1.17mmol) as previously described  
(Intermediate 3). Silica gel chromatography (2mm  
10 Chromatotron® plate, eluting with 20/1 ethyl  
acetate/hexanes) afforded N-(3-(2-  
methoxyphenylsulfinyl)phenyl)pyrid-3-ylmethylamine (162mg,  
82%) as an oil. <sup>1</sup>H NMR(400MHz): 3.77(3H,s), 4.35(2H,br s),  
6.61(1H,m), 6.83(1H,d,J=8.0), 6.96(1H,br t), 7.01-  
15 7.04(1H,m), 7.11(1H,dt,J=7.8,1.0), 7.18(1H,t,J=7.8), 7.21-  
7.28(1H,br s), 7.35-7.41(1H,m), 7.63(1H,br d),  
7.82(2H,dd,J=7.8,1.5), 8.40-8.80(2H,br s); MS(ESI)m/z  
339.1([M+H]<sup>+</sup>,100)

          N-(3-(2-Methoxyphenylsulfinyl)phenyl)pyrid-3-  
20 ylmethylamine (159mg, 0.44mmol) was taken up in 1,2-  
dichloroethane (2.0ml) and anhydrous pyridine (1.0ml) and  
cooled under nitrogen to 0°C. Ethanesulfonyl chloride  
(103mg, 0.80mmol) is added dropwise via syringe and the  
reaction allowed to stir at 0°C for thirty minutes. The ice  
25 bath is removed and the mixture stirred 88 hours at room  
temperature, as a matter of convenience. Solid potassium  
carbonate was added, the reaction stirred an additional  
hour, and then filtered and concentrated to a dark oil.  
Radial silica gel chromatography (2mm Chromatotron® plate,

-75-

eluting with 20/1 methylene chloride/methanol) afforded the title compound (16mg, 8%).  $^1\text{H}$  NMR(400MHz): 1.39(3H,t,J=7.3), 3.04(2H,q,J=7.3), 3.78(3H,s), 4.88(2H,s), 6.86(1H,d,J=7.8), 7.11-7.21(2H,m), 7.25-7.30(1H,m), 7.35(1H,t,J=7.8), 7.42(1H,dt,J=7.3,2.0), 7.56-7.62(2H,m), 7.68(1H,m), 7.81(1H,dd,J=7.3,1.5), 8.31(1H,br s), 8.46(1H,br s); MS(HRMS)m/z 431.1116([M+H]<sup>+</sup> calcd for C<sub>21</sub>H<sub>23</sub>N<sub>2</sub>O<sub>4</sub>S 431.1099).

#### Preparation 5

##### 4-(2,4-Difluorophenoxy)aniline

##### 10      1-(2,4-Difluorophenoxy)-4-nitrobenzene

A biphasic mixture of potassium carbonate (304 g, 2.20mol), 1-fluoro-4-nitrobenzene (148mL, 1.40mol) and 2,4-difluorophenol (200g, 1.54mol) in N,N-dimethylformamide (1.0 L) was heated at 90°C under nitrogen for 22 h, after which the mixture was cooled to room temperature. The mixture was poured into water (1.0 L) and was extracted with ethyl acetate (2 x 1.5 L). The combined organic extracts were washed with water (1 x 1.0 L), dried over MgSO<sub>4</sub>, filtered and the solvent was removed under reduced pressure to afford the crude product as a yellow semi-solid. This residue was triturated with hexanes (500mL) and gravity filtered, washing with hexanes (2 x 100mL) to provide 1-(2,4-difluorophenoxy)-4-nitrobenzene as off-white crystals (209 g, 60%), mp 58-62°C.  $^1\text{H}$  NMR(300 MHz, DMSO-d<sub>6</sub>): 7.15(2H,d,J=9.4), 7.27-7.20(1H,m), 7.62-7.48(2H,m), 8.25(2H,d,J=9.4).

A suspension of 1-(2,4-difluorophenoxy)-4-nitrobenzene (209g, 1.17mol) and 10% palladium-on-carbon (20g, 50% wet) in methanol (1.0 L) was shaken under an atmosphere of hydrogen (60psi) at room temperature for 5 h. The mixture was filtered through a plug of Celite® (150g), washing with methanol (2 x 300mL) and the solvent was removed under reduced pressure to afford the crude product as a dark red oil. This residue was triturated with hexanes (500mL) and




-76-

filtered to provide the title compound as a gray solid (175 g, 95%).  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ ): 6.55 (2H, d,  $J=8.8$ ), 6.73 (2H, d,  $J=8.8$ ), 7.02-6.95 (2H, m), 7.40 (1H, t,  $J=11.4$ ).

5

Example 341

N-(4-(2,4-Difluorophenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylamine



4-(2,4-Difluorophenoxy)aniline (2.21g, 10.0mmol) and  
10 pyridine-3-carboxaldehyde (1.12g, 10.5mmol) were dissolved  
in anhydrous methanol (35ml) and the solution was heated at  
55°C for three hours. The solution was cooled and  
concentrated in vacuo to a residual oil, which was  
maintained under a 90 $\mu\text{m}$  Hg vacuum for twenty-four hours. N-  
15 (4-(2,4-Difluorophenoxy)phenyl)pyrid-3-ylmethyleamine was  
obtained in quantitative yield.  $^1\text{H}$  NMR (400MHz): 6.83-  
6.90 (1H, m), 6.93-7.00 (3H, m), 7.05-7.13 (1H, m), 7.21-  
7.27 (2H, m), 7.40 (1H, dd,  $J=8.1, 4.9$ ), 8.27 (1H, dd,  $J=8.1, 1.7$ ),  
8.50 (1H, s), 8.69 (1H, dd,  $J=4.9, 1.7$ ), 9.00 (1H, d,  $J=1.7$ );  
20 MS (ESI)  $m/z$  311.1 ( $[\text{M}+\text{H}]^+$ , 100); Anal. Calcd. for  $\text{C}_{18}\text{H}_{12}\text{F}_2\text{N}_2\text{O}$   
C, 69.67; H, 3.90; N, 9.03. Found: C, 69.28; H, 3.99; N, 8.91.

N-(4-(2,4-Difluorophenoxy)phenyl)pyrid-3-  
ylmethyleamine (205mg, 0.66mmol) was taken up in anhydrous  
methanol (3.0ml) and was treated, at room temperature, with  
25 solid sodium borohydride (50mg, 1.32mmol) in one portion.  
There was an initial foaming and mild exotherm to 29 C,  
after which the mixture was stirred under ambient conditions  
for four hours. Aqueous 1N sodium hydroxide was added and  
the mixture extracted four times with methylene chloride.

-77-

The combined organics were washed once with brine, dried over magnesium sulfate and the solvent removed in vacuo to provide N-4-(2,4-difluorophenoxy)phenylpyrid-3-ylmethanamine (200mg, 97%) as a colorless oil which crystallized on standing. <sup>1</sup>H NMR(400MHz): 4.34(2H,s), 6.56-6.62(2H,m), 6.73-6.80(1H,m), 6.82-6.87(2H,m), 6.87-6.94(2H,m), 7.28(1H,dd,J=7.8,2.9), 7.70(1H,ddd,J=7.8,1.5,1.5).

N-4-(2,4-Difluorophenoxy)phenylpyrid-3-ylmethanamine (184mg, 0.59mmol) was stirred in anhydrous acetonitrile (2.0ml) under a nitrogen cover. The solution was treated, via syringe, with anhydrous triethylamine (99μl, 0.71mmol) and then cooled to -8°C. Trifluoroethanesulfonyl chloride (87μl, 0.71mmol) was added neat, dropwise via syringe, at such a rate that the reaction temperature did not exceed -3°C. The mixture was stirred for two hours at -8-(-2)°C and then quenched with 1M potassium carbonate (3.0ml). The upper layer was separated and the aqueous portion extracted three times with ethyl acetate. The combined organics were washed once with saturated sodium chloride, dried over magnesium sulfate and concentrated in vacuo to provide the crude product as a yellow solid. Radial silica gel chromatography (2mm Chromatotron® plate, eluting with 1/1 hexanes/ethyl acetate) provided the title compound (193mg, 71%) as an off-white solid. <sup>1</sup>H NMR(400MHz): 3.81(2H,q,J=9.0), 4.85(2H,s), 6.83-6.92(3H,m), 6.92-6.99(1H,m), 7.06-7.16(3H,m), 7.24-7.31(1H,m), 7.70(1H,ddd,J=9.5,1.7,1.7), 8.33(1H,br s), 8.53(1H,br s); MS(ESI)m/z 454.1([M+H]<sup>+</sup>,100); Anal. Calcd. for C<sub>20</sub>H<sub>15</sub>F<sub>2</sub>N<sub>2</sub>O<sub>3</sub>S, 52.40; H,3.30; N,6.11. Found: C,52.13; H,3.28; N,5.99.


-78-

Preparation 6(4-Bromophenyl)pyridin-3-ylmethylaniline

A solution of 4-bromoaniline (Aldrich; 1.3 equiv.) in methanol (0.4 M) was treated with 3-pyridinecarboxaldehyde (Aldrich; 1.0 equiv.). The solution was heated to reflux for 2 hours. The reaction was cooled to 0°C and treated slowly with sodium borohydride (3.4 equiv.). After 1 hour the cooling bath was removed and stirring continued overnight. The reaction was concentrated in vacuo. The residue was partitioned between DCM and water. The aqueous phase was back-extracted once with DCM. The combined organics were washed with water and brine. The organic layer was concentrated to a white solid, which was purified via recrystallization from 1.2:4 ethyl acetate/hexanes to give the title compound. Anal Calcd for  $C_{12}H_{11}BrN_2 \cdot 0.1H_2O$ : C, 54.40; H, 4.26; N, 10.57. Found: C, 54.30; H, 4.12; N, 10.44. MS found 262.84, 264.83 [M+H]<sup>+</sup>.

Example 342

N-(4-(2-Methoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline



A solution of N-N-(4-bromophenyl)pyridin-3-ylmethylaniline (1.0 equiv.) in pyridine (0.9 M) was treated with the guaiacol (Aldrich) (3.0 equiv.) and anhydrous potassium carbonate (1.5 equiv.). The reaction was warmed to 90°C then treated with cupric oxide. The mixture was heated to reflux for 48 hours. The reaction was cooled


-79-

and diluted with ethyl acetate. The mixture was extracted with 10% aqueous ethylenediamine (3-6 times), 0.1 M potassium carbonate, and brine. The organic layer was dried over sodium sulfate, filtered, and concentrated. The reaction mixture was purified via preparative HPLC eluting with 70:30 to 60:40 DCM/ethyl acetate to give N-(4-(2-methoxyphenoxy)phenyl)pyridin-3-ylmethylamine. <sup>1</sup>H NMR (400 MHz) 3.88 (3H, s), 4.33 (2H, s), 6.60 (2H, m), 6.79-6.89 (4H, m), 6.95-7.04 (2H, m), 7.23-7.29 (1H, m), 7.70 (1H, m), 8.53 (1H, m), 8.64 (1H, bs). MS found 307.0 [M+H]<sup>+</sup>

A solution of N-(4-(2-methoxyphenoxy)phenyl)pyridin-3-ylmethylamine (1.0 equiv.) in 2:1 DCE/Pyridine (0.15 M) was cooled to 0°C and treated with 2,2,2-trifluoroethanesulfonyl chloride (1.8 equiv.). The mixture was allowed to warm to ambient temperature and stir overnight. The reaction was treated with anhydrous potassium carbonate, stirred for 40 min, and filtered. The filtrate was purified via preparative HPLC eluting with 98:2 to 96:4 DCM/MeOH to give the title compound. Anal Calcd for C<sub>21</sub>H<sub>19</sub>F<sub>3</sub>N<sub>2</sub>O<sub>4</sub>S: C, 55.75; H, 4.23; N, 6.19. Found: C, 55.65; H, 4.27; N, 6.19. MS found 453.0 [M+H]<sup>+</sup>

#### Example 343

N-(4-(3-Benzyloxyphenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylamine



A solution of an 3-benzyloxyaniline (1.2 equiv.) in methanol (0.4 M) was treated with 3-pyridinecarboxaldehyde (1.0 equiv.). The solution was heated to reflux for 2.5

-80-


hours. The reaction was cooled to 0°C and treated with sodium borohydride (2.5 equiv.) in 3 portions. The cooling bath was removed and stirring continued for 2 hours. The reaction was concentrated in vacuo. The residue was taken  
5 up in DCM and filtered through celite. The filtrate was concentrated in vacuo and purified via preparative HPLC eluting with 85:15 to 70:30 DCM/ethyl acetate to give N-(4-(3-benzyloxyphenyl)pyridin-3-yl)methylamine. Anal Calcd for  $C_{19}H_{18}N_2O \cdot 0.1H_2O$ : C, 78.11; H, 6.28; N, 9.59. Found: C,  
10 77.86; H, 6.21; N, 9.56. MS found 291.1 (M + H)+

A solution of N-(4-(3-benzyloxyphenyl)pyridin-3-yl)methylamine (1.0 equiv.) in 2:1 DCE/Pyridine (0.15 M) was cooled to 0°C and treated with 2,2,2-trifluoroethanesulfonyl chloride (1.8 equiv.). The mixture was allowed to warm to  
15 ambient temperature and stir overnight. The reaction was treated with anhydrous potassium carbonate, stirred for 40 min, and filtered. The filtrate was purified via preparative HPLC eluting with 70:30 to 60:40 DCM/ethyl acetate, evaporated, and then triturated with diethyl ether  
20 to give the title compound. Anal Calcd for  $C_{21}H_{19}F_3N_2O_3S$ : C, 57.79; H, 4.39; N, 6.42. Found: C, 57.60; H, 4.47; N, 6.41.  $^1H$  NMR (400 MHz) 3.77 (2H, q, J=9.0 Hz), 4.86 (2H, d), 4.99 (2H, d), 6.82 (2H, m), 6.95 (1H, m), 7.25 (2H, m), 7.34-7.40 (5H, m), 7.67 (1H, td, J=2.0, 8.0 Hz), 8.32 (1H, bs), 8.50  
25 (1H, d, J=4.0 Hz). MS found 437.1 [M+H]+

-81-

Example 344


N-(4-(3-Hydroxyphenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylamine



- 5 To a mixture of 2 N-4-(3-benzyloxyphenoxy)phenyl-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylamine in ethanol (0.1 M) in a well-purged par bottle was added Raney® Nickel. The catalyst was rinsed into the mixture with ethyl acetate. The reaction was placed under 37 psi of
- 10 hydrogen gas for 5 hours. The contents were filtered through celite and concentrated in vacuo to give the title compound. Anal Calcd for  $C_{14}H_{13}F_3N_2O_3S \cdot 0.3H_2O$ : C, 47.81; H, 3.90; N, 7.96. Found: C, 47.81; H, 3.80; N, 7.86.  $^1H$  NMR (400 MHz) 3.84 (2H, q,  $J=9.1$  Hz), 4.88 (2H, s), 6.68 (1H, m), 6.72-6.78 (2H, m), 7.19 (1H, t,  $J=8.2$  Hz), 7.34 (1H, dd,  $J=5.1, 7.9$  Hz), 7.75 (1H, d,  $J=7.6$  Hz), 8.30 (1H, s), 8.46 (1H, d,  $J=3.9$  Hz). MS found 347  $[M]^+$
- 15

Example 345

- 20 N-(4-t-Butyldimethylsilyloxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine



To a mixture of 4-aminophenol (Aldrich; 1.0 equiv.) in benzene (1 M) was added 3-pyridenecarboxaldehyde (Aldrich;

-82-

1.0 equiv.). The flask was fitted with a Dean-Stark trap and heated to reflux overnight. (A solid mass formed on the bottom of the flask, which was difficult to break up). The reaction was cooled to ambient and concentrated in vacuo.

5 The solid was taken up in methanol (0.2 M), cooled to 0°C, and treated with sodium borohydride (1.5 equiv.) in 4 portions. The cooling bath was removed and stirring continued for 2 hours. A small amount of was added to quench unreacted hydride. The reaction was concentrated in  
10 vacuo. The residue was partitioned between ethyl acetate and water. The aqueous layer was back-extracted with ethyl acetate. The combined organics were washed with brine, dried over sodium sulfate, filtered, and concentrated. The crude solid was recrystallized from hexanes/ethyl acetate to  
15 give N-(4-hydroxyphenyl)pyridin-3-ylmethylaniline. <sup>1</sup>H NMR (400 MHz, DMSO d<sub>6</sub>) 4.16 (2H, s), 6.41 (2H, m), 6.48 (2H, m), 7.28 (1H, ddd, J=0.73, 4.8, 7.7 Hz), 7.7 (1H, td, J=2.2, 7.3 Hz), 8.38 (1H, m), 8.39 (1H, s), 8.52 (1H, d, J=1.5 Hz).

Alternately, a solution of 4-aminophenol (1.0 equiv.)  
20 (Aldrich) in 2:1 methanol/ethanol was treated with 3-pyridinecarboxaldehyde (1.05 equiv.). The solution was allowed to stir for 1 hour. The reaction was cooled to 0°C and treated with sodium borohydride (4.2 equiv.) in 5 portions. The cooling bath was removed and stirring  
25 continued for 20 hours. The reaction mixture was concentrated in vacuo. The residue was mixed with 1M HCl, warmed, and partitioned between ethyl acetate and 0.1M K<sub>2</sub>CO<sub>3</sub>. The pH of the aqueous phase was adjusted to pH=7 (using 1M HCl and 1M K<sub>2</sub>CO<sub>3</sub>) and the product extracted 4-6  
30 times with ethyl acetate. The organic phases were combined, washed with brine, dried over anhydrous MgSO<sub>4</sub>, and concentrated to afford N-(4-hydroxyphenyl)pyridin-3-ylmethylaniline. <sup>1</sup>H NMR(400 MHz, DMSO-d<sub>6</sub>): d 4.18(2H, s),

-83-

5.65 (1H, s), 6.41-6.53 (4H, m), 7.29-7.32 (1H, m), 7.70-7.72 (1H, m), 8.39-8.40 (2H, m), 8.53-8.54 (1H, m).


MS Found 201.2 [M+H]<sup>+</sup>.

A solution of N-(4-hydroxyphenyl)pyridin-3-ylmethanamine (1.0 equiv.) in THF (0.1 M) was cooled to 0°C and treated with imidazole (1.0 equiv.) and t-butyldimethylsilyl chloride (1.1 equiv.). The reaction was warmed to ambient temperature and stirred overnight. The THF was removed in vacuo. The residue was dissolved in ethyl acetate and washed with 0.1 N HCl, saturated aqueous NaHCO<sub>3</sub>, water, and brine. The residue was purified via preparative HPLC eluting with 80:20 to 68:32 DCM/ethyl acetate to give N-(4-(t-butyldimethylsilyloxy)phenyl)pyridin-3-ylmethanamine. Anal Calcd for C<sub>18</sub>H<sub>26</sub>N<sub>2</sub>OSi: C, 68.74; H, 8.33; N, 8.91. Found: C, 68.62; H, 8.28; N, 8.74. <sup>1</sup>H NMR (400 MHz) 0.15 (6H, s), 0.96 (9H, s), 3.77 (1H, bs), 4.30 (2H, s), 6.53 (2H, m), 6.70 (2H, m), 7.26 (1H, m), 7.70 (1H, td, J=1.9, 7.7 Hz), 8.53 (1H, dd, J=1.4, 4.6 Hz), 8.62 (1H, d, J=1.6 Hz). MS found 315.2 [M+H]<sup>+</sup>.

Following the procedure of Example 342 using N-(4-(t-butyldimethylsilyloxy)phenyl)pyridin-3-ylmethanamine and ethanesulfonyl chloride (Aldrich) and purifying via preparative HPLC eluting with 90:10 to 82:18 DCM/ethyl acetate gave the title compound. Anal Calcd for C<sub>20</sub>H<sub>30</sub>N<sub>2</sub>O<sub>3</sub>SSi: C, 59.08; H, 7.43; N, 6.89. Found: C, 58.87; H, 7.11; N, 6.85. MS found 407.2 [M+H]<sup>+</sup>.




-84-

Example 346N-(4-Hydroxyphenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline

A mixture of N-((4-t-butyldimethylsilyloxy)phenyl)-N-  
5 (ethanesulfonyl)pyrid-3-ylmethylaniline 4-aminophenol was  
added THF (0.5 M), water (0.5 M), and acetic acid (0.16 M)  
was heated to reflux for 3 days. The reaction was cooled to  
ambient temperature and concentrated in vacuo. The  
resultant solid was recrystallized from ethyl acetate to  
10 give the title compound. Anal Calcd for  $C_{14}H_{16}N_2O_3S$ : C, 57.52;  
H, 5.52; N, 9.58. Found: C, 57.25; H, 5.50; N, 9.48. MS  
found 293.1  $[M+H]^+$

Example 347

15 N-(4-t-Butyldimethylsilyloxy)phenyl)-N-(2,2,2-trifluoro  
ethanesulfonyl)pyrid-3-ylmethylaniline



Following the procedure of Example 342 using N-(4-(t-  
butyldimethylsilyloxy)phenyl)pyridin-3-ylmethylaniline and  
20 2,2,2-trifluoroethanesulfonyl chloride (Aldrich) and  
purifying via preparative HPLC eluting with 92:8 to 86:14  
DCM/ethyl acetate gave the title compound. Anal Calcd for  
 $C_{20}H_{27}F_3N_2O_3SSi$ : C, 52.15; H, 5.91; N, 6.08. Found: C, 52.10;  
H, 5.80; N, 6.01.  $^1H$  NMR (400 MHz) 0.18 (6H, s), 0.96 (9H,


-85-

s), 3.80 (2H, q, J=9.2 Hz), 4.83, (2H, s), 6.77 (2H, m), 7.04 (2H, m), 7.24 (1H, dd, J=4.5, 7.4 Hz), 7.66 (1H, td, J=1.8, 8.0 Hz), 8.33 (1H, d, J=1.8 Hz), 8.51 (1H, dd, J=1.5, 4.8 Hz). MS found 461.1 [M+H]<sup>+</sup>

5

Example 348

N-(4-Hydroxyphenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylamine




10 To N-(4-t-butyldimethylsilyloxy)phenyl-N-(2,2,2-trifluoro ethanesulfonyl)pyrid-3-ylmethylamine was added THF (0.5 M), water (0.5 M), and acetic acid (0.16 M). The solution was heated to reflux for 3 days. The reaction was cooled to ambient temperature and concentrated in vacuo.

15 The resultant solid was recrystallized from ethyl acetate to give the title compound. <sup>1</sup>H NMR (400 MHz) 3.82 (2H, q, J=8.8 Hz), 4.84 (2H, s), 6.73 (2H, m), 6.99 (2H, m), 7.43 (1H, dd, J=5.4, 7.3 Hz), 7.91 (1H, m), 8.22 (1H, bs), 8.52 (1H, m).

20

Example 349

N-(4-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylamine



-86-


Using the method of Example 343 using 4-phenoxyaniline (Aldrich) and 3-pyridinecarboxaldehyde (Aldrich) and purifying via preparative HPLC eluting with 95:5 to 60:40 DCM/ethyl acetate gave N-(4-phenoxyphenyl)pyridin-3-

5 ylmethylamine. Anal Calcd for  $C_{18}H_{16}N_2O$ : C, 78.24; H, 5.84; N, 10.14. Found: C, 78.19; H, 5.67; N, 10.03.  $^1H$  NMR (400 MHz, DMSO) 4.00 (1H, bt), 4.35 (2H, d,  $J=5.1$  Hz), 6.62 (2H, m), 6.89-6.93 (4H, m), 6.99-7.03 (1H, m), 7.25-7.30 (3H, m), 7.71 (1H, m), 8.54 (1H, dd,  $J=1.8, 4.8$  Hz), 8.64 (1H, d,  $J=1.8$  Hz). MS found 277.0  $[M+H]^+$

Using the method of Example 343 using N-(4-phenoxyphenyl)pyridin-3-ylmethylamine and 2,2,2-trifluoroethanesulfonyl chloride (Aldrich) and purifying via preparative HPLC eluting with 90:10 to 70:30 DCM/ethyl acetate gave the title compound. Anal Calcd for  $C_{20}H_{17}F_3N_2O_3S$ : C, 56.87; H, 4.06; N, 6.63. Found: C, 56.76; H, 4.09; N, 6.69. MS found 423.1  $[M+H]^+$

#### Example 350

20 N-(4-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)-1-oxypyrid-3-ylmethylamine



A solution of N-(4-(phenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)-1-oxypyrid-3-ylmethylamine (1.0 equiv.) in DCM (0.1 M) was treated with mCPBA (55%; 1.1 equiv.; Lancaster). The resulting mixture was stirred at ambient temperature for 3 hours. The reaction was diluted with DCM, washed with saturated aqueous  $NaHCO_3$ , and brine. The organic layer was dried over sodium sulfate, filtered,


-87-

and concentrated. The residue was purified via preparative HPLC eluting with 96:4 DCM/MeOH to give the title compound. Anal Calcd for  $C_{20}H_{17}F_3N_2O_4S$ : C, 54.79; H, 3.91; N, 6.39. Found: C, 54.68; H, 3.78; N, 6.30. MS found 439.0  $[M+H]^+$

5

Example 351

N-(4-(2-Cyanophenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylamine




10 A solution of N-(4-hydroxyphenoxyphenyl)pyridin-3-ylmethylamine (1.0 equiv.) in DMF (0.15 M) was treated with sodium hydride (1.2 equiv.; as a 60% dispersion in oil). After stirring for 40 min., 2-fluorobenzonitrile (Aldrich; 1.7 equiv.) was added. After 2 hours the reddish reaction  
15 was diluted with ethyl acetate and washed with 0.25 M phosphate buffer, water, and brine. The organic phase was dried over  $Na_2SO_4$ , filtered, and concentrated. The oil was purified via radial chromatography eluting with 70:30 to 50:50 DCM/ethyl acetate to give N-(4-(2-  
20 cyanophenoxy)phenyl)pyridin-3-ylmethylamine. Anal Calcd for  $C_{19}H_{15}N_3O \cdot 0.2H_2O$ : C, 74.83; H, 5.09; N, 13.78. Found: C, 74.86; H, 4.99; N, 13.97.  $^1H$  NMR (400 MHz) 4.10 (1H, bs), 4.38 (2H, s), 6.64 (2H, m), 6.77 (1H, d,  $J=8.3$  Hz), 6.94 (2H, m), 7.04 (1H, t,  $J=7.3$  Hz), 7.30 (1H, dd,  $J=4.9$ , 7.8 Hz), 7.41 (1H, m), 7.61 (1H, dd,  $J=1.5$ , 7.3 Hz), 7.72 (1H, d,  $J=7.8$  Hz), 8.55 (1H, d,  $J=3.4$  Hz), 8.65 (1H, d,  $J=1.5$  Hz). MS found 302.1  $[M+H]^+$   
25

-88-

Using the method of Example 342 using N-(4-(2-cyanophenoxy)phenyl)pyridin-3-ylmethylamine and 2,2,2-trifluoroethanesulfonyl chloride (Aldrich) and purifying via preparative HPLC eluting with 90:10 to 78:22 DCM/ethyl acetate gave the title compound. Anal Calcd for  $C_{21}H_{16}F_3N_3O_3S \cdot 0.2H_2O$ : C, 55.92; H, 3.67; N, 9.32. Found: C, 55.77; H, 3.51; N, 9.18. MS found 447.8  $[M+H]^+$

Example 352

10 N-(4-(2-Cyanophenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine




Using the method of Example 342 using N-(4-(2-cyanophenoxy)phenyl)pyridin-3-ylmethylamine and ethanesulfonyl chloride (Aldrich) and purifying via preparative HPLC eluting with 98:2 to 96:4 DCM/MeOH. Anal Calcd for  $C_{21}H_{19}N_3O_3S \cdot 0.2H_2O$ : C, 63.52; H, 4.92; N, 10.58. Found: C, 63.51; H, 4.79; N, 10.55. MS found 394.0  $[M+H]^+$

-89-

Example 353

N-(3-(2-Methoxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylamine



5           A solution of 3-bromoaniline (Aldrich; 1.3 equiv.) in methanol (0.4 M) was treated with 3-pyridinecarboxaldehyde (Aldrich; 1.0 equiv.). The solution was heated to reflux for 2 hours. The reaction was cooled to 0°C and treated slowly with sodium borohydride (3.4 equiv.). After 1 hour  
10 the cooling bath was removed and stirring continued overnight. The reaction was concentrated in vacuo. The residue was partitioned between DCM and water. The aqueous phase was back-extracted once with DCM. The combined organics were washed with water and brine. The organic  
15 layer was concentrated to a white solid, which was purified via column chromatography eluting with 90:10 DCM/ethyl acetate to give N-(3-bromophenyl)pyridin-3-ylmethylamine. Anal Calcd for  $C_{12}H_{11}BrN_2$ : C, 54.77; H, 4.21; N, 10.64. Found: C, 54.67; H, 4.19; N, 10.57. MS found 262.99, 264.99  
20 [M+H]<sup>+</sup>

          Using the method of Example 342 using N-(3-bromophenyl)pyridin-3-ylmethylamine and guaiacol (Aldrich) and purifying via preparative HPLC eluting with 1:1 DCM/ethyl acetate gave N-(3-(2-Methoxyphenoxy)phenyl) pyrid-  
25 3-ylmethylamine. Anal Calcd for  $C_{19}H_{18}N_2O_2 \cdot 0.5H_2O$ : C, 72.36; H, 6.07; N, 8.88. Found: C, 72.43; H, 5.80; N, 8.96. MS found 307.2 [M+H]<sup>+</sup>

          Using the method of Example 342 using N-(3-(2-methoxyphenoxy)phenyl)pyridin-3-ylmethylamine and ethanesulfonyl


-90-

chloride (Aldrich) and purifying via preparative HPLC eluting with 98:2 to 96:4 DCM/MeOH gave the title compound. Anal Calcd for  $C_{21}H_{22}N_2O_4S$ : C, 63.30; H, 5.56; N, 7.03. Found: C, 63.16; H, 5.51; N, 6.64. MS found 398.9  $[M+H]^+$

5

Example 354


N-(3-(2-phenylphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethamphetamine




10 Using the procedure of Example 342 using N-(3-bromophenyl)pyridin-3-ylmethamphetamine and 2-phenylphenol (Aldrich) and purifying via preparative HPLC eluting with 90:10 to 80:20 DCM/ethyl acetate gave N-(3-(2-phenylphenoxy)phenyl)pyrid-3-ylmethamphetamine. Anal Calcd for  
15  $C_{24}H_{20}N_2O \cdot 0.2H_2O$ : C, 80.96; H, 5.78; N, 7.87. Found: C, 81.07; H, 5.73; N, 7.63. MS found 353.1  $[M+H]^+$

Using the procedure of Example 342 using N-(3-(2-phenylphenoxy)phenyl)pyrid-3-ylmethamphetamine and ethanesulfonyl chloride (Aldrich) and purifying via  
20 preparative HPLC eluting with 99:1 to 97:3 DCM/MeOH gave the title compound. Anal Calcd for  $C_{26}H_{24}N_2O_3S$ : C, 70.25; H, 5.44; N, 6.30. Found: C, 70.13; H, 5.39; N, 6.26. MS found 445.1  $[M+H]^+$

- 91 -

Example 355N-(4-Bromophenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline

Using the procedure of Example 343 using N-(4-bromophenyl)pyridin-3-ylmethylaniline and ethanesulfonyl chloride (Aldrich) and purifying via preparative HPLC eluting with 90:10 to 70:30 DCM/ethyl acetate gave the title compound. Anal Calcd for  $C_{14}H_{15}BrN_2O_2S$ : C, 47.33; H, 4.26; N, 7.88. Found: C, 47.24; H, 4.23; N, 7.79. MS found 354.0, 355.9 FD.

Example 356N-(4-(2-Fluorophenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline


Using the method of Example 342 using N-(4-bromophenyl)pyridin-3-ylmethylaniline and 2-fluorophenol (Aldrich) and purifying via preparative HPLC eluting with 90:10 to 80:20 DCM/ethyl acetate gave N-(4-(2-fluorophenoxy)phenyl)pyrid-3-ylmethylaniline. Anal Calcd for  $C_{18}H_{15}FN_2O$ : C, 73.45; H, 5.14; N, 9.52. Found: C, 73.36; H, 5.22; N, 9.35. MS found 295.1  $[M+H]^+$



- 92 -

Using the method of Example 342 using N-(4-(2-fluorophenoxy)phenyl)pyrid-3-ylmethylaniline and ethanesulfonyl chloride (Aldrich) and purifying via preparative HPLC eluting with 99:1 to 97:3 DCM/MeOH gave the title compound. Anal Calcd for  $C_{20}H_{19}FN_2O_3S \cdot 0.2H_2O$ : C, 61.59; H, 5.01; N, 7.18. Found: C, 61.65; H, 4.85; N, 7.14. MS found 387.1  $[M+H]^+$

By the method of Example 356 the following compounds were prepared and purified by preparative HPLC:




No.	R <sub>1'</sub>	R <sub>c</sub>	Data
357	H	3-fluoro	Anal Calcd for $C_{18}H_{15}FN_2O \cdot 0.4H_2O$ : C, 71.70; H, 5.28; N, 9.29. Found: C, 71.87; H, 5.04; N, 9.47. MS found 295.1 $[M+H]^+$
358	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-fluoro	Anal Calcd for $C_{20}H_{19}FN_2O_3S$ : C, 62.16; H, 4.96; N, 7.25. Found: C, 61.91; H, 4.95; N, 7.24. MS found 387.1 $[M+H]^+$
359	H	4-fluoro	Anal Calcd for $C_{18}H_{15}FN_2O$ : C, 73.45; H, 5.14; N, 9.52. Found: C, 73.23; H, 5.10; N, 9.32. MS found 295.1 $[M+H]^+$
360	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-fluoro	Anal Calcd for $C_{20}H_{19}FN_2O_3S$ : C, 62.16; H, 4.96; N, 7.25. Found: C, 62.18; H, 4.98; N, 7.15. MS found 387.2 $[M+H]^+$

- 93 -


Example 361

N-(4-(2,4-Difluorophenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine



5 Using the method of Example 342 using N-(4-Bromophenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine and 2,4-difluorophenol and purifying via reverse phase chromatography affording the HCl salt. Anal Calcd for  $C_{20}H_{18}F_2N_2O_3S \cdot 1.2HCl$ : C, 53.60; H, 4.32; N, 6.25. Found: C, 53.55; H, 4.18; N, 6.23. MS found 405.1 [M+H]<sup>+</sup>

By the method of Example 342 the following compounds were prepared and purified via reverse phase chromatography affording the HCl salt, except where noted:



No.	Rc	Data
362	2-trifluoromethoxy	Anal Calcd for $C_{21}H_{19}F_3N_2O_4S \cdot 1.2HCl$ : C, 50.83; H, 4.10; N, 5.65. Found: C, 50.66; H, 3.79; N, 5.62. MS found 453.0 [M+H] <sup>+</sup>
363	4-trifluoromethoxy	Anal Calcd for $C_{21}H_{19}F_3N_2O_4S$ : C, 55.75; H, 4.23; N, 6.19. Found: C, 55.60; H, 4.26; N, 6.16. MS found 453.1 [M+H] <sup>+</sup> (analysis on free-base)
364	4-fluoro-2-methyl	Anal Calcd for $C_{21}H_{21}FN_2O_3S$ : C, 62.98; H, 5.28; N, 7.00. Found: C, 62.76; H, 5.28; N, 6.95. MS found 401.1 [M+H] <sup>+</sup> (analysis on free-base)
365	2,4-dimethyl	Anal Calcd for $C_{22}H_{24}N_2O_3S \cdot H_2O \cdot HCl$ : C, 58.59; H, 6.03; N, 6.21. Found: C, 58.34; H, 5.67; N, 6.23. MS found


		397.0 [M+H] <sup>+</sup>
366	4-methyl	Anal Calcd for C <sub>21</sub> H <sub>22</sub> N <sub>2</sub> O <sub>3</sub> S•0.5H <sub>2</sub> O: C, 64.43; H, 5.92; N, 7.16. Found: C, 64.19; H, 5.52; N, 7.03. HRMS calculated 383.1429 [M+H] <sup>+</sup> Found 383.1428 (analysis on free-base)
367	4-isopropyl	Anal Calcd for C <sub>23</sub> H <sub>26</sub> N <sub>2</sub> O <sub>3</sub> S•0.1H <sub>2</sub> O: C, 67.00; H, 6.40; N, 6.79. Found: C, 66.87; H, 6.27; N, 6.78. MS found 411.1 [M+H] <sup>+</sup> (analysis on free-base)
368	4-t-butyl	Anal Calcd for C <sub>24</sub> H <sub>28</sub> N <sub>2</sub> O <sub>3</sub> S: C, 67.90; H, 6.65; N, 6.60. Found: C, 68.02; H, 6.63; N, 6.57. MS found 425.1 [M+H] <sup>+</sup> (analysis on free-base)
369	2-methyl	Anal Calcd for C <sub>21</sub> H <sub>22</sub> N <sub>2</sub> O <sub>3</sub> S•0.3H <sub>2</sub> O: C, 65.03; H, 5.87; N, 7.22. Found: C, 64.94; H, 5.73; N, 7.14. MS found 383.1 [M+H] <sup>+</sup> (analysis on free-base)
370	2-C(O)NH <sub>2</sub>	Anal Calcd for C <sub>21</sub> H <sub>21</sub> N <sub>3</sub> O <sub>4</sub> S: C, 61.30; H, 5.14; N, 10.21. Found: C, 61.16; H, 4.89; N, 9.95. MS found 412.1 [M+H] <sup>+</sup> (analysis on free-base)
372	2-S(O)CH <sub>3</sub>	Anal Calcd for C <sub>21</sub> H <sub>22</sub> N <sub>2</sub> O <sub>4</sub> S <sub>2</sub> •0.5H <sub>2</sub> O: C, 57.38; H, 5.27; N, 6.37. Found: C, 57.31; H, 5.32; N, 5.98. MS found 431.1 [M+H] <sup>+</sup> (purified by radial chromatography eluting with 60:40 to 90:10 THF/hexanes)
373	4-t-butyl-2-fluoro	Anal Calcd for C <sub>24</sub> H <sub>27</sub> FN <sub>2</sub> O <sub>3</sub> S•0.2H <sub>2</sub> O: C, 64.61; H, 6.19; N, 6.28. Found: C, 64.28; H, 5.83; N, 6.19. MS found 443.3 [M+H] <sup>+</sup> (purified via radial chromatography eluting with 50:50 to 70:30 THF/hexanes)
374	4-fluoro-2-n-propyl	<sup>1</sup> H NMR (400 MHz) 0.89 (3H, t, J=7.3 Hz), 1.44 (3H, t, J=7.3 Hz), 1.56 (2H, sextet, J=7.8 Hz), 2.48 (2H, t, J=7.8 Hz), 3.10 (2H, q, J=7.3 Hz), 4.85 (2H, s), 6.78 (2H, m), 6.85 (2H, m), 6.94 (1H, m), 7.13 (2H, m), 7.32 (1H, bs), 7.79 (1H, d, J=7.8 Hz), 8.4 (1H, bs), 8.55 (1H, bs). HRMS calculated 429.1648 [M+H] <sup>+</sup> Found 429.1656 (analysis of free-base)
375	4-C(O)OCH <sub>3</sub>	<sup>1</sup> H NMR (400 MHz, DMSO-d <sub>6</sub> ) 1.28 (3H, t, J=7.3 Hz), 3.26 (2H, q, J=7.3 Hz), 3.80 (3H, s), 4.96 (2H, s), 7.00-7.05 (4H, m), 7.40 (2H, m), 7.56 (1H, bs), 7.86 (1H, d, J=7.8 Hz), 7.93 (2H, m), 2 protons alpha to the pyridyl N broadened into the baseline. HRMS calculated 427.1328 [M+H] <sup>+</sup> Found 427.1338
376	4-CH <sub>2</sub> C(O)NH <sub>2</sub>	Anal Calcd for C <sub>22</sub> H <sub>23</sub> N <sub>3</sub> O <sub>4</sub> S: C, 53.40; H, 5.07; N, 8.49. Found: C, 53.50; H, 4.90; N, 8.46. MS found 426.0 [M+H] <sup>+</sup> (analysis on free-base)
377	3-t-butyl	Anal Calcd for C <sub>24</sub> H <sub>28</sub> N <sub>2</sub> O <sub>3</sub> S: C, 67.90; H, 6.65; N, 6.60. Found: C, 68.02; H, 6.34; N, 6.49. MS found 425.1 [M+H] <sup>+</sup> purified by radial chromatography eluting with 50:50 hexanes/THF)
378	2-methoxy	Anal Calcd for C <sub>21</sub> H <sub>22</sub> N <sub>2</sub> O <sub>4</sub> S: C, 63.30; H, 5.56; N, 7.03. Found: C, 63.25; H, 5.59; N, 7.02. MS found 399.1 [M+H] <sup>+</sup> (purified via radial chromatography eluting with 50:50

- 95 -

		hexanes/THF)
379	3-methyl	Anal Calcd for $C_{21}H_{22}N_2O_3S$ : C, 65.95; H, 5.80; N, 7.32. Found: C, 65.55; H, 5.64; N, 7.21. MS found 383.1 [M+H] <sup>+</sup> (analysis on free-base)
380	2-chloro	Anal Calcd for $C_{20}H_{19}ClN_2O_3S$ : C, 59.62; H, 4.75; N, 6.95. Found: C, 59.63; H, 4.61; N, 6.88. MS found 403.1 [M+H] <sup>+</sup> (purified via radial chromatography eluting with 60:40 to 50:50 hexanes/THF)
381	4-C(O)HN <sub>2</sub>	Anal Calcd for $C_{21}H_{21}N_3O_4S \cdot 1.1HCl \cdot 1H_2O$ : C, 53.71; H, 5.17; N, 8.95. Found: C, 53.65; H, 4.79; N, 8.39. MS found 412.1 [M+H] <sup>+</sup>
382	2-methyl	Anal Calcd for $C_{21}H_{22}N_2O_3S \cdot 0.3H_2O$ : C, 65.03; H, 5.87; N, 7.22. Found: C, 64.94; H, 5.73; N, 7.14. MS found 383.1 [M+H] <sup>+</sup> . (isolated as the free-base)
383	4-S(O) <sub>2</sub> CH <sub>3</sub>	Anal Calcd for $C_{21}H_{22}N_2O_5S_2$ : C, 56.48; H, 4.96; N, 6.27. Found: C, 56.33; H, 5.03; N, 6.20. MS found 447.1 [M+H] <sup>+</sup> (purified via radial chromatography eluting with 50:50 to 70:30 THF/hexanes)
384	2-(prop-2-yl)	Anal Calcd for $C_{23}H_{26}N_2O_3S$ : C, 67.29; H, 6.38; N, 6.82. Found: C, 66.91; H, 6.30; N, 6.68. MS found 411.2 [M+H] <sup>+</sup> (purified via normal phase radial chromatography eluting with 1:1 THF/hexanes)
385	2-ethyl	Anal Calcd for $C_{22}H_{24}N_2O_3S$ : C, 66.64; H, 6.10; N, 7.07. Found: C, 66.49; H, 6.02; N, 6.87. MS found 397.2 [M+H] <sup>+</sup> (purified via flash chromatography eluting with 1:1 ethyl acetate/hexanes to 100% ethyl acetate)
386	4-S(O) <sub>2</sub> NH <sub>2</sub>	Anal Calcd for $C_{20}H_{21}N_3O_5S_2$ : C, 53.68; H, 4.73; N, 9.39. Found: C, 53.29; H, 4.70; N, 9.29. MS found 448.1 [M+H] <sup>+</sup> (purified via normal phase radial chromatography eluting with 5% (methanol/7M ammonia)/dichloromethane)

Example 387

N- (4- (2,4-Difluorophenoxy) phenyl) -N- (ethanesulfonyl) -1-  
pyrid-3-ylethylamine



-96-


A solution of N-(4-(2,4-difluorophenoxy)phenyl)amine (1.0 equiv.) in dichloromethane (0.2 M) was cooled to 0°C and treated with 3-acetylpyridine (Aldrich; 1.4 equiv.), acetic acid (5.0 equiv.), and sodium triacetoxyborohydride (2.5 equiv.). The reaction was allowed to warm to ambient temperature and stirred an additional 22 hours. The reaction was cautiously quenched with saturated aqueous NaHCO<sub>3</sub> then adjusted to pH=10 with 1 N NaOH. The mixture was stirred for 4 hours then the layers were separated. The aqueous layer was extracted twice with DCM. The combined organics were washed with brine, dried over sodium sulfate, filtered, and concentrated. The crude material was purified via radial chromatography eluting with 50:50 to 60:40 ethyl acetate/hexanes to give N-(4-(2,4-difluorophenoxy)phenyl)-1-pyrid-3-ylethylamine. <sup>1</sup>H NMR (400 MHz) 1.54 (3H, d, J=6.7 Hz), 3.96 (1H, bs), 4.48 (1H, q, J=6.7 Hz), 6.44 (2H, m), 6.76 (2H, m), 6.83-6.93 (2H, m), 7.24 (1H, m), 7.44 (1H, m), 7.68 (1H, td, J=1.8, 7.9 Hz), 8.50 (1H, d, J=3.8 Hz), 8.64 (1H, bs). HRMS calcd 327.1309. Found 327.1308.

Using the method of Example 342 using N-(4-(2,4-difluorophenoxy)phenyl)-1-pyrid-3-ylethylamine and 2,2,2-trifluoroethanesulfonyl chloride and purifying via radial chromatography eluting with 95:5 to 85:15 CHCl<sub>3</sub>/ethyl acetate gave the title compound. Anal Calcd for C<sub>21</sub>H<sub>17</sub>F<sub>5</sub>N<sub>2</sub>O<sub>3</sub>S•0.2H<sub>2</sub>O: C, 52.98; H, 3.68; N, 5.89. Found: C, 52.88; H, 3.73; N, 5.67. MS found 473.1 [M+H]<sup>+</sup>

-97-

Example 388

N-((3-Benzyloxyphenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine




5        Using the method of Example 342 using (3-benzyloxy-phenyl)pyridin-3-ylmethylamine and ethanesulfonyl chloride and purifying via preparative HPLC eluting with 90:10 to 75:25 DCM/ethyl acetate gave, after trituration with diethyl ether, the title compound. Anal Calcd for  $C_{21}H_{22}N_2O_3S$ : C, 65.95; H, 5.80; N, 7.32. Found: C, 65.85; H, 5.75; N, 7.38.  
10        MS found 383.1  $[M+H]^+$

Preparation 7

2-(3-Hydroxy-3-methyl)butyl-4-fluorophenol


15        A solution of 6-fluoro-2,2-dimethyl-chroman-4-one (EP 193493; 1 equiv.) in ethanol (0.24 M) was treated with copper chromite (0.6 equiv.). The reaction was heated to 200°C for 6 hours at 3000 psi of hydrogen gas. The reaction was filtered and concentrated to a solid. The crude  
20        material was recrystallized from hot hexanes to give the title compound. MS found 198  $[M+H]^+$

- 98 -

Example 389N- (3- (Bromophenyl) -N- (ethylsulfonyl) pyrid-3-yl)methylamine

Using the method of Example 342 using (3-  
 5 bromophenyl)pyridin-3-ylmethylamine and ethanesulfonyl  
 chloride and purifying via preparative HPLC eluting with  
 90:10 to 70:30 DCM/ethyl acetate gave the title compound.  
 Anal Calcd for  $C_{14}H_{15}BrN_2O_2S$ : C, 47.33; H, 4.26; N, 7.88.  
 Found: C, 47.07; H, 4.05; N, 7.67. MS found 354.9, 356.9  
 10  $[M+H]^+$ .

By the method of Example 342 using N-(3-(bromophenyl)-  
 N-(ethylsulfonyl)pyrid-3-yl)methylamine the following  
 compounds were prepared and purified via reverse phase  
 chromatography and isolated as the free-base, except where  
 15 noted:



No.	Rc	Data
390	2-fluoro	Anal Calcd for $C_{20}H_{19}FN_2O_3S$ : C, 62.16; H, 4.96; N, 7.25. Found: C, 61.85; H, 4.92; N, 7.02. MS found 387.1 $[M+H]^+$
391	3-fluoro	Anal Calcd for $C_{20}H_{19}FN_2O_3S$ : C, 62.16; H, 4.96; N, 7.25. Found: C, 61.84; H, 5.00; N, 7.13. MS found 387.1 $[M+H]^+$
392	4-fluoro	Anal Calcd for $C_{20}H_{19}FN_2O_3S$ : C, 62.16; H, 4.96; N, 7.25. Found: C, 62.02; H, 4.70; N, 7.23. MS found 387.1 $[M+H]^+$
393	2, 4-difluoro	Anal Calcd for $C_{20}H_{17}F_3N_2O_3S$ : C, 56.87; H, 4.06; N, 6.63. Found: C, 56.93; H, 4.00; N, 6.43. MS found 423.0 $[M+H]^+$
394	2-ethoxy	Anal Calcd for $C_{22}H_{24}N_2O_4S$ : C, 64.06; H, 5.86; N, 6.79. Found: C, 64.10; H, 5.61; N, 6.75. MS found 413.0 $[M+H]^+$

395	2-phenyl	Anal Calcd for $C_{26}H_{24}N_2O_3S$ : C, 70.25; H, 5.44; N, 6.30. Found: C, 70.43; H, 5.35; N, 6.18. MS found 445.1 [M+H] <sup>+</sup>
396	4-trifluoromethoxy	Anal Calcd for $C_{21}H_{19}F_3N_2O_4S$ : C, 55.75; H, 4.23; N, 6.19. Found: C, 55.46; H, 4.20; N, 6.10. MS found 453.0 [M+H] <sup>+</sup>
397	4-methyl	<sup>1</sup> H NMR (400 MHz) 1.42 (3H, t, J=7.3 Hz), 2.34 (3H, s), 3.10 (2H, q, J=7.3 Hz), 4.85 (2H, s), 6.83 (4H, m), 6.95 (1H, m), 7.13 (2H, m), 7.23 (2H, m), 7.70 (1H, d, J=7.8 Hz), 8.37 (1H, bs), 8.51 (1H, bs). HRMS calculated 383.1429 [M+H] <sup>+</sup> Found 383.1426.
398	2-methyl-4-fluoro	<sup>1</sup> H NMR (400 MHz) 1.42 (3H, t, J=7.3 Hz), 2.09 (3H, s), 3.11 (2H, q, J=7.3 Hz), 4.84 (2H, s), 6.68-6.95 (6H, m), 7.22 (2H, m), 7.69 (1H, d, J=7.8 Hz), 8.33 (1H, bs), 8.51 (1H, bs). HRMS calculated 401.1335 [M+H] <sup>+</sup> Found 401.1327.
399	2-trifluoromethoxy	Anal Calcd for $C_{21}H_{19}F_3N_2O_4S$ : C, 55.75; H, 4.23; N, 6.19. Found: C, 55.60; H, 4.12; N, 6.05. MS found 453.1 [M+H] <sup>+</sup>
400	4-CH <sub>2</sub> C(O)NH <sub>2</sub>	Anal Calcd for $C_{22}H_{23}N_3O_4S \cdot 0.5H_2O$ : C, 60.81; H, 5.27; N, 9.67. Found: C, 60.69; H, 5.27; N, 9.48. MS found 426.1 [M+H] <sup>+</sup>
401	4-C(O)NH <sub>2</sub>	Anal Calcd for $C_{21}H_{21}N_3O_4S \cdot 2.1HCl \cdot 1.1H_2O$ : C, 49.66; H, 5.02; N, 8.27. Found: C, 49.67; H, 4.92; N, 8.11. MS found 412.1 [M+H] <sup>+</sup> (isolated as the HCl salt)
402	4-S(O) <sub>2</sub> CH <sub>3</sub>	Anal Calcd for $C_{21}H_{22}N_2O_5S_2 \cdot 0.2H_2O$ : C, 56.03; H, 5.02; N, 6.22. Found: C, 56.02; H, 4.78; N, 6.12. MS found 447.1 [M+H] <sup>+</sup> (purified via radial chromatography eluting with 50:50 to 70:30 THF/hexanes)
403	2-S(O)CH <sub>3</sub>	Anal Calcd for $C_{21}H_{22}N_2O_4S_2 \cdot 0.4H_2O$ : C, 57.62; H, 5.25; N, 6.40. Found: C, 57.62; H, 5.25; N, 6.24. HRMS calculated 431.1099 [M+H] <sup>+</sup> Found 431.1114 (purified via radial chromatography eluting with 50:50 to 70:30 THF/hexanes)
404	4-t-butyl	<sup>1</sup> H NMR (400 MHz) 1.33 (9H, s), 1.42 (3H, t, J=7.8 Hz), 3.11 (2H, q, J=7.8 Hz), 4.87 (2H, s), 6.83-6.97 (5H, m), 7.22-7.36 (4H, m), 7.76 (1H, d, J=7.8 Hz), 8.38 (1H, bs), 8.53 (1H, bs). HRMS calculated 425.1899 [M+H] <sup>+</sup> Found 425.1906 (purified via radial chromatography eluting with 50:50 hexanes/THF)
405	3-methyl	Anal Calcd for $C_{21}H_{22}N_2O_3S$ : C, 65.95; H, 5.80; N, 7.32. Found: C, 66.09; H, 5.86; N, 7.29. MS found 383.1 [M+H] <sup>+</sup>
406	2-n-propyl-4-fluoro	Anal Calcd for $C_{23}H_{25}FN_2O_3S$ : C, 64.47; H, 5.88; N, 6.54. Found: C, 64.31; H, 5.79; N, 6.45. MS found 429.1 [M+H] <sup>+</sup>
407	2-chloro	<sup>1</sup> H NMR (400 MHz) 1.42 (3H, t, J=7.3 Hz), 3.11 (2H, q, J=7.3 Hz), 4.86 (2H, s), 6.8-6.92 (3H, m), 7.00 (1H, dd, J=2.0, 7.8 Hz), 7.12 (1H, dt, J=2.0, 7.8 Hz), 7.21-7.28 (3H, m), 7.45 (1H, dd, J=2.0, 8.3 Hz), 7.68 (1H, d, J=7.8 Hz), 8.36 (1H, bs), 8.50 (1H, bs). HRMS calculated 403.0883 [M+H] <sup>+</sup> Found 403.0896 (purified via radial




		chromatography eluting with 60:40 to 50:50 hexanes/THF)
408	2-methyl	<sup>1</sup> H NMR (400 MHz) 1.42 (3H, t, J=7.3 Hz), 2.13 (3H, s), 3.11 (2H, q, J=7.3 Hz), 4.84 (2H, s), 6.74-6.80 (3H, m), 6.93-6.95 (1H, m), 7.07-7.24 (5H, m), 7.68 (1H, td, J=2.0, 7.8 Hz), 8.34 (1H, bs), 8.50 (1H, d, J=3.4 Hz). HRMS calculated 383.1429 [M+H] <sup>+</sup> Found 383.1429 (purified via radial chromatography eluting with 60:40 to 50:50 hexanes/THF)
409	H	Anal Calcd for C <sub>20</sub> H <sub>20</sub> N <sub>2</sub> O <sub>3</sub> S: C, 65.20; H, 5.47; N, 7.60. Found: C, 64.86; H, 5.25; N, 7.26. MS found 369.1 [M+H] <sup>+</sup> (purified via radial chromatography eluting with 60:40 to 50:50 hexanes/THF)
410	4-C(O)OCH <sub>3</sub>	Anal Calcd for C <sub>22</sub> H <sub>22</sub> N <sub>2</sub> O <sub>5</sub> S•1HCl: C, 57.07; H, 5.01; N, 6.05. Found: C, 56.73; H, 5.01; N, 6.01. MS found 427.1 [M+H] <sup>+</sup> (isolated as the HCl salt)
411	3-t-butyl	Anal Calcd for C <sub>24</sub> H <sub>28</sub> N <sub>2</sub> O <sub>3</sub> S: C, 67.90; H, 6.65; N, 6.60. Found: C, 68.17; H, 6.45; N, 6.51. MS found 425.2 [M+H] <sup>+</sup> (purified via radial chromatography eluting with 60:40 to 50:50 hexanes/THF)
412	4-fluoro-2-(3-hydroxy-3-methylbutyl	Anal Calcd for C <sub>25</sub> H <sub>29</sub> FN <sub>2</sub> O <sub>4</sub> S: C, 63.54; H, 6.18; N, 5.93. Found: C, 63.16; H, 6.05; N, 5.84. MS found 473.2 [M+H] <sup>+</sup>
413	2-fluoro-4-t-butyl	Anal Calcd for C <sub>24</sub> H <sub>27</sub> FN <sub>2</sub> O <sub>3</sub> S: C, 65.14; H, 6.15; N, 6.33. Found: C, 65.28; H, 6.24; N, 6.30. MS found 442.1 (FD). (purified via radial chromatography eluting with 50:50 to 70:30 THF/hexanes)
414	4-CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	Anal Calcd for C <sub>23</sub> H <sub>24</sub> N <sub>2</sub> O <sub>5</sub> S•0.5H <sub>2</sub> O: C, 61.45; H, 5.61; N, 6.23. Found: C, 61.43; H, 5.43; N, 6.45. MS found 441.1 [M+H] <sup>+</sup> (purified via radial chromatography eluting with 50:50 to 75:25 THF/hexanes)
415	3-C(O)NH <sub>2</sub>	<sup>1</sup> H NMR (400 MHz) 1.44 (3H, t, J=7.3 Hz), 3.13 (2H, q, J=7.8 Hz), 4.87 (2H, s), 5.7 (1H, bs), 6.74 (1H, bs), 6.86 (1H, t, J=2.0 Hz), 6.96 (1H, m), 7.06 (1H, m), 7.12 (1H, ddd, J=0.98, 2.4, 8.3 Hz), 7.28 (2H, m), 7.33 (1H, t, J=7.8 Hz), 7.43 (1H, t, J=7.8 Hz), 7.60 (1H, m), 7.72 (1H, d, J=7.8 Hz), 8.32 (1H, bs), 8.48 (1H, bs). HRMS calculated 412.1331 [M+H] <sup>+</sup> Found 412.1329 (purified via radial chromatography eluting with 60:40 to 70:30 THF/hexanes)
416	2,4-difluoro	Anal Calcd for C <sub>20</sub> H <sub>18</sub> F <sub>2</sub> N <sub>2</sub> O <sub>3</sub> S•0.1DCM: C, 58.46; H, 4.44; N, 6.78. Found: C, 58.65; H, 4.19; N, 6.59. HRMS calcd 405.1084. Found 405.1083 [M+H] <sup>+</sup>
417	4-CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	Anal Calcd for C <sub>23</sub> H <sub>24</sub> N <sub>2</sub> O <sub>5</sub> S•0.5H <sub>2</sub> O: C, 61.45; H, 5.61; N, 6.23. Found: C, 61.43; H, 5.43; N, 6.45. MS found 441.1 [M+H] <sup>+</sup> (purified via radial chromatography eluting with 50:50 to 75:25 THF/hexanes)
418	2,4,6-trifluoro	Anal Calcd for C <sub>20</sub> H <sub>17</sub> F <sub>3</sub> N <sub>2</sub> O <sub>3</sub> S: C, 56.87; H, 4.06; N, 6.63. Found: C, 56.93; H, 4.00; N, 6.43. MS found 423.0 [M+H] <sup>+</sup>

-101-

419	2-(prop-2-yl)	Anal Calcd for $C_{23}H_{26}N_2O_3S$ : C, 67.29; H, 6.38; N, 6.82. Found: C, 67.37; H, 6.25; N, 6.66. MS found 411.2 $[M+H]^+$ (purified via normal phase radial chromatography eluting with 1:1 THF/hexanes)
420	2-ethyl	Anal Calcd for $C_{22}H_{24}N_2O_3S$ : C, 66.64; H, 6.10; N, 7.07. Found: C, 66.72; H, 6.09; N, 6.88. MS found 397.2 $[M+H]^+$ (purified via flash chromatography eluting with 1:1 ethyl acetate/hexanes to 100% ethyl acetate)
421	4-S(O) <sub>2</sub> NH <sub>2</sub>	Anal Calcd for $C_{20}H_{21}N_3O_5S_2 \cdot HCl$ : C, 49.63; H, 4.58; N, 8.68. Found: C, 49.58; H, 4.42; N, 8.43. MS found 448.1 $[M+H]^+$ (isolate as the HCl salt from reverse phase chromatography)

Example 422

N-(4-(3-Methoxybenzyloxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine



5


A solution of N-(4-(4-hydroxyphenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine (1.0 equiv.) in DMSO (0.15 M) was cooled to 0°C. Sodium hydride (1.1 equiv.; 60% dispersion in oil) was added; after 20 minutes at 0°C 3-methoxybenzyl bromide (Aldrich; 1.3 equiv.) was added and the reaction warmed to ambient temperature. After stirring overnight the reaction was diluted with ethyl acetate and washed with 0.1 M potassium carbonate, and brine. The organic layer was dried over sodium sulfate, filtered, and concentrated in vacuo. The residue was purified via preparative HPLC eluting with 90:10 to 70:30 DCM/ethyl acetate to give the title compound. Anal Calcd for  $C_{22}H_{24}N_2O_4S \cdot 0.1H_2O$ : C, 64.06; H, 5.86; N, 6.79. Found: C, 63.78; H, 5.89; N, 6.76. MS found 413.0  $[M+H]^+$ .

20

-102-

Example 443

N-(3-(2-Fluorophenoxy)phenyl)-N-(2,2,2-  
trifluoroethanesulfonyl)pyrid-3-ylmethylamine




5        Using the method of Example 342 using N-(3-(2-fluorophenoxy)phenyl)pyridin-3-ylmethylamine and 2,2,2-trifluoroethanesulfonyl chloride and purifying via radial chromatography eluting with 40:60 to 50:50 THF/hexanes gave the title compound. Anal Calcd for  $C_{20}H_{16}F_4N_2O_3S$ : C, 54.54; H, 3.66; N, 6.36. Found: C, 54.23; H, 3.40; N, 6.31. MS found 441.2  $[M+H]^+$

10

Example 444

N-(3-(2-Methylphenoxy)phenyl)-N-(2,2,2-  
trifluoroethanesulfonyl)pyrid-3-ylmethylamine

15




Using the method of Example 342 using N-(3-(2-methylphenoxy)phenyl)pyridin-3-ylmethylamine and 2,2,2-trifluoroethanesulfonyl chloride and purifying via radial chromatography eluting with 40:60 to 50:50 THF/hexanes gave the title compound. Anal Calcd for  $C_{21}H_{19}F_3N_2O_3S \cdot 0.1H_2O$ : C, 57.55; H, 4.42; N, 6.39. Found: C, 57.28; H, 4.15; N, 6.33. MS found 437.2  $[M+H]^+$

20

-103-

Example 445

N-(3-(4-Carboxymethylphenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline




5           N-(3-(4-Carbomethoxymethylphenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline (1.0 equiv.) was dissolved in THF (3.3 M) and treated with 1N NaOH (3 equiv.). The reaction was stirred overnight then concentrated in vacuo to remove THF. The pH was adjusted to  
10   4 with HCl, which caused the title compound to precipitate. The solid was collected via filtration washing with pH=4 water to give the title compound. Anal Calcd for  $C_{22}H_{22}N_2O_5S \cdot 0.2H_2O$ : C, 61.44; H, 5.25; N, 6.51. Found: C, 61.39; H, 5.12; N, 6.59. MS found 427.0 [M+H]<sup>+</sup>

15

Example 446

N-(4-(2-Fluoro-4-methylphenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline




20           Using the method of Example 342 using N-(4-hydroxyphenyl)-N-(ethanesulfonyl)pyridin-3-ylmethylaniline and 4-bromo-2-fluorotoluene (Strem) and purifying via preparative HPLC eluting with 90:10 to 70:30 DCM/ethyl

-104-

acetate) gave the title compound. Anal Calcd for  $C_{21}H_{21}FN_2O_3S \cdot 0.2H_2O$ : C, 62.42; H, 5.34; N, 6.93. Found: C, 62.48; H, 5.33; N, 6.72.  $^1H$  NMR (400 MHz) 1.42 (3H, t,  $J=7.3$  Hz), 2.35 (3H, s), 3.10 (2H, q,  $J=7.3$  Hz), 4.82 (2H, s), 6.82-7.00 (5H, m), 7.12 (2H, m), 7.30 (1H, bs), 7.78 (1H, d,  $J=7.8$  Hz), 8.40 (1H, bs), 8.55 (1H, bs).

By a method similar to the method of Example 1 the following compounds were prepared:



No.	R1	R2	Data
447	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-(2-methoxy-phenylsulfanyl)	$^1H$ NMR: 8.30 (d, 1H, $J=3$ Hz), 7.62 (s, 1H), 7.30 (d, 1H, $J=7$ Hz), 7.20 (m, 1H), 7.05-7.18 (m, 6H), 4.79 (s, 2H), 3.79 (s, 3H), 3.00-3.05 (m, 2H), 1.35 (t, 3H, $J=7$ Hz). MS calcd. 414.5; MS (M+1) 416.1.
448	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-(2-methoxy-phenylsulfanyl)	$^1H$ NMR: 8.48 (d, 1H, $J=4$ Hz), 8.26 (s, 1H), 7.60 (m, 1H), 7.32-7.36 (m, 1H), 7.15-7.22 (m, 6H), 6.90-7.00 (m, 2H), 4.78 (s, 2H), 3.69-3.79 (m, 5H). MS calcd. 468.5; MS (M+1) 469.9.
449	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-(2-methoxy-phenoxy)-	$^1H$ NMR: 8.50 (d, 1H, $J=4$ Hz), 8.30 (s, 1H), 7.64 (d, 1H, $J=8$ Hz), 7.14-7.24 (m, 4H), 6.99 (d, 1H, $J=8$ Hz), 6.93 (s, 1H), 6.84-6.86 (m, 2H), 6.76 (s, 1H), 4.82 (s, 2H), 3.75-3.82 (m, 5H). MS calcd. 452.5; MS (M+1) 453.3.
450	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(2-benzyloxy-phenoxy)	$^1H$ NMR: 8.5 (s, 1H), 8.28 (s, 1H), 7.69 (d, 1H, $J=8$ Hz), 7.26 (m, 2H), 7.02-7.12 (m, 4H), 6.84-6.91 (m, 4H), 4.83 (s, 2H), 3.76-3.83 (dd, 2H). MS calcd. 438.4; MS (M+1) 439.4.
451	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-(4-fluoro-2-methoxy-phenoxy)	$^1H$ NMR: 8.5 (dd, 1H), 8.32 (d, 1H, $J=1$ Hz), 7.67 (d, 1H, $J=8$ Hz), 7.16-7.22 (m, 3H), 6.87 (m, 2H), 6.75 (m, 2H), 6.62 (m, 1H), 4.83 (s, 2H), 3.73 (s, 3H), 3.08 (dd, 2H), 1.37-1.41 (t, 3H). MS calcd. 416.5; MS (M+1) 417.6.
452	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-(4-fluoro-2-methoxy-	$^1H$ NMR: 8.49 (d, 1H, $J=4$ Hz), 8.28 (s, 1H), 7.65 (d, 1H, $J=8$ Hz), 7.20-7.22 (m, 3H), 6.80-

		phenoxy)	6.90 (m,2H), 6.70 (m,1H), 6.60 (m,1H), 4.82 (s, 2h), 3.72-3.82 (m,4H). MS calcd. 470.5; MS (M+1)471.6
453	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-(2-bromo - phenoxy)	<sup>1</sup> H NMR: 8.47 (dd,1H), 8.33(d,1H,J=2Hz), 7.65(m,1H), 7.61 (m,1H), 7.19-7.27 (m,4H), 6.96-7.05 (m,3H), 6.78-6.86 (m,3H), 4.84 (s,2H), 3.06-3.10 (m,2H), 1.37-1.41 (t,3H). MS calcd. 447.4, MS (M+1)449.0
454	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-(2,3-dimethoxy-phenoxy)	<sup>1</sup> H NMR: 8.47 (dd,1H), 8.35 (d,1H,J=1Hz), 7.65 (dd,1H), 7.17-7.22 (m,3H), 6.81-7.00 (m,3H), 6.84 (m,2H), 6.72 (dd,1H), 6.47 (dd,1H), 4.83(s, 2H), 3.87 (s,3H), 3.71 (s,3H), 3.05-3.11 (dd,2H), 1.38 (t,3H). MS calcd. 428.5; MS (M+1)429.6
455	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-(2-chloro-phenoxy)	<sup>1</sup> H NMR: 8.47 (dd,1H), 8.33 (d,1H,J=2Hz), 7.65 (d,1H,J=8Hz), 7.42 (dd,1H), 7.19-7.25 (m,4H), 7.11 (dd,1H), 6.97 (dd,1H), 6.86-6.89 (m,1H), 6.82 (d,2H,J=2Hz), 6.78-6.80 (m,2H), 4.84 (s,3H), 3.11 (dd,3H), 1.39 (t,2H). MS calcd. 402.9; MS (M+1)403.8.
456	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-(2-chloro-6-methyl-phenoxy)	<sup>1</sup> H NMR: 8.45 (m,1H), 8.31 (d,1H,J=1Hz), 7.61 (d,1H,J=8Hz), 7.06-7.26 (m,5H), 6.90 (dd,1H), 6.69 (dd,1H), 6.58 (t,1H), 4.81 (s,2H), 3.05-3.10 (dd,2H), 2.07 (s,3H), 1.37 (t,3H). MS calcd. 416.9; MS (M+1)417.7
457	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-(2-isopropoxy-phenoxy)	<sup>1</sup> H NMR: 8.47 (dd,1H), 8.35 (d,1H,J=2Hz), 7.68 (d,1H,J=8Hz), 7.21 (dd,1H), 7.09-7.22 (m,3H), 6.84-6.98 (m,4H), 6.82 (s,1H), 6.74 (dd,1H), 4.84 (s,2H), 4.39 (m,1H), 3.08 (dd,2H), 1.38 (t,3H), 1.12 (d,6H,J=7Hz). MS calcd. 426.5; MS (M+1)427.6
458	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-fluoro-5-(2-methoxy-phenoxy)	<sup>1</sup> H NMR (CD <sub>3</sub> OD): 8.79 (m,2H), 8.56 (d,1H,J=8Hz), 8.05 (dd,1H), 7.22 (t,1H), 7.11 (d,J=7Hz), 6.90-7.03 (m,3H), 6.68 (s,1H), 6.49-6.53 (m,1H), 5.14 (s,2H), 4.41 (dd,2H), 3.70 (s,3H). MS calcd. 470.45; MS (M+1) 471.6
459	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-fluoro-5-(2-methoxy-phenoxy)	<sup>1</sup> H NMR (CD <sub>3</sub> OD): 8.77 (s,2H), 8.57 (s, 1H), 8.04 (s,1H), 7.21 (s,1H), 7.09 (s,1H), 6.95 (m,3H), 6.71 (s,1H), 6.42 (d,1H,J=9Hz), 5.15 (s,2H), 3.69 (s,3H), 3.27 (m,2H), 1.34 (m,3H). MS calcd. 416.48; MS (M+1)417.7
460	SO <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-(2-methoxy-phenoxy)	<sup>1</sup> H NMR (CD <sub>3</sub> OD): 8.53-8.83 (m,3H), 7.90-8.08 (m,1H), 6.74-7.26 (m,8H), 5.13 (s,2H), 3.66 (s,3H), 3.25 (m,2H), 2.65 (m,2H). MS calcd. 466.38; MS (M+1) 467.6
461	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-(2-bromo-	<sup>1</sup> H NMR: 7.63 (m,2H), 7.37 (m,3H), 7.04-

-106-

		phenoxy)	7.19 (m,4H), 6.91 (s,2H), 6.77 (br,1H), 4.10 (dd,1H), 3.85 (s, NH), 2.02 (s,1H), 1.24 (t,2H). MS calcd. 501.1; MS (M+1)501.4 (502.4, due to Br).
462	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-(2,3-dimethoxy-phenoxy)	<sup>1</sup> H NMR (CD <sub>3</sub> OD): 8.69-8.77 (m,2H), 8.51 (s,1H), 8.00 (s,1H), 7.33 (s,1H), 7.03-7.11 (m,2H), 6.89 (m,2H), 6.69 (s,1H), 6.56 (m,1H), 5.10 (s,2H), 4.35 (m,2H), 3.85 (s,3H), 3.58 (s,3H). MS calcd. 482.18; MS (M+1)483.8
463	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-(2-chloro-phenoxy)	<sup>1</sup> H NMR: 8.47 (dd,1H), 8.33 (d,1H,J=2Hz), 7.65 (d,1H,J=8Hz), 7.42 (dd,1H), 7.19-7.25 (m,4H), 7.11 (dd,1H), 6.97 (dd,1H), 6.86-6.89 (m,1H), 6.82 (d,2H,J=2Hz), 6.78-6.80 (m,2H), 4.84 (s,3H), 3.11 (dd,3H), 1.39 (t,2H). MS calcd. 402.9; MS (M+1)403.8.
464	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-(2-chloro-phenoxy)	<sup>1</sup> H NMR: 7.34-7.50 (m,5H), 7.19 (m,3H), 6.91 (m,4H), 4.10 (dd,1H), 3.85 (br, NH), 2.02 (s,1H), 1.24 (t,1H). MS calcd. 456.87; MS (M+1)457.
465	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-(2-chloro-6-methyl-phenoxy)	<sup>1</sup> H NMR: 6.65-7.90 (m,11H), 3.70 (br, NH), 3.43 (m,1H), 2.13 (s,3H), 1.95 (m,2H), 1.20 (m,1H). MS calcd. 470.9; MS (M+1)471.6
466	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-(2-isopropoxy-phenoxy)	<sup>1</sup> H NMR: 8.73 (m,2H), 8.50 (s,1H), 7.98 (s,1H), 7.27 (s,1H), 6.94-7.12 (m,6H), 6.77 (s,2H), 5.10 (br, NH), 4.29-4.36 (m,4H), 1.02 (s,3H). MS calcd. 480.51; MS (M+1)481.4

-107-

Preparation 8N-(4-Methoxycarbonylphenyl)pyrid-3-ylmethylamine

A mixture of 3-pyridinecarboxaldehyde (4.72 mL, 50 mmol) and methyl 4-aminobenzoate (9.83 g, 65 mmol) in 140 mL of methanol was heated to reflux for 2.5 h then cooled to 0 °C. Solid NaBH<sub>4</sub> was added slowly and the reaction mixture was warmed to ambient temperature and stirred for 14 h. The mixture was concentrated and partitioned between DCM and H<sub>2</sub>O. The aqueous layer was extracted with DCM and the combined organic layers were washed with H<sub>2</sub>O and brine and concentrated. The residue was chromatographed on silica gel with DCM to 2% MeOH in DCM to give 8.08 g of a white solid (67% yield). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 8.57 (d, J=1.8Hz, 1H), 8.46 (dd, J=4.7, 1.8Hz, 1H), 7.73 (dt, J=8.4, 2.2Hz, 1H), 7.68 (d, J=8.8Hz, 2H), 7.35 (ddd, J=8.1, 4.8, 0.7Hz, 1H), 7.13 (br t, J=5.9Hz, 1H), 6.64 (d, J=8.8Hz, 2H), 4.38 (d, J=6.2Hz, 2H), 2.75 (s, 3H). MS calcd. 242.1; MS (M+1) 243.2.

Alternately, 3-pyridinecarboxaldehyde ((61.6 g, 0.575 moles, 1 eq) and methyl 4-aminobenzoate (103.87 g, 0.687 moles, 1.2 eq) with p-toluenesulfonic acid (2.19 g, .012 moles, 0.02 eq) were refluxed in methanol (25 volumes) for 2 hours. The reaction mixture was cooled to 0°C and sodium borohydride (67.6 g, 1.787 moles, 3.5 eq) added slowly then it was allowed to warm to ambient overnight. The reaction was concentrated, dissolved with dichloromethane and washed with water then brine, and evaporated to give the title compound.

Preparation 9N-(3-Methoxycarbonylphenyl)pyrid-3-ylmethylamine

The title compound was prepared using a method similar to Preparation 8 using 3-aminobenzoate. <sup>1</sup>H NMR: 8.64 (d, J=2.4Hz, 1H), 8.54 (dd, J=4.9, 1.5Hz, 1H), 7.69 (dt, J=7.8, 2.0Hz, 1H), 7.41 (dt, J=7.8, 1.3Hz, 2H), 7.32 (dd, J=2.5, 1.5, Hz, 1H), 7.27 (dd, J=7.8, 3.9Hz, 1H), 7.23



-108-

(t, J=8.3Hz, 1H), 6.79 (ddd, J=7.8, 2.5, 1.0Hz, 1H), 4.40 (d, J=5.9Hz, 2H), 4.19 (br t, J=5.9Hz, 1H) 3.88 (s, 3H).


#### Preparation 10

##### 5 N-(4-Benzylphenyl)pyrid-3-ylmethylaniline

The title compound was prepared using a method similar to Preparation 8 using 4-aminodiphenylmethane (purchased from TCI). <sup>1</sup>H NMR: 8.63 (br s, 1H), 8.53 (br d, J=4.4Hz, 1H), 7.50 (br d, J=7.8Hz, 1H), 7.26 (m, 3H), 7.18 (m, 3H), 7.00 (m, 2H), 6.57 (m, 2H), 4.34 (s, 2H), 3.99 (br s, 1H), 3.88 (s, 2H).


#### Example 467

##### 15 N-(4-(Benzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline




To a solution of N-(4-benzylphenyl)pyrid-3-ylmethylaniline (0.517 g, 1.88 mmol) dissolved in 8 mL dichloroethane and 4 mL pyridine at 0 °C was added 2,2,2-trifluoroethylsulfonyl chloride (0.313 mL, 2.83 mmol). The reaction was stirred at ambient temperature overnight. The mixture was treated with 1.5 g anhydrous K<sub>2</sub>CO<sub>3</sub> and stirred for 2 h. Then the mixture was filtered and concentrated. The residue was chromatographed on silica gel with 3:1 to 25 2:1 DCM/EtOAc to give an amorphous white solid. <sup>1</sup>H NMR: 8.52 (br d, J=3.3Hz, 1H), 8.32 (br s, 1H), 7.71 (dt, J=6.8, 1.4Hz, 1H), 7.33-7.20 (m, 4H), 7.18-7.10 (m, 6H), 4.88 (s, 2H), 3.95 (s, 2H), 3.79 (q, J=9.3Hz, 2H).

-109-

Example 468N-(4-(Benzyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline

The title compound was prepared using a method similar  
 5 to the method of Example 467 using ethanesulfonyl chloride.  
<sup>1</sup>H NMR: 8.50 (br d, J=3.4Hz, 1H), 8.36 (br s, 1H), 7.77  
 (dd, J=7.8, 1.5Hz, 1H), 7.31-7.19 (m, 4H), 7.17-7.09 (m, 6H),  
 4.87 (s, 2H), 3.92 (s, 2H), 3.08 (q, J=7.3Hz, 2H), 1.42  
 (t, J=7.3Hz, 3H). MS calcd. 366.1; MS (M+1) 367.1.

10

Example 469N-(4-Methoxycarbonylphenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline


15 The title compound was prepared using a method similar  
 to the method of Example 467 using N-(4-methoxycarbonyl-  
 phenyl)pyrid-3-ylmethylaniline. <sup>1</sup>H NMR: 8.52 (br s, 1H), 8.37  
 (br s, 1H), 8.02 (m, 2H), 7.68 (d, J=7.8Hz, 1H), 7.33 (m, 2H),  
 7.27 (m, 1H), 4.95 (s, 2H), 3.90 (s, 3H), 3.82 (q, J=9.3Hz, 2H).

20

-110-

Example 470

N-(3-Methoxycarbonylphenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylamine



5


The title compound was prepared using a method similar to the method of Example 467 using N-(3-methoxycarbonylphenyl)pyrid-3-ylmethylamine. <sup>1</sup>H NMR: 8.51

(dd, J=4.8, 1.8 Hz, 1H), 8.33 (d, J=1.8 Hz, 1H), 8.00  
 10 (dt, J=7.3, 1.5 Hz, 1H), 7.93 (t, J=1.5 Hz, 1H), 7.68  
 (dt, J=7.7, 2.2 Hz, 1H), 7.43 (t, J=7.7 Hz, 1H), 7.39  
 (dt, J=7.7, 1.5 Hz, 1H), 7.25 (ddd, J=7.3, 4.8, 1.5 Hz, 1H), 4.93  
 (s, 2H), 3.92 (s, 2H), 3.82 (q, J=9.3 Hz, 2H).

15

Example 471

N-(4-(Hydoxymethyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylamine




To a solution of N-(4-hydroxymethyl-phenyl)pyrid-3-ylmethylamine (0.472 g, 1.22 mmol) dissolved in 15 mL THF at,  
 20 -78 °C was DIBAL (1.0 M in toluene, 3.04 mL, 3.04 mmol).  
 The reaction was stirred for 1h and then warmed to 0 °C for  
 1 h. It was quenched with 0.5 mL methanol and stirred  
 overnight with 60 mL ether and 30 mL saturated Rochelle's

-111-

salt. The mixture was extracted with ether, dried over  $\text{MgSO}_4$ , filtered and concentrated. The residue was chromatographed on silica gel with 1:1 DCM/EtOAc to give the title compound as an amorphous white solid.  $^1\text{H}$  NMR: 8.43 (d,  $J=3.8\text{Hz}$ , 1H), 8.27 (s, 1H), 7.68 (d,  $J=7.8\text{Hz}$ , 1H), 7.32 (m, 2H), 7.23 (m, 1H), 7.19 (m, 2H), 4.88 (s, 2H), 4.63 (s, 2H), 3.80 (q,  $J=9.3\text{Hz}$ , 2H), 3.10 (br s, 1H).

Example 472

10 N-(3-(Hydroxymethyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline




The title compound was prepared using a method similar to the method of Example 471 using N-(3-methoxycarbonylphenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline.  $^1\text{H}$  NMR: 8.50 (dd,  $J=4.9, 1.5\text{Hz}$ , 1H), 8.31 ( $J=2.0\text{Hz}$ , 1H), 7.70 (dt,  $J=7.8, 2.0\text{Hz}$ , 1H), 7.34 (m, 2H), 7.25 (m, 2H), 7.14 (dt,  $J=7.3, 2.5\text{Hz}$ , 1H), 4.90 (s, 2H), 4.67 (d,  $J=5.4\text{Hz}$ , 2H), 3.81 (q,  $J=9.3\text{Hz}$ , 1H), 1.92 (br t,  $J=5.4\text{Hz}$ , 1H).

20

Example 473

N-(4-Formylphenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline




-112-

To a solution of DMSO (0.190 mL, 2.68 mmol) in 4 mL DCM at -78 °C was added oxalyl chloride (0.146 mL, 1.67 mmol) dropwise. After 15 min a solution of N-(4-(hydroxymethyl)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethanamine (0.241 g, 0.669 mmol) was added via cannula. The reaction was stirred for 30 min and triethylamine (0.932 mL, 6.69 mmol) was added and warmed to 0 °C over 1 h. The mixture was poured into saturated NaHCO<sub>3</sub> solution and diluted with ether. The organic layer was washed with H<sub>2</sub>O and brine, dried over MgSO<sub>4</sub>, filtered, and concentrated. The residue was chromatographed on silica gel with 5:1 DCM/EtOAc to give 173 mg of a clear oil (72%). <sup>1</sup>H NMR: 9.97 (s, 1H), 8.50 (dd, J=4.9, 1.5 Hz, 1H), 8.35 (d, J=2.0 Hz, 1H), 7.87 (m, 2H), 7.66 (dt, J=7.8, 2.0 Hz, 1H), 7.44 (m, 2H), 7.24 (ddd, J=7.8, 4.9, 1.0 Hz, 1H), 4.97 (s, 2H), 3.84 (q, J=9.3 Hz, 2H).

Example 474

N-(3-Formylphenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethanamine



-113-

The title compound was prepared using a method similar to the method of Example 473 using N-(3-hydroxymethyl-phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-

ylmethylamine.  $^1\text{H}$  NMR: 9.97 (s, 1H), 8.51

5 (dd, J=4.9, 1.5Hz, 1H), 8.33 (d, J=2.0Hz, 1H), 7.85

(dt, J=7.8, 2.0Hz, 1H), 7.77 (t, J=2.0Hz, 1H), 7.69

(dt, J=7.8, 2.0Hz, 1H), 7.54 (t, J=7.8Hz, 1H), 7.48


(dt, J=7.8, 2.0Hz, 1H), 7.26 (dd, J=7.8, 4.9Hz, 1H), 4.95 (s, 2H),

3.83 (q, J=9.3Hz, 2H).

10

#### Example 475

N-(3-( $\alpha$ -Hydroxybenzyl)phenyl)-N-(2,2,2-  
trifluoroethanesulfonyl)pyrid-3-ylmethylamine



15 To a solution of N-(3-formylphenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylamine (0.174 g, 0.486 mmol) in 4 mL THF at 0 °C was added phenyl magnesium chloride (2.0 M in THF, 0.364 mL, 0.728 mmol). After 1 h at ambient temperature, the reaction was poured in to saturated

20  $\text{NaHCO}_3$  solution and extracted with DCM. The extracts were dried over  $\text{MgSO}_4$ , filtered and concentrated. The residue was chromatographed on silica gel with 1:1 DCM/EtOAc to give a clear oil.  $^1\text{H}$  NMR: 8.41 (dd, J=4.9, 1.5Hz, 1H), 8.24


(d, J=2.0Hz, 1H), 7.67 (dt, J=7.8, 2.0Hz, 1H), 7.40=7.15 (m, 10H),

25 5.78 (s, 1H), 4.86 (s, 2H), 3.77 (q, J=9.3Hz, 2H), 3.21 (br s, 1H). MS calcd. 436.1; MS (M+1) 437.1. The HCl salt was prepared by adding 1.0 M HCl in ether solution to the free base in ether suspension, stirring for 10 min then collecting the ppt by filtration.

-114-

Optionally, the HCl salt can be prepared by adding 1.0 M HCl in ether solution to the free base in ether suspension, stirring for 10 min then collecting the ppt by filtration.

5 By a method similar to the method of Example 475 the following compounds were prepared:




No.	R2	Data
476	4-( $\alpha$ -hydroxybenzyl)	<sup>1</sup> H NMR: 8.38 (dd, J=4.9, 2.0 Hz, 1H), 8.21 (d, J=1.5 Hz, 1H), 7.61 (dt, J=7.8, 1.5 Hz, 1H), 7.33–7.09 (m, 10H), 5.72 (d, J=2.5 Hz, 1H), 4.79 (s, 2H), 3.71 (q, J=9.3 Hz, 2H), 2.71 (d, J=3.4 Hz, 1H)
477	4-( $\alpha$ -hydroxy-2-methylsulfanyl benzyl)	<sup>1</sup> H NMR: 8.39 (dd, J=4.9, 2.0 Hz, 1H), 8.24 (d, J=1.5 Hz, 1H), 7.65 (dt, J=7.8, 1.5 Hz, 1H), 7.43–7.38 (m, 3H), 7.27–7.13 (m, 6H), 6.20 (d, J=1.5 Hz, 1H), 4.85 (s, 2H), 3.78 (q, J=9.3 Hz, 2H), 3.60 (br s, 1H), 2.38 (s, 3H). MS calcd. 482.1; MS (M+1) 483.1
478	3-( $\alpha$ -hydroxy-2-methylsulfanyl benzyl)	<sup>1</sup> H NMR: 8.40 (dd, J=4.9, 2.0 Hz, 1H), 8.22 (d, J=2.0 Hz, 1H), 7.62 (dt, J=7.8, 1.5 Hz, 1H), 7.39–7.33 (m, 2H), 7.32–7.15 (m, 6H), 7.06 (d, J=7.8 Hz, 1H), 6.14 (d, J=3.4 Hz, 1H), 4.85 (A of AB, d, J=14.8 Hz, 1H), 4.80 (B of AB, d, J=14.8 Hz, 1H), 3.84–3.70 (m, 2H), 3.53 (d, J=4.3 Hz, 1H), 2.39 (s, 3H). MS calcd. 482.1; MS (M+1) 483.1
479	4-( $\alpha$ -hydroxy-2-methoxybenzyl)	<sup>1</sup> H NMR: 8.45 (dd, J=4.9, 1.5 Hz, 1H), 8.29 (d, J=2.0 Hz, 1H), 7.67 (dd, J=8.3 Hz, 1H), 7.38 (d, J=8.3 Hz, 2H), 7.29–7.20 (m, 3H), 7.16 (d, J=8.6 Hz, 2H), 6.94 (dt, J=1.0, 7.8 Hz, 1H), 6.88 (d, J=8.3 Hz, 1H), 6.01 (d, J=4.9 Hz, 1H), 4.87 (s, 2H), 3.78 (q, J=9.3 Hz, 2H), 3.77 (s, 3H)
480	3-( $\alpha$ -hydroxy-2-methoxybenzyl)	<sup>1</sup> H NMR: 8.42 (dd, J=4.9, 1.5 Hz, 1H), 8.24 (d, J=2.0 Hz, 1H), 7.63 (dt, J=7.8, 2.0 Hz, 1H), 7.37 (d, J=7.3 Hz, 1H), 7.34–7.24 (m, 3H), 7.22–7.17 (m, 2H), 7.09 (br d, J=9.0 Hz, 1H), 6.96 (dt, J=1.0, 7.3 Hz, 1H), 6.88 (d, J=8.3 Hz, 1H), 6.03 (s, 1H), 4.87 (s, 2H), 3.80 (br s, 1H), 3.79 (q, J=9.3 Hz, 2H), 3.77 (s, 3H).
481	4-( $\alpha$ -hydroxy-3,4-	<sup>1</sup> H NMR: 8.41 (dd, J=4.9, 2.0 Hz, 1H), 8.23

- 115 -

	difluorobenzyl)	(d,J=1.5Hz,1H), 7.69 (dt,J=7.8,2.0Hz,1H), 7.34 (d,J=7.3Hz,2H), 7.24 (dd,J=4.9,4.0Hz,1H), 7.21-7.07 (m,4H), 7.03 (m,1H), 5.73 (s,1H), 4.87 (s,2H), 3.79 (q,J=9.3Hz,2H), 3.34 (br s,1H). MS calcd. 472.1; MS (M+1) 473.1
482	3-( $\alpha$ -hydroxy-3,4-difluorobenzyl)	<sup>1</sup> H NMR: 8.29 (d,J=3.9Hz,1H), 8.12 (d,J=1.5Hz,1H), 7.57 (dt,J=7.8,2.0Hz,1H), 7.34-7.24 (m,2H), 7.18-7.11 (m,3H), 7.09-6.96 (m,2H), 6.89 (m,1H), 5.63 (s,1H), 4.82 (s,2H), 4.67 (br s,1H), 3.77 (q,J=9.3Hz,2H). MS calcd. 472.1; MS (M+1) 473.1
483	4-( $\alpha$ -hydroxy-2-phenylbenzyl)	<sup>1</sup> H NMR: 8.46 (d,1H,J=3.4Hz), 8.26 (s,1H), 7.76-7.60 (m,1H), 7.56-7.45 (m,1H), 7.42-7.30 (m,5H), 7.44-7.15 (m,4H), 7.15-7.00 (m,5H), 5.89 (d,1H,J=3.4Hz), 4.85 (s,2H), 3.76 (q,2H,J=9.8Hz)
484	3-( $\alpha$ -hydroxy-2-phenylbenzyl)	<sup>1</sup> H NMR: 8.45 (m,1H), 8.22 (d,1H), 7.62-7.57 (m,1H), 7.43-7.30 (m,6H), 7.28-7.14 (m,5H), 7.10-6.98 (m,3H), 5.90 (s,1H), 4.80 (q,2H), 3.82-3.62 (q,2H)
485	4-(1-hydroxy-1-(5-methylpyrid-2-yl)methyl)	<sup>1</sup> H NMR: 8.50 (d,1H), 8.37 (s,1H), 8.31 (s,1H), 7.63 (d,1H), 7.15 (m,6H), 7.00 (d,1H), 5.68 (d,1H), 4.84 (s,2H), 3.78 (q,2H), 2.32 (s,3H).
486	3-(1-hydroxy-1-(5-methylpyrid-2-yl)methyl)	<sup>1</sup> H NMR: 8.49 (m,1H), 8.35 (d,1H,J=1.5Hz), 7.70-7.65 (d,2H,7.8Hz), 7.57-7.52 (m,1H), 7.46-7.16 (m,7H), 6.26 (d,1H), 4.87(s,1H), 3.88 (q,2H)
487	4-( $\alpha$ -hydroxy-2-trifluoromethylbenzyl)	<sup>1</sup> H NMR: 8.47 (q,1H,J=2.0, 2.93), 8.24 (s,1H,J=2.0Hz), 7.70-7.52 (m,3H), 7.46-7.08 (m,8H), 6.24 (d,1H), 4.83 (s,2H), 3.75 (q,2H)
488	3-( $\alpha$ -hydroxy-2-fluorobenzyl)	<sup>1</sup> H NMR: 8.43 (d,1H), 8.24 (s,1H), 7.68 (m,1H), 7.42-6.88 (m,9H), 5.78 (s,1H), 4.87 (s,2H), 3.86 (q,2H)
489	4-( $\alpha$ -hydroxy-2-fluorobenzyl)	<sup>1</sup> H NMR: 8.46 (d,1H,J=3.4), 8.30 (s,1H), 7.65 (m,1H), 7.47-7.38 (m,3H), 7.30-6.82 (m,6H), 6.12 (d,1H), 4.87 (s,2H), 3.75 (q,2H,J=8.8Hz)
490	4-( $\alpha$ -hydroxy-3-fluorobenzyl)	<sup>1</sup> H NMR: 8.43 (d,1H), 8.24 (s,1H), 7.68 (m,1H), 7.42-6.88 (m,9H), 5.78 (s,1H), 4.87 (s,2H), 3.86 (q,2H).

## Example 491

N-(3-(Benzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylamine






-116-

To a solution of N-3-( $\alpha$ -hydroxybenzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylamine (0.050 g, 0.115 mmol) in 2 mL DCM and 0.5 mL triethylsilane was added trifluoroacetic acid (0.3 mL). The reaction was stirred at ambient temperature overnight and poured into saturated NaHCO<sub>3</sub> solution and extracted with EtOAc. The combined extracts were dried over MgSO<sub>4</sub>, filtered and concentrated. The residue was chromatographed on silica gel with 4:1 to 2:1 DCM/EtOAc to give 0.025 g of a clear oil (52% yield). <sup>1</sup>H NMR: 8.51 (dd, J=4.9, 1.5 Hz, 1H), 8.22 (d, J=2.0 Hz, 1H), 7.69 (dt, J=7.8, 2.0 Hz, 1H), 7.32–7.20 (m, 4H), 7.18–7.11 (m, 6H), 4.86 (s, 2H), 3.94 (s, 2H), 3.79 (q, J=9.3 Hz, 2H). MS calcd. 420.1; MS (M+1) 421.1. Optionally, the HCl salt can be prepared by adding 1.0 M HCl in ether solution to the free base in ether suspension, stirring for 10 min then collecting the ppt by filtration.

By a method similar to the method of Example 491 the following compounds were prepared:




No.	R <sub>2</sub>	Data
492	4-(2-methylsulfanyl benzyl)	<sup>1</sup> H NMR: 8.50 (vbr s, 2H), 7.71 (br d, J=7.8 Hz, 1H), 7.29 (br s, 1H), 7.23 (m, 2H), 7.17 (br d, J=8.3 Hz, 2H), 7.11 (m, 2H), 7.05 (br d, 8.3 Hz, 2H), 3.87 (s, 2H), 4.04 (s, 2H), 3.79 (q, J=9.3 Hz, 2H), 2.41 (s, 3H). MS calcd. 466.1; MS (M+1) 467.0.
493	3-(2-methylsulfanyl benzyl)	<sup>1</sup> H NMR: 8.50 (br s, 1H), 8.31 (br s, 1H), 7.63 (d, J=7.8 Hz, 1H), 7.28–7.15 (m, 5H), 7.11 (dt, J=1.5, 6.8 Hz, 1H), 7.02 (m, 3H), 4.73 (s, 2H), 4.03 (s, 2H), 3.78 (q, J=9.3 Hz, 2H), 2.42 (s, 3H). MS calcd. 466.1; MS (M+1) 467.0.
494	4-(2-methoxybenzyl)	<sup>1</sup> H NMR: 8.50 (br d, J=3.9 Hz, 1H), 8.33 (br d, J=7.8 Hz, 1H), 7.68 (br d, J=7.8 Hz, 1H), 7.28–7.16

-117-

		(m,4H), 7.12–7.20 (m,3H), 6.92–6.85 (m,2H), 4.86 (s,2H), 3.93 (s,2H), 3.79 (q,J=9.3Hz,2H), 3.78 (s,3H)
495	3-(2-methoxybenzyl)	<sup>1</sup> H NMR: 8.48 (d,J=3.4Hz,1H), 8.30 (d,J=1.5Hz,1H), 7.61 (dt,J=7.8,2.0Hz,1H), 7.24–7.15 (m,4H), 7.06 (br s,1H), 7.03–6.95 (m,2H), 6.91–6.84 (m,2H), 4.84 (s,2H), 3.91 (s,2H), 3.77 (s,3H), 3.75 (q,J=9.3Hz,2H)
496	4-(3,4-difluorobenzyl)	<sup>1</sup> H NMR: 8.51 (dd,J=4.9,1.5Hz,1H), 8.31 (d,J=2.0Hz,1H), 7.69 (dt,J=7.8,2.0Hz,1H), 7.25 (dd,J=7.8,4.9Hz,1H), 7.17–7.04 (m,5H), 6.93 (m,1H), 6.86 (m,1H), 4.87 (s,2H), 3.89 (s,2H), 3.80 (q,J=9.3Hz,2H)
497	3-(3,4-difluorobenzyl)	<sup>1</sup> H NMR: 8.50 (dd,J=4.9,1.5Hz,1H), 8.29 (d,J=1.5Hz,1H), 7.62 (dt,J=7.8,2.0Hz,1H), 7.31 (t,J=7.8Hz,1H), 7.21 (dd,J=7.8,4.9Hz,1H), 7.15–7.02 (m,3H), 6.91 (s,1H), 6.82–6.71 (m,2H), 4.85 (s,2H), 3.85 (s,2H), 3.79 (q,J=9.3Hz,2H). MS calcd. 456.1; MS (M+1) 457.1
498	4-(2-phenylbenzyl)	<sup>1</sup> H NMR: 8.52 (br s,1H), 8.31 (br s,1H), 7.66 (d,1H,J=7.8Hz), 7.30 (m,4H), 7.27–7.17 (m,4H), 7.13 (m,2H), 7.01 (d,2H,J=8.4Hz), 6.88 (d,2H,J=8.4Hz), 4.84 (s,2H), 3.92 (s,2H), 3.76 (q,2H,J=9.3Hz). MS calcd 496.1; MS (M+1) 497.1
499	3-(2-phenylbenzyl)	<sup>1</sup> H NMR: 8.42 (br s,1H), 7.19 (br s,1H), 7.51 (d,1H,J=7.8Hz), 7.28–7.09 (m,7H), 7.06 (m,3H), 6.99 (m,1H), 6.88 (d,1H,J=8.3Hz), 6.82 (d,1H,J=7.8Hz), 6.67 (s,1H), 4.70 (s,2H), 3.83 (s,2H), 3.62 (q,2H,J=9.3Hz). MS calcd 496.1; MS (M+1) 497.1

Example 500

N-(3-Methoxycarbonylphenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine



5

Into a 0°C solution of N-(3-methoxycarbonylphenyl)pyrid-3-ylmethylanine (204 g, 0.842 mole, 1 eq) in dichloroethane (3610 mL) and pyridine (1775 mL) was added ethanesulfonyl chloride (114 mL, 1.2 mole, 1.43 eq). The reaction was allowed to warm to room temperature overnight.


10

-118-

The dark red reaction was stirred with potassium carbonate (204 g) for 30 minutes and then filtered and concentrated to dryness. The residue was purified by Biotage, eluting with 3:1 to 2:1 DCM/ EtOAc mixture to give the title compound as a white solid (137.0 g, 49% yield).

Example 501

N-(3-(Hydroxymethyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine




10

Into a  $-78^{\circ}\text{C}$  solution of N-(3-methoxycarbonylphenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine (75 g, 0.224 mole, 1eq) in THF was added slowly DIBAL-H (561 mL, 1M in toluene, 2.5 eq). The mixture was stirred for 1 hour and then was warmed to  $0^{\circ}\text{C}$  with an ice water bath and stirred for 2 h. The reaction was carefully quenched with methanol (18.50 mL, 0.396 mole, 1.8 eq). It was then diluted with ether (4650 mL) and Saturated Rochelle's salt (2775 mL) and stirred vigorously overnight. The layers were separated and the aqueous extracted with EtOAc (2000 mL). The combined organics were washed with brine, dried with  $\text{MgSO}_4$ , filtered and concentrated. The residue was chromatographed, eluting with 1:1 DCM/EtOAc to give the title compound as a white solid (51.4 g, 43% Yield).

20

-119-


Example 502N-(3-Formylphenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline

Into a  $-78^{\circ}\text{C}$  solution of DMSO (20.8 mL, 0.269 mole, 3.1  
5 eq) in DCM (460 mL) was added oxalyl chloride (16 mL, 0.109  
mole, 1.25 eq) dropwise and stirred for 15 minutes. N-(3-  
(Hydroxymethyl)phenyl)-N-(ethanesulfonyl)pyrid-3-  
ylmethylaniline (26.4 g, 0.86 mole, 1 eq) was added dropwise  
via addition funnel as a solution in DCM (146 mL). The  
10 reaction mixture was stirred for 30 min. then TEA (102 mL, 1  
mole, 11.7 eq) was added via addition funnel. The reaction  
mixture was allowed to warm to  $0^{\circ}\text{C}$  with an ice bath and  
stirred for 1 hour with a  $\text{N}_2$  sweep through a bleach/caustic  
scrubber. The reaction mixture was poured into saturated  
15  $\text{NaHCO}_3$  and diluted with diethyl ether (2000 mL). The layers  
were separated and the aqueous layer backextracted with  
diethyl ether (1000 mL). The combined organics were washed  
with brine (1000 mL), dried with  $\text{MgSO}_4$ , filtered and  
concentrated. The residue was purified by flash  
20 chromatography, eluting with 1:1 DCM/EtOAc to give the title  
compound as a white solid (15.0 g, 57.2%).

-120-

Example 503

N-(4-(Hydoxymethyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine



5 To a  $-78^{\circ}\text{C}$  solution of N-(4-methoxycarbonylphenyl)pyrid-3-ylmethylamine (105 g, 0.433 moles, 1eq) in THF (17 volumes) was added slowly DIBAL-H (1M in toluene, 1084 mL, 1.084 moles, 2.5 eq,) and held cold for 1 hour. The reaction was then allowed to warm to  $0^{\circ}\text{C}$  for 2 h. The  
10 reaction was carefully quenched with methanol (1.8 eq). It was then diluted with ether (62 volumes) and Saturated Rochelle's salt (37 volumes) and stirred vigorously overnight. The layers were separated and the aqueous extracted with EtOAc. The combined organics were washed  
15 with brine, dried with  $\text{MgSO}_4$ , filtered, and concentrated to give N-(4-hydroxymethylphenyl)pyrid-3-ylmethylamine (88.2 g, 95 % Yield).

To a  $0^{\circ}\text{C}$  solution of give N-(4-hydroxymethylphenyl)pyrid-3-ylmethylamine (75 g, 0.35 moles, 1eq) and imidazole  
20 (26.21 g, 0.384 moles, 1.1 eq) in DCM was added slowly t-butyldimethylsilyl chloride (85.09 g, 564.5 moles, 1.6 eq) and held cold for 1/4 hour. The reaction was then allowed to warm to  $0^{\circ}\text{C}$  overnight. The reaction was partitioned with water and DCM. The organic layer was stripped to give N-(4-t-butyldimeythsilyloxymethylphenyl) pyrid-3-ylmethylamine a  
25 yellow solid (115 g, 100% yield)


To a  $0^{\circ}\text{C}$  solution of N-(4-t-butyldimeythsilyloxy methylphenyl)pyrid-3-ylmethylamine (115.25 g, 0.351 moles, 1 eq) in dichloromethane (DCM, 17.7volumes) and pyridine

-121-

(8.7volumes) was added ethanesulfonyl chloride (50.26 g, 0.391 moles, 1.1 eq). The reaction was allowed to warm to room temperature overnight. The dark red reaction was stirred with  $K_2CO_3$  (1 wt eq) for 30 minutes and then  
5 filtered and concentrated to dryness. The residue was purified by Biotage with 3:1 to 2:1 DCM/ EtOAc to give N-(4-t-butyl dimeythsilyloxymethylphenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylamine a yellow solid (91.24 g, 62% yield).

To a 0°C solution of N-(4-t-butyl dimeythsilyloxymethylphenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylamine (90.61 g, 0.215 moles, 1 eq) in DCM (10  
10 volumes) was added tetrabutylammonium fluoride (1.0 M in THF, 252.4 mL, 1.17 eq) dropwise and stirred for 15 minutes. The bath was removed and the reaction allowed to stir  
15 overnight. Washed with sat.  $Na_2CO_3$  solution, water, and then brine. Dried the organics with  $Na_2SO_4$ . Stripped to yellow oil. Combined lots and prepped clean with 50/50 EtOAc/DCM on Biotage 75. Triturated solid product to give the title compound (30.0 g, 46% yield).

20

Example 504N-(4-Formylphenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine

To a solution of DMSO (0.190 mL, 2.68 mmol) in 4 mL DCM  
25 at -78 °C was added oxalyl chloride (0.146 mL, 1.67 mmol) dropwise. After 15 min a solution of N-(4-(hydroxymethyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine (0.241 g, 0.669 mmol) was added via cannula. The reaction was stirred for 30 min and triethylamine (0.932

-122-

mL, 6.69 mmol) was added and warmed to 0 °C over 1 h. The mixture was poured into saturated NaHCO<sub>3</sub> solution and diluted with ether. The organic layer was washed with H<sub>2</sub>O and brine, dried over MgSO<sub>4</sub>, filtered, and concentrated.


- 5 The residue was chromatographed on silica gel with 5:1 DCM/EtOAc to give 173 mg of a clear oil (72%). <sup>1</sup>H NMR: 9.97 (s, 1H), 8.50 (dd, J=4.9, 1.5 Hz, 1H), 8.35 (d, J=2.0 Hz, 1H), 7.87 (m, 2H), 7.66 (dt, J=7.8, 2.0 Hz, 1H), 7.44 (m, 2H), 7.24 (ddd, J=7.8, 4.9, 1.0 Hz, 1H), 4.97 (s, 2H), 3.84 (q, J=9.3 Hz, 2H).

10

Example 505

N-(3-(Phenoxymethyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline

-123-




To a solution of triphenylphosphine (0.141 g, 0.539 mmol) in 3.5 mL THF was added diethylazodicarboxylate (0.165 g, 0.539 mmol). After 5 min phenol (0.061 g, 0.646 mmol) was added as a solution in 2 mL THF. After 5 min N-(3-(hydroxymethyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethanamine (0.165 g, 0.539 mmol) was added as a solution in 2 mL THF and the reaction was stirred for 3 days. The mixture was diluted with EtOAc, washed with 1 N NaOH and brine, dried over MgSO<sub>4</sub>, filtered and concentrated. The residue was chromatographed on silica gel with 4:1 to 2:1 DCM/EtOAc to give the title compounds as a clear oil. <sup>1</sup>H NMR: 8.48 (d, J=1.5Hz, 1H), 8.36 (s, 1H), 7.64 (d, J=7.8Hz, 1H), 7.36-7.23 (m, 5H), 7.19 (m, 2H), 6.95 (t, J=7.8Hz, 1H), 6.91 (d, J=7.8Hz, 1H), 5.01 (s, 2H), 4.84 (s, 2H), 3.03 (q, J=7.6Hz, 2H), 1.39 (t, J=7.6Hz, 3H). MS calcd. 382.1; MS (M+1) 383.1.



-124-


Example 506

N-(4-(Phenoxymethyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine



5 The title compound was prepared using a method similar to the method of Example 505 using N-(4-hydroxymethylphenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylamine. <sup>1</sup>H NMR: 8.41 (d, 1H, J=3.4Hz), 8.29 (d, 1H, J=1.5Hz), 7.63 (d, 1H, J=7.8Hz), 7.00-7.40 (m, 7H), 6.86 (m, 3H), 4.92 (s, 2H),  
 10 4.81 (s, 2H), 3.00 (q, 2H, J=7.3Hz), 1.34 (t, 3H, J=7.3Hz). MS calcd 382.1; MS (M+1) 383.1.

By a method similar to the method of Example 475 the following compounds were prepared:



15


No.	R2	Data
507	3-(α-hydroxybenzyl)	<sup>1</sup> H NMR: 8.43 (dd, J=4.9, 1.5Hz, 1H), 8.29 (d, J=2.0Hz, 1H), 7.63 (dt, J=8.3, 1.5Hz, 1H), 7.36-7.21 (m, 8H), 7.19-7.12 (m, 2H), 5.77 (s, 1H), 4.84 (s, 2H), 3.06 (q, J=7.3Hz, 2H), 2.68 (br s, 1H), 1.39 (t, J=7.3Hz, 3H)
508	4-(α-hydroxybenzyl)	<sup>1</sup> H NMR: 8.48 (d, 1H, J=3.9Hz), 8.35 (br s, 1H), 7.73 (d, 1H, J=7.8Hz), 7.34 (m, 5H), 7.30-7.23 (m, 3H), 7.21 (d, 2H, J=8.3Hz), 5.80 (s, 2H), 4.88 (s, 2H), 3.07 (q, 2H, J=7.3Hz), 1.41 (t, 3H, J=7.3Hz). MS calcd

-125-

		382.1; MS (M+1) 383.1
509	3-( $\alpha$ -hydroxy-2-fluorobenzyl)	$^1\text{H}$ NMR: 8.46 (d,1H,J=3.4Hz), 8.32 (s,1H), 7.62 (m,1H), 7.42-6.92 (m,9H), 6.00 (s,1H), 4.85 (s,2H), 3.06 (m,2H), 1.38 (m,3H). MS calcd 400.1; MS (M+1) 401.1
510	3-( $\alpha$ -hydroxy-3-fluorobenzyl)	$^1\text{H}$ NMR: 8.46 (d,1H), 8.34 (s,1H), 7.62 (d,1H), 7.40-7.15 (m,6H), 7.04-6.88 (m,3H), 5.75 (s,1H), 4.85 (s,2H), 3.07 (q,2H,J=7.3Hz), 1.40 (t,3H,7.3Hz). MS calcd 400.1; MS (M+1) 401.1

Example 511

N-(3-( $\alpha$ -Hydroxy-4-ethoxycarbonylbenzyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine



5

A solution of ethoxycarbonylphenylzinc iodide in THF (Rieke Metals, Inc.) (0.5 M, 2.8 ml) was evaporated to dryness under nitrogen. To the residue was added chromium(III) chloride (0.210 g, 1.33 mmole), N-(3-formylphenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylamine (0.210 g, 0.7 mmole), and tetramethylurea (0.5 + 1.5 ml), followed by chlorotrimethylsilane (0.7 ml, 0.55 mmole) dropwise. The mixture was stirred overnight at room temperature. A few pieces of ice, 1 ml of saturated  $\text{NaHCO}_3$  and 1 ml of ethylene diamine were added to the reaction mixture, which was filtered. The organic layer was washed with brine, after extraction with ethyl acetate. Evaporation afforded the crude product, that was chromatographed on a silica-gel column eluted with ethyl acetate and chloroform to give the title compound.  $^1\text{H}$  NMR: 8.44 (d,1H, J=4.1Hz), 8.29 (s,1H), 7.99 (d,2H,J=8Hz), 7.62 (d,1H), 7.34-7.14 (m,7H), 5.80 (s,1H), 4.85 (s,2H), 4.36 (q,2H, J=7.1Hz), 3.05


20

-126-

(q, 2H, J=7.4Hz), 1.41 (m, 6H). MS calcd. 454.2; MS (M+1) 455.2.


Example 512

N- (3- (α-Hydroxy-3-ethoxycarbonylbenzyl) phenyl) -N-  
5 (ethanesulfonyl)pyrid-3-ylmethylaniline



The title compound was prepared using a method similar to the method of Example 511 using 3-ethoxycarbonylphenylzinc chloride (Rieke Metals Inc.) as  
10 starting materials. <sup>1</sup>H NMR: 8.43-8.29 (m, 2H), 7.63 (d, 1H), 7.40-7.00 (m, 9H), 5.80 (s, 1H), 4.85 (s, 2H), 4.36 (q, 2H), 3.08 (m, 4H), 1.38 (m, 6H). MS calcd 454.2; MS (M+1) 455.2.

By a method similar to the method of Example 491 the  
15 following compounds were prepared:




No.	R2	Data
513	3-(4-ethoxycarbonyl benzyl)	<sup>1</sup> H NMR: 8.48 (d, 1H, J=4.3Hz), 8.33 (s, 1H), 7.95 (d, 1H, J=8.2Hz), 7.62 (d, 1H, J=7.8Hz), 7.27-6.94 (m, 7H), 4.85 (s, 2H), 4.36 (q, 2H, J=7.3Hz), 3.95 (s, 2H), 3.10 (q, 2H, J=7.0Hz), 2.04 (1H), 1.38 (m, 6H). MS calcd 438.2; MS (M+1) 439.1
514	3-(2-fluorobenzyl)	<sup>1</sup> H NMR: 8.47 (m, 1H), 8.36 (d, 1H, J=2.0Hz), 7.64 (m, 1H), 7.26-6.96 (m, 9H), 4.85 (s, 2H), 3.93 (s, 2H), 3.08 (q, 2H). MS calcd 384.1; MS (M+1) 385.2

-127-

Example 515


N-(3-( $\alpha$ -Fluorobenzyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine



5 To a solution of N-(3-( $\alpha$ -hydroxybenzyl)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylamine (0.106 g, 2.77 mmol) in DCM was added DAST (0.040 mL, 3.05 mmol). The reaction was stirred overnight and poured into saturated NaHCO<sub>3</sub> solution and extracted with EtOAc, dried over MgSO<sub>4</sub>, filtered, and  
10 concentrated. The residue was chromatographed on silica gel with 2:1 DCM/EtOAc to give the title compound as a clear oil. <sup>1</sup>H NMR: 8.47 (dd, J=4.9, 1.5 Hz, 1H), 8.33 (d, J=2.0 Hz, 1H), 7.63 (dt, J=7.8, 2.0 Hz, 1H), 7.38-7.28 (m, 4H), 7.24-7.16 (m, 6H), 4.89 (A of AB, d, J=14.8 Hz, 1H), 4.83 (B of AB, d, J=14.8 Hz, 1H), 3.07 (q, J=7.3 Hz, 2H), 1.40 (t, J=7.3 Hz, 3H). MS calcd. 384.1; MS (M+1) 385.1.

Example 516

20 N-(4-( $\alpha$ -Fluorobenzyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine




-128-

By the method of Example 515 using N-(4-( $\alpha$ -hydroxybenzyl)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline gave the title compound.  $^1\text{H}$  NMR: 8.49 (d, 1H,  $J=3.9\text{Hz}$ ), 8.37 (br s, 1H), 7.72 (d, 1H,  $J=7.8\text{Hz}$ ), 7.37 (m, 3H), 7.31-7.23 (m, 7H), 6.41 (d, 1H,  $J=49.9\text{Hz}$ ), 4.90 (s, 2H), 3.08 (q, 2H,  $J=7.3\text{Hz}$ ), 1.43 (t, 3H,  $J=7.3\text{Hz}$ ). MS calcd 384.1; MS (M+1) 385.2.


#### Example 517

10 N-(3-Benzoylphenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline




To a solution of N-(3-( $\alpha$ -hydroxybenzyl)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline (0.717 g, 1.87 mmol) in 10 mL DCM was added activated  $\text{MnO}_2$  (0.489 g, 5.62 mmol).  
15 The reaction was gently heated and sonicated for 5 min then stirred overnight. Another portion of  $\text{MnO}_2$  (0.489 g, 5.62 mmol) was added and it was sonicated and stirred overnight again. The mixture was filtered through Celite and concentrated to give a pure white solid.  $^1\text{H}$  NMR: 8.52  
20 (dd,  $J=4.9, 1.5\text{Hz}$ , 1H), 8.40 (d,  $J=2.0\text{Hz}$ , 1H), 7.74-7.66 (m, 5H), 7.61 (m, 1H), 7.51-7.42 (m, 4H), 7.25 (dd,  $J=7.8, 4.9\text{Hz}$ , 1H), 4.93 (s, 2H), 3.14 (q,  $J=7.3\text{Hz}$ , 2H), 1.45 (t,  $J=7.3\text{Hz}$ , 3H). MS calcd. 380.1; MS (M+1) 381.1.

-129-

Example 518N-(4-Benzoylphenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline

By the method of Example 517 using N-(4-( $\alpha$ -  
5 hydroxybenzyl)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline  
the title compound was obtained.  $^1\text{H}$  NMR: 8.51  
(d, 1H, J=3.9Hz), 8.44 (br s, 1H), 7.79-7.71 (m, 4H), 7.59  
(m, 1H), 7.48 (m, 2H), 7.41 (d, 2H, J=8.3Hz), 7.26 (m, 2H), 4.99  
(s, 2H), 3.15 (q, 2H, J=7.3Hz), 1.46 (t, 3H, J=7.3Hz). MS calcd  
10 380.1; MS (M+1) 381.1.

Example 519N-(3-(1-Phenylvinyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline

15


To a suspension of methyltriphenylphosphonium bromide  
(0.269 g, 0.752) in 3 mL THF at 0 °C was added KHMDS (0.5 M  
in toluene, 1.5 mL, 0.752 mmol). The mixture was warmed to  
ambient temperature for 1 h and N-(3-benzoylphenyl)-N-  
20 (ethylsulfonyl)pyrid-3-ylmethylaniline (0.143 g, 0.376 mmol)  
as a solution in 3 mL THF was added via cannula. After 3 h  
the mixture was poured into saturated  $\text{NaHCO}_3$  solution and  
Extracted with EtOAc, dried over  $\text{MgSO}_4$ , filtered and

-130-

concentrated. The residue was chromatographed with 10:1 to 5:1 DCM/EtOAc to give the title compound as a clear oil. <sup>1</sup>H NMR: 8.50 (dd, J=4.9, 1.5 Hz, 1H), 8.35 (d, J=2.0 Hz, 1H), 7.70 (dt, J=7.8, 2.0 Hz, 1H), 7.37-7.19 (m, 9H), 7.14 (m, 1H), 5.48 (d, J=1.0 Hz, 1H), 5.39 (d, J=1.0 Hz, 1H), 4.85 (s, 2H), 3.09 (q, J=7.3 Hz, 2H), 1.41 (t, J=7.3 Hz, 3H). MS calcd. 378.1; MS (M+1) 379.1.

Example 520


10 N-(4-(1-Phenylvinyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine



By the method of Example 519 using N-(4-benzoylphenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylamine the title compound was obtained. <sup>1</sup>H NMR: 8.51 (d, 1H, J=3.9 Hz), 8.44 (br s, 1H), 7.79-7.71 (m, 4H), 7.59 (m, 1H), 7.48 (m, 2H), 7.41 (d, 2H, J=8.3 Hz), 7.26 (m, 2H), 4.99 (s, 2H), 3.15 (q, 2H, J=7.3 Hz), 1.46 (t, 3H, J=7.3 Hz). MS calcd 380.1; MS (M+1) 381.1.

Example 521

20 N-(3-(α-Methylbenzyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine




-131-

The title compound was prepared using a method similar to the method of Example 491 using N-[3-(1-phenylvinyl)phenyl]-N-(ethylsulfonyl)pyrid-3-ylmethanamine.  
<sup>1</sup>H NMR: 8.47 (dd, J=4.9, 1.5 Hz, 1H), 8.32 (d, J=2.0 Hz, 1H), 7.61 (dt, J=7.8, 2.0 Hz, 1H), 7.30-7.05 (m, 9H), 6.99 (m, 1H), 4.82 (s, 2H), 4.08 (q, J=7.3 Hz, 1H), 3.04 (q, J=7.3 Hz, 2H), 1.56 (d, J=7.3 Hz, 3H), 1.38 (t, J=7.3 Hz, 3H). MS calcd. 380.2; MS (M+1) 381.2.

10

Example 522

N-(4-(α-Methylbenzyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethanamine



The title compound was prepared using a method similar to the method of Example 491 using N-[3-(1-phenylvinyl)phenyl]-N-(ethylsulfonyl)pyrid-3-ylmethanamine.  
<sup>1</sup>H NMR: 8.49 (dd, 1H, J=4.4, 1.0 Hz), 8.35 (d, 1H, J=1.5 Hz), 7.72 (d, 1H, J=7.8 Hz), 7.31-7.11 (m, 10H), 4.86 (s, 2H), 4.10 (q, 1H, J=7.3 Hz), 3.07 (q, 2H, J=7.3 Hz), 1.59 (d, 3H, J=7.3 Hz), 1.41 (q, 3H, J=7.3 Hz). MS calcd 380.2; MS (M+1) 381.3.




-132-

Preparation 11N-(3-Iodo-4-methylphenyl)pyrid-3-ylmethylaniline

The title compound was prepared using a method similar to the method of Preparation 8 with 3-iodo-4-methylaniline as starting material. <sup>1</sup>H NMR: 8.61 (d, J=1.5Hz, 1H), 8.53 (dd, J=4.9, 2.0Hz, 1H), 7.67 (dt, J=7.8, 1.5Hz, 1H), 7.27 (ddd, J=7.8, 4.9, 1.0Hz, 1H), 7.15 (d, J=2.5Hz, 1H), 7.00 (d, J=8.3Hz, 1H), 6.52 (dd, J=7.8, 2.5Hz, 1H), 4.30 (d, J=5.4Hz, 2H), 3.97 (br s, 1H), 2.31 (s, 3H).

10

Example 523N-(3-Iodo-4-methylphenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline


The title compound was prepared using a method similar to the method of Example 467 using N-(3-iodo-4-methylphenyl)pyrid-3-ylmethylaniline. <sup>1</sup>H NMR: 8.50 (d, J=3.4Hz, 1H), 8.36 (d, J=1.6Hz, 1H), 7.69 (m, 2H), 7.26 (t, J=5.9Hz, 1H), 7.14 (d, J=7.8Hz, 1H), 7.08 (dd, J=7.8, 2.0Hz, 1H), 4.84 (s, 2H), 3.08 (q, J=7.3Hz, 2H), 1.42 (t, J=7.3Hz, 1H).

20

-133-

Example 524

N-(3-Formyl-4-methylphenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine




- 5 A mixture of N-(3-iodo-4-methylphenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylamine (0.415 g, 0.997 mmol), sodium formate (0.102 g, 1.50 mmol), and dichloro-bis(triphenylphosphine)palladium (0.014 g, 0.020 mmol) were combined in 1.0 mL DMF and purged with anhydrous CO gas.
- 10 The mixture was heated to 100 °C for 1.5 h. The reaction was poured into saturated NaHCO<sub>3</sub> solution and extracted with EtOAc. The combined fractions were washed with H<sub>2</sub>O and brine, dried over MgSO<sub>4</sub>, filtered and concentrated. The mixture was chromatographed on silica gel with 1:1 EtOAc/DCM
- 15 to give the title compound as a clear oil. <sup>1</sup>H NMR: 10.11 (s,1H), 8.38 (br s,1H), 8.26 (br s,1H). 7.60 (m,2H), 7.13 (m,1H), 7.00 (s,2H), 4.80 (s,2H), 3.00 (q,J=7.3Hz,2H), 2.52 (s,3H), 2.35 (t,J=7.3Hz,3H).

20

Example 525

N-(3-(α-Hydroxybenzyl)-4-methylphenyl)-N-ethanesulfonylpyrid-3-ylmethylamine




-134-

The title compound was prepared using a method similar to the method of Example 475 using N-(3-formyl-4-methylphenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline. <sup>1</sup>H NMR: 8.46 (br s, 1H), 8.30 (br s, 1H), 7.69 (br d, J=7.8Hz, 1H), 7.45 (s, 1H), 7.36-7.19 (m, 4H), 7.13 (d, J=6.3Hz, 2H), 7.06 (m, 2H), 5.87 (s, 1H), 4.91 (AofAB, d, J=14.7Hz, 1H), 4.82 (BofAB, d, J=14.7Hz, 1H), 3.08 (q, J=7.3Hz, 2H), 2.54 (br s, 1H), 2.13 (s, 3H), 1.41 (t, J=7.3Hz, 3H). MS calcd. 396.2; MS (M+1) 397.2.

10

Example 526

N-(3-Benzyl-4-methylphenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline




15 The title compound was prepared using a method similar to the method of Example 491 using N-(3-( $\alpha$ -hydroxybenzyl)-4-methylphenyl)-N-ethanesulfonylpyrid-3-ylmethylaniline. <sup>1</sup>H NMR: 8.49 (d, J=3.9, 1H), 8.32 (br s, 1H), 7.67 (br d, J=7.8Hz, 1H), 7.28-7.17 (m, 4H), 7.09 (d, J=7.8Hz, 1H), 7.02 (dd, J=7.8, 2.0Hz, 1H), 6.97 (d, J=7.4Hz, 2H), 6.88 (d, J=2.3Hz, 1H), 4.81 (s, 2H), 3.88 (s, 2H), 3.06 (q, J=7.3Hz, 2H), 2.18 (s, 3H), 1.39 (t, J=7.3Hz, 3H). MS calcd. 380.2; MS (M+1) 381.2.

20

-135-

Example 527

N-(3-(4-*t*-Butylphenoxy)-4-methylphenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline




The title compound was prepared following Example 342 using N-(3-iodo-4-methylphenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline and 4-*tert*-butylphenol and purifying via radial chromatography eluting with 50:50 to 53:47

THF/hexanes gave the title compound. <sup>1</sup>H NMR (400 MHz) 1.32 (9H, s), 1.39 (3H, t, J=7.8 Hz), 2.19 (3H, s), 3.07 (2H, q, J=7.8 Hz), 4.79 (2H, s), 6.70 (3H, m), 6.92 (1H, dd, J=2.0, 7.8 Hz), 7.16 (1H, d, J=7.8 Hz), 7.20-7.31 (3H, m), 7.67 (1H, d, J=8.3 Hz), 8.31 (1H, bs), 8.50 (1H, bs). HRMS calculated 439.2055 [M+H]<sup>+</sup> Found 439.2073.

Example 528

N-(3-(2-Methoxyphenoxy)benzyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline



The title compound was prepared following Example 1 using 3-(2-benzyloxyphenoxy)benzyl amine gave the title compound which was isolated as its HCl salt. <sup>1</sup>H NMR: 8.53 (s, 1H), 8.40 (s, 1H), 7.62 (d, 1H, J=8Hz), 7.14-7.24 (m, 4H),


-136-

6.85-7.00 (m, 4H), 6.77 (s, 1H), 4.39 (s, 2H), 4.30 (s, 2H), 3.80 (s, 3H), 3.56-3.58 (m, 3H). MS calcd. 466.48; MS (M+1) 467.9.

5

Example 529

N-(3-(2-Methoxyphenoxy)benzyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine




The title compound was prepared following Example 1 using 3-(2-benzyloxyphenoxy)benzyl amine gave the title compound which was isolated as its HCl salt. <sup>1</sup>H NMR: 8.51 (d, 1H, J=4Hz), 8.39 (s, 1H), 7.65 (d, 1H, J=8Hz), 7.13-7.24 (m, 4H), 6.84-7.02 (m, 4H), 6.78 (s, 1H), 4.34 (s, 2H), 4.25 (s, 2H), 3.81 (s, 3H), 2.92-2.96 (m, 2H), 1.28 (t, 3H, J=8Hz). MS calcd. 412.5; MS (M+1) 413.6.

15

Example 530

N-(3-(2-Phenylvinyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine



20

Benzyltriphenylphosphonium bromide (0.765 g, 1.77 mmol) was dispersed into THF (12 ml) and cooled at 0°C. To the cooled suspension was added dropwise a solution of potassium bis(trimethylsilyl)amide (0.5 M, 3.5 ml, 1.75 mmol). After stirring at 0 °C for 1 h, a solution of ) . N-

25


-137-

(3-formyl-phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylamine (0.304 g, 1.0 mmol) in THF was added. The reaction mixture was stirred at ambient temperature for 2 h. Evaporation and chromatography on silica-gel eluting with toluene-EtOAc to  
5 give the title compound.  $^1\text{H}$  NMR: 8.48 (d, 1H,  $J=3.4\text{Hz}$ ), 8.30 (d, 1H,  $J=2.0\text{Hz}$ ), 7.65 (d, 1H,  $J=7.8\text{Hz}$ ), 7.30-6.74 (m, 8H), 6.60 (q, 2H,  $J=12.2, 37.1\text{Hz}$ ), 4.68 (s, 2H), 2.87 (q, 2H,  $J=7.3\text{Hz}$ ), 1.28 (t, 3H,  $J=7.4\text{Hz}$ ). MS calcd 378.1; MS (M+1) 379.1.

10

Example 531

N-(3-(2-Phenylethyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine




N-(2-phenylvinylphenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylamine (0.50 g, 0.13 mmol) was dissolved in ethanol (5 ml), followed by addition of 5% platinum-on-carbon (0.38 g). After flushing with hydrogen, the mixture was stirred under hydrogen-ballon at ambient temperature overnight and filtered. Upon evaporation, it underwent chromatography on  
20 silica-gel with eluent of toluene and EtOAc and gave 0.32 g (64% yield).  $^1\text{H}$  NMR: 8.48 (d, 1H,  $J=3.4$ ), 8.34 (s, 1H), 7.68 (s, 1H), 7.30-6.86 (m, 10H), 4.81 (s, 2H), 3.04-2.82 (m, 6H), 1.38 (t, 3H,  $J=7.3\text{Hz}$ ). MS calcd 380.2; MS (M+1) 381.1.

-138-

Example 532


N-(3-(Prop-1-en-1-yl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethamphetamine



5       The title compound was prepared using a method similar to the method of Example 530 using N-(3-formylphenyl)-N-(ethylsulfonyl)pyrid-3-ylmethamphetamine. <sup>1</sup>H NMR: 8.39 (d, 1H, J=1.5Hz), 8.30 (s, 1H), 7.62 (d, 1H, J=8.3Hz), 7.22-6.80 (m, 5H), 6.25 (d, 1H, J=10.3), 5.70 (m, 1H), 4.80 (s, 1H), 3.01  
10 (q, 2H, 7.4Hz), 1.72 (d, 3H, J=6.4Hz), 1.34 (t, 3H, J=7.3Hz). MS calcd 316.1; MS (M+1) 317.1.


Example 533

15       N-(3-(2-N,N-Diethylamidovinyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethamphetamine




The title compound was prepared using a method similar to the method of Example 530 using N-(3-formylphenyl)-N-(ethylsulfonyl)pyrid-3-ylmethamphetamine. <sup>1</sup>H NMR: 8.41  
20 (d, 1H, J=3.4Hz), 8.32 (s, 1H), 7.70-7.03 (m, 9H), 6.61 (d, 1H, 15.1Hz), 4.83 (s, 1H), 3.39 (m, 2H), 3.05 (m, 2H), 1.36 (t, 3H, J=7.3Hz), 1.26-1.04 (m, 6H). MS calcd 401.2; MS (M+1) 402.2.

-139-

Example 534N-(3-(2-Ethoxycarbonylvinyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethamphetamine

5 The title compound was prepared using a method similar to the method of Example 530 using N-(3-formylphenyl)-N-(ethylsulfonyl)pyrid-3-ylmethamphetamine. <sup>1</sup>H NMR: 8.50 (d, 1H, J=1.5Hz), 8.37 (s, 1H), 7.80-7.50 (m, 2H), 7.30-7.10 (m, 4H), 4.90 (s, 2H), 4.25 (q, 2H, J=7.3Hz), 3.10 (q, 2H, 7.3Hz), 1.50-1.28 (m, 6H). MS calcd 374.1; MS (M+1) 375.1.

Example 535N-(3-Propylphenyl)-N-(ethanesulfonyl)pyrid-3-ylmethamphetamine


15 The title compound was prepared using a method similar to the method of Example 431. <sup>1</sup>H NMR: 8.48 (d, 1H, J=3.4Hz), 8.35 (d, 1H, J=1.5Hz), 7.70 (d, 1H, 7.8Hz), 7.21 (m, 2H), 7.05 (m, 3H), 4.87 (s, 2H), 3.10 (q, 2H, J=7.8Hz), 2.50 (t, 2H, J=7.3Hz), 1.75-1.20 (m, 6H), 0.85 (t, 3H, J=7.3Hz). MS calcd 318.1; MS (M+1) 319.1.



-140-

Example 536

N-(3-Trifluoromethoxyphenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline




5           3-(Trifluoromethoxy)aniline (0.602 g, 3.4 mmol) and pyridine-3-carboxaldehyde (0.400g, 3.7 mmol) were dissolved in methanol (10 ml). The solution was refluxed for 3 h and then cooled in an ice-bath. To the solution was added sodium borohydride (0.212 g, 5.5 mmol) in portions. The  
10 reaction mixture was stirred at ambient temperature overnight. After addition of small pieces of ice, the solvent was evaporated to near dryness. The residue was dissolved in cold water and extracted with ethyl acetate. The organic phase was washed with brine and dried over  $\text{MgSO}_4$ .  
15 Filtration and evaporation afforded N-(3-trifluoromethoxyphenyl)pyrid-3-ylmethylaniline.  $^1\text{H}$  NMR: 8.43-8.44 (m, 2H), 7.60 (d, 1H), 7.00-7.30 (m, 3H), 6.20-6.40 (m, 2H), 4.27 (d, 5.4 Hz). MS calcd 268.1; MS (M+1) 269.0.

          N-(3-trifluoromethoxyphenyl)pyrid-3-ylmethylaniline (0.270  
20 g, 1 mmol) and pyridine were dissolved in dichloroethane (2 ml). The solution was cooled in an ice-bath. To the solution was added 2,2,2-trifluoroethylsulfonyl chloride (0.274 g, 1.5 mmol). The mixture was stirred at room temperature overnight. Upon removing solvent, the residue  
25 was dissolved in water and extracted with EtOAc. The organic phase was washed with brine concentrated. The residue was chromatographed to afford the title compound.  $^1\text{H}$  NMR: 8.42 (d, 1H,  $J=6.4\text{ Hz}$ ), 8.20 (d, 1H,  $J=2.0\text{ Hz}$ ), 7.57

-141-

(d, 1H, J=9.8Hz), 7.40-6.85 (m, 5H), 4.81 (s, 2H), 3.72 (q, 2H, J=7.8Hz). MS calcd 414.1; MS (M+1) 415.1.

By a method similar to the method of Example 536 the following compounds were prepared:




No.	R1	R2	Data
537	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	3-trifluoromethoxy	<sup>1</sup> H NMR: 8.42 (s, 1H), 8.29 (s, 1H), 7.60 (d, 1H), 7.00-7.30 (m, 5H), 4.81 (s, 2H), 3.01 (q, 2H, J=7.3Hz), 1.35 (t, 3H, J=7.3). MS calcd 360.1; MS (M+1) 361.2.
538	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-trifluoromethyl	<sup>1</sup> H NMR: 8.43 (d, 1H, J=3.9Hz), 8.26 (s, 1H), 7.56 (m, 3H), 7.00-7.30 (m, 3H), 4.85 (s, 2H), 3.72 (q, 2H, J=8.3Hz). MS calcd 398.05; MS (M+1) 399.1
539	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-trifluoromethoxy	<sup>1</sup> H NMR: 8.44 (d, 1H, J=3.9Hz), 8.25 (d, 1.5Hz), 7.59 (d, 1H, J=7.8Hz), 5.00-7.30 (m, 5H), 4.80 (s, 2H), 3.72 (q, 2H, J=8.8Hz). MS calcd 414.1; MS (M+1) 415.1
540	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	4-trifluoromethoxy	<sup>1</sup> H NMR: 8.42 (d, 1H, J=3.4Hz), 8.29 (d, 1H, J=2.0Hz), 7.61 (d, 1H, J=7.8Hz), 7.00-7.25 (m, 5H), 4.79 (s, 2H), 3.01 (q, 2H, J=7.3Hz), 1.35 (t, 3H, J=7.3Hz). MS calcd 360.1; MS (M+1) 361.2
541	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-methoxy	<sup>1</sup> H NMR: 8.41 (d, 1H, J=3.4Hz), 8.26 (s, 1H), 7.59 (d, 1H, J=7.8Hz), 7.17 (m, 2H), 6.60-6.86 (m, 3H), 4.79 (s, 2H), 3.50-3.82 (m, 5H). MS calcd 360.1; MS (M+1) 361.1
542	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	4-(prop-2-yl)	<sup>1</sup> H NMR: 8.42 (d, 1H, J=3.9Hz), 8.23 (1H), 7.62 (d, 1H, J=7.8Hz), 6.96-7.26 (m, 5H), 4.78 (s, 2H), 3.71 (q, 2H, J=9.3Hz), 2.79 (quint, 1H, J=7.3Hz), 1.13 (d, 6H, J=7.8Hz). MS calcd 372.1; MS (M+1) 373.2.
543	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	2-trifluoromethoxy	<sup>1</sup> H NMR: 8.44 (m, 1H), 8.18 (d, 1H, J=2.0Hz), 7.62 (d, 1H, J=7.8Hz), 6.80-7.40 (m, 5H), 3.80 (q, 2H, J=9.3Hz).

-142-

			MS calcd 414.1; MS (M+1) 415.1
544	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	3-(prop-2-yloxy)	<sup>1</sup> H NMR: 8.41 (d,1H,J=3.4Hz), 8.26 (d,1H,J=1.5Hz), 7.59 (d,1H,J=7.8Hz), 7.10-7.22 (m,2H), 6.56-6.80 (m, 3H), 4.79 (s,2H), 4.34 (m,1H,J=5.9Hz), 3.73 (q,2H,J=9.2Hz),1.17 (d,6H,J=5.9Hz ). MS calcd 388.1; MS (M+1) 389.2

Example 545

N-(3-Benzyl-4-chlorophenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylamine



2-Chloro-5-nitrobenzophenone (1.3 g, 5.0 mmol), tin(II) chloride (4.5 g, 20 mmol), and ethanol (20 ml) were mixed and heated at 70 °C for 1 h. After cooling to room temperature, to the reaction mixture were added ice, EtOAc and aqueous NaHCO<sub>3</sub> and it was stirred for 30 min. The organic layer was washed with brine and dried over MgSO<sub>4</sub>. Filtration and evaporation afforded 5-amino-2-chlorobenzophenone. <sup>1</sup>H NMR, 7.85 (m, 2H), 7.30-7.66 (m, 3H), 7.20 (d, 1H, J=8.8Hz), 6.60-6.80 (m, 2H). MS calcd 231.1; MS (M+1) 232.0.

5-Amino-2-chlorobenzophenone (1.2 g, 5.2 mmol) and pyridine-3-carboxaldehyde (0.642 g, 6 mmol) were dissolved in methanol. The solution was refluxed for 2 h and then cooled in ice-bath. To the cold solution was added sodium borohydride (0.741 g, 19.5 mmol) in portions. The reaction mixture was stirred at ambient temperature overnight. After addition of small pieces of ice, most of the solvent was evaporated. The residue was dissolved in water and

-143-

extracted with EtOAc. The combined organic layer was washed with brine and dried over  $\text{MgSO}_4$ . Upon filtering and evaporation, solid precipitated, when chloroform was added to the oil. After cooling for a while, filtration and drying gave N-(3-( $\alpha$ -hydroxybenzyl)-4-chlorophenyl)pyrid-3-methylamine.  $^1\text{H}$  NMR, 8.40-8.70 (m, 2H), 7.65 (d, 1H), 7.00-7.50 (m, 7H), 6.83 (d, 1H,  $J=2.5\text{Hz}$ ), 6.43 (d, 1H), 6.14 (s, 1H). MS calcd 324.1; MS (M+1) 325.1.

N-(3-( $\alpha$ -hydroxybenzyl)-4-chlorophenyl)pyrid-3-methylamine (0.105 g, 0.32 mmol) was dispersed into methylene chloride (5 ml) and triethylsilane (1 ml). The mixture was cooled in an ice-bath. To the cooled suspension was added trifluoroacetic acid (0.6 ml) and then stirred overnight. The reaction mixture was cooled at 0 °C, to which was added ice, aqueous  $\text{NaHCO}_3$  and chloroform. The organic phase was washed with brine and evaporated to dryness. Chromatography afforded N-(3-benzyl-4-chlorophenyl)pyrid-3-ylmethylamine.  $^1\text{H}$  NMR: 8.53 (d, 2H,  $J=14\text{Hz}$ ), 7.19 (d, 1H,  $J=8\text{Hz}$ ), 7.00-7.40 (m, 8H), 6.20-6.40 (m, 2H), 4.27 (s, 2H), 3.99 (s, 2H). MS calcd 308.1; MS (M+1) 309.1.

N-(3-benzyl-4-chlorophenyl)pyrid-3-ylmethylamine (0.04 g, 0.13 mmol) and pyridine (1 ml) was dissolved in DCM. The reaction was cooled in an ice-bath to which was added dropwise 2,2,2-trifluoroethylsulfonyl chloride (0.59 g, 0.32 mmol). The mixture was stirred at room temperature overnight. After evaporation of solvent, the residue was dissolved in EtOAc and washed with cold  $\text{NaHCO}_3$  and brine, and evaporated to dryness. Chromatography gave the title compound.  $^1\text{H}$  NMR: 8.51 (d, 1H,  $J=3.9\text{Hz}$ ), 8.27 (s, 1H), 7.59 (d, 1H,  $J=7.8\text{Hz}$ ), 7.34 (t, 1H,  $J=8.3\text{Hz}$ ), 7.28 (d, 2H,  $J=7.8\text{Hz}$ ), 7.21 (m, 2H), 8.03 (m, 3H), 6.89 (d, 1H,  $J=2.4\text{Hz}$ ), 4.78 (s, 2H), 4.01 (s, 2H), 3.73 (q, 2H,  $J=9.3\text{Hz}$ ). MS calcd 454.1; MS (M+1) 455.0.


-144-

Preparation 12N-(4-(3-Hydroxyphenyl)pyrid-3-ylmethylamine

3-Pyridinecarboxaldehyde (0.550 g, 5 mmol) and 3-aminophenol (0.540 g, 5 mmol) were dissolved into methanol (15 ml). To the solution was added a mixture of sodium cyanoborohydride (0.378 g, 6 mmol) and zinc chloride (0.408 g, 3.0 mmol) in methanol at room temperature. The reaction mixture was stirred for 2.5 h. Most of the methanol was evaporated under reduced pressure. Ice water was added to the residue. It was filtered and the layers were separated. The aqueous layer was extracted with EtOAc. The combined organic layers were washed with brine, dried over  $\text{MgSO}_4$ , filtered, and concentrated to give the title compound.  $^1\text{H}$  NMR: 8.53 (s, 1H), 8.45 (d, 1H,  $J=3.9\text{Hz}$ ), 7.72 (d, 1H,  $J=7.8\text{Hz}$ ), 7.24-7.30 (m, 2H), 6.95-7.01 (m, 1H), 4.27 (s, 2H). MS calcd. 200.1; MS (M+1) 201.1.

Example 546

N-(4-(3-Cyclobutyloxyphenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylamine




Sodium hydride (60% in mineral oil, 0.072 g, 1.8 mmol) was suspended in DMF at 0 °C. To the dispersion was added dropwise N-(3-hydroxyphenyl)pyrid-3-ylmethylamine (0.250 g, 1.24 mmol) as a solution in DMF. The reaction mixture was stirred at room temperature for 30 min. Then bromocyclobutane (0.25 g, 1.84 mmol) was added as a solution in DMF. The reaction was concentrated and the residue was diluted with water, extracted with EtOAc, washed with brine,

-145-

dried over  $\text{MgSO}_4$ , filtered and concentrated. The residue was purified by silica-gel chromatography to give N-(4-(3-Cyclobutyloxyphenyl)pyrid-3-yl)methylamine.  $^1\text{H}$  NMR: , 8.61 (d, 1H,  $J=2.3\text{Hz}$ ), 8.51 (q, 1H,  $J=1.6, 3.13$ ), 7.68 (m, 1H), 7.25 (m, 1H), 7.04 (t, 1H,  $1.7\text{Hz}$ ), 6.18-6.24 (m, 2H), 6.09 (t, 1H,  $J=2.3\text{Hz}$ ), 4.6 (quint, 1H,  $J=6.7\text{Hz}$ ), 2.33-2.42 (m, 2H), 2.06-2.18 (m, 2H), 1.76-1.88 (m, 1H), 1.57-1.70 (m, 1H). MS calcd. 254.1; MS (M+1) 255.1.

To a solution of give N-(4-(3-Cyclobutyloxyphenyl)pyrid-3-yl)methylamine (0.180 g, 0.71 mmol) in DCM (5 ml) and pyridine (1 ml) was dropped a solution of 2,2,2-trifluoroethylsulfonyl chloride (0.156 g, 0.85 mmol) in DCM (1ml) at  $0^\circ\text{C}$ . The reaction mixture was stirred overnight ambient temperature. After evaporation of solvent, the residue was diluted with ice water and extracted with EtOAc. The organic layer was washed with brine, evaporated and chromatographed on silica gel eluting with chloroform / ethyl acetate to give the title compound.  $^1\text{H}$  NMR: 8.50 (d, 1H,  $J=4.7\text{Hz}$ ), 8.34 (s, 1H), 7.65-7.69 (m, 1H), 7.21-7.26 (m, 2H), 6.74-6.80 (m, 2H), 6.63 (t, 1H,  $J=2.0\text{Hz}$ ), 4.53 (q, 1H,  $J=7.0\text{Hz}$ ), 3.80 (q, 2H,  $J=8.0\text{Hz}$ ), 2.82-2.90 (m, 2H), 2.03-2.15 (m, 2H), 1.77-1.89 (m, 1H), 1.60-1.73 (m, 2H). MS calcd. 400.1; MS (M+1) 401.2.

By a method similar to the method of Example 546 the following compounds were prepared:



No.	R2	Data
547	3-isobutyloxy	$^1\text{H}$ NMR: 8.51 (m, 1H), 8.35 (d, 1H, $J=2.0\text{Hz}$ ), 7.68


		(m,1H), 7.25 (m,2H), 6.66-6.90 (m,3H), 4.88 (s,2H), 3.80 (q,2H,J=9.0Hz), 3.61 (d,2H,J=5.7Hz), 2.01 (m,1HJ=6.6Hz), 1.00 (d,6H,J=6.5Hz). MS calcd ; MS (M+1)
548	3-sec-butoxyl	<sup>1</sup> H NMR: 8.48 (d,1H,J=3.5Hz), 8.35 (s,1H), 7.67 (m,1H), 7.20-7.28 (m,2H), 6.60-6.90 (m,3H), 4.87 (s,2H), 4.17 (m,1H,J=6.2Hz), 3.82 (q,2H,J=9.0Hz), 1.45-1.75 (m, 2H), 1.20 (d,3H,J=5.9Hz), 0.92 (t,3H,J=7.4Hz). MS calcd 402.1; MS (M+1) 403.2
549	3-cyclopentyloxy	<sup>1</sup> H NMR: 8.52 (d,1H,J=4.3Hz), 8.35 (s,1H), 7.68 (m,1H), 7.26 (m, 2H), 6.60-6.90 (m, 3H), 4.87 (s,2H), 4.62 (m,2H), 3.82 (q,2H,J=9.0Hz), 1.40-2.00 (m,8H). MS calcd 414.1; MS (M+1) 415.2
550	3-cyclohexyloxy	<sup>1</sup> H NMR: 8.20-8.40 (m,2H), 7.69 (m,1H), 7.25 (m,2H), 6.60-6.94 (m,3H), 4.87 (s,2H), 4.11 (m,1H), 3.83 (q,2H,J=8.8Hz), 1.10-2.00 (m,10H). MS calcd 428.1; MS (M+1) 429.1
551	3-(1-ethoxycarbonyl-1,1-dimethyl methoxy)	<sup>1</sup> H NMR: 8.49 (m,1H), 8.33 (d,1H,J=1.9Hz), 7.66 (m,1H),7.24 (m,2H), 6.60-6.90 (m,3H), 4.85 (s,2H), 4.17 (q,2H,J=7.5Hz), 3.80 (q,2H,J=9.4Hz), 1.52 (s,6H), 1.20 (t,3H,J=7.0Hz). MS calcd 460.1; MS (M+1) 461.1.
552	3-(1-ethoxycarbonyl-1-methylmethoxy)	<sup>1</sup> H NMR: 8.51 (m,1H), 8.35 (d,1H,J=1.9Hz), 7.66 (m,1H), 7.24 (m,2H), 6.82 (m,3H), 4.87 (s,2H), 4.64 (q,1H,J=6.6Hz), 4.18 (q,2H,J=7.0Hz), 3.80 (q,2H,J=9.0Hz), 1.60 (s,6H), 1.23 (t,3H,J=8.9Hz). MS calcd 446.1; MS (M+1) 447.0.
553	3-(1-(ethoxycarbonyl)-cyclobut-1-yloxy	<sup>1</sup> H NMR: 8.50 (m,1H), 8.32 (d,1H,J=2.0Hz), 7.66 (m,1H), 7.25 (m,2H), 6.50-6.90 (m,3H), 4.85 (s,2H), 3.88 (q,2H,J=7.3Hz), 3.78 (q,2H,J=9.0Hz), 2.68 (m,2H), 2.36 (m,2H), 2.00 (m,2H), 1.13 (t,3H,J=7.0Hz). MS calcd 472.1; MS (M+1) 473.1.
554	3-pent-3-yloxy	<sup>1</sup> H NMR: 8.51 (m,1H), 8.35 (d,1H,J=2.0Hz), 7.67 (m,1H), 7.24 (m,2H), 6.60-6.90 (m,3H), 4.87 (s,2H), 3.98 (m,1H,J=5.5Hz), 3.82 (q,2H,J=9.3Hz), 1.59 (m,4H), 0.89 (t,6H,J=7.4Hz). MS calcd 416.1; MS (M+1) 417.1
555	3-pent-2-yloxy	<sup>1</sup> H NMR: 8.50 (m,1H), 8.35 (d,1H,J=2.0Hz), 7.67 (m,1H), 7.23 (m,2H), 6.60-6.90 (m,3H), 4.87 (s,2H), 4.24 (m, 1H,J=5.9Hz), 3.82 (q,2H,J=9.0Hz), 1.27-1.80 (m,4H), 1.21 (d,3H,J=5.9Hz), 0.90 (t,3H,J=7.1Hz). MS calcd 416.1; MS (M+1) 417.2
556	3-(2,2,2-trifluoroethoxy)-	<sup>1</sup> H NMR: 8.52 (m,1H), 8.35 (d,1H,J=1.6Hz), 7.67 (m,1H), 7.26 (m,2H), 6.70-6.98 (m,3H), 4.89 (s,2H), 4.26 (q,2H,J=7.8Hz), 3.81 (q,2H,J=8.4Hz). MS calcd 428.1; MS (M+1) 429.1
557	3-(2-methyl-	<sup>1</sup> H NMR: 8.51 (s,1H), 8.35 (d,1H,J=2.4Hz), 7.67

-147-

	butyloxy)	(m,1H), 7.24 (m,2H), 6.60-6.90 (m,3H), 4.88 (s,2H), 3.80 (q,2H,J=9.0Hz), 3.62 (m,2H), 1.78 (m,1H), 1.52 (m,1H), 1.23 (m,1H), 0.95 (m,6H). MS calcd 416.1; MS (M+1) 417.2
558	3-(ethoxycarbonyl methoxy)	<sup>1</sup> H NMR: 8.50 (d,1H,J=4.9Hz), 8.35 (d,1H,J=2.0Hz), 7.65 (m,1H), 7.26 (m, 2H), 6.60-6.90 (m,3H), 4.87 (s,2H), 4.53 (s,2H), 4.24 (q,2H,J=7.3Hz), 3.80 (q,2H,J=8.8Hz), 1.27 (t,3H,J=7.3Hz). MS calcd 432.1; MS (M+1) 433.0

Example 559

N-(3-(2-Fluoro-4-cyanophenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline



5

Prepare by the method of Example 342 using 3,4-difluoro-4-cyanobenzene and N-(3-hydroxyphenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, purifying via radial chromatography eluting with 50:50 to 60:40 THF/hexanes, to give the title compound. Anal Calcd for C<sub>21</sub>H<sub>18</sub>FN<sub>3</sub>O<sub>3</sub>S: C, 61.30; H, 4.41; N, 10.21. Found: C, 61.32; H, 4.50; N, 10.07. MS found 412.0 [M+H]<sup>+</sup>.


10



-148-

Example 560

N-(4-(2-Fluoro-4-amidophenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline



5           A solution of N-(4-hydroxyphenyl)-3-pyridylmethylaniline (1.0 equiv.) in DMF (0.1-0.3M) was treated with sodium hydride (1.0-1.5 equiv.) (60% dispersion in mineral oil). The reaction mixture was allowed to stir at room temperature for 30 minutes. The solution was then treated with 3,4-  
10   difluorobenzonitrile (1.6 equiv.) and allowed to stir. When complete (about 3 hours) the reaction mixture was partitioned between ethyl acetate/0.1M K<sub>2</sub>CO<sub>3</sub>, the organic layer separated, and the aqueous layer extracted 3 times with ethyl acetate. The organic layers were combined,  
15   washed with brine, and dried over MgSO<sub>4</sub>. The organic phase was concentrated in vacuo and purified by recrystallization from ethyl acetate/hexanes to give N-(4-(2-fluoro-4-cyanophenoxy)phenyl)-3-pyridylmethyl-amine <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 4.29 (2H, d, J=5.9 Hz), 6.40 (1H, t, J=5.5 Hz),  
20   6.63-6.65 (2H, m), 6.83 (1H, t, J=8.6 Hz), 6.90-6.92 (2H, m), 7.33-7.36 (1H, m), 7.55-7.57 (1H, m), 7.74-7.76 (1H, m), 7.93-7.97 (1H, m), 8.43-8.44 (1H, m), 8.58 (1H, s). MS Found 320.1 [M+H]<sup>+</sup>.

25           A solution (0.1M) of N-(4-(2-fluoro-4-cyanophenoxy)phenyl)-3-pyridylmethylaniline (1.0 equiv.) in 85% v/v H<sub>2</sub>SO<sub>4</sub> was allowed to stir at room temperature for 18 to 24 hours. The reaction mixture was poured over ice and neutralized with an appropriate amount of concentrated NaOH.

-149-

The mixture was extracted 3 times with ethyl acetate, washed with brine, dried over  $\text{MgSO}_4$ , and concentrated in vacuo to give N-(4-(2-fluoro-4-amidophenoxy)phenyl)-3-pyridylmethylamine as a crystalline solid.  $^1\text{H}$  NMR(400MHz,  $\text{DMSO-d}_6$ ):

5 4.28(2H, d,  $J=5.9$  Hz), 6.31(1H, t,  $J=6.1$  Hz), 6.61-6.64(2H, m), 6.81(1H, t,  $J=8.6$  Hz), 6.84-6.87(2H, m), 7.33-7.36(1H, m), 7.38(1H, s), 7.60-7.63(1H, m), 7.73-7.80(2H, m), 7.93(1H, s), 8.43-8.44(1H, m), 8.58(1H, d,  $J=2.0$  Hz).

MS Found 338.1  $[\text{M}+\text{H}]^+$ .

10 A solution of N-(4-(2-fluoro-4-amidophenoxy)phenyl)-3-pyridylmethylamine (1.0 equiv.) in DCM / pyridine(1/1 v/v) (0.2M) was cooled to  $0^\circ\text{C}$  and treated with 2,2,2-trifluoroethanesulfonyl chloride (1.2 equiv.). The mixture was allowed to stir while the temperature was maintained at  
15  $0-2^\circ\text{C}$ . The reaction was complete in approximately 5 minutes. The reaction mixture was partitioned between ethyl acetate/0.1 M  $\text{K}_2\text{CO}_3$ , the organic layer separated, and the aqueous layer extracted 2-4 times with ethyl acetate. The organic layers were combined, washed with brine, and dried  
20 over  $\text{MgSO}_4$ . The organic phase was concentrated in vacuo and purified via preparative HPLC eluting with methanol(2%) / ethyl acetate. The title compound was afforded by recrystallization from ethyl acetate/hexanes. Anal Calcd for  $\text{C}_{21}\text{H}_{17}\text{F}_4\text{N}_3\text{O}_4\text{S}$ : C, 52.17; H, 3.54; N, 8.69. Found: C, 51.78;  
25 H, 3.47; N, 8.45. MS found 484.1  $[\text{M}+\text{H}]^+$ .


The HCl salt was obtained by combining N-(4-(2-fluoro-4-amidophenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylamine (1 equiv.) in acetonitrile (0.4M) and adding 1.1 equiv. of HCl (0.27M), the solution was cooled to  
30  $-78^\circ\text{C}$ , and lyophilized over a 72 hour period to afford the HCl salt. Anal Calcd for  $\text{C}_{21}\text{H}_{17}\text{F}_4\text{N}_3\text{O}_4\text{S}\cdot\text{HCl}\cdot 0.8\text{H}_2\text{O}$ : C, 47.21; H, 3.70; N, 7.86; Cl, 6.64. Found: C, 46.97; H, 3.56; N, 7.74; Cl, 7.03.

-150-

<sup>1</sup>H NMR(400 MHz, DMSO-d<sub>6</sub>): 4.83(2H, q, J=9.8 Hz), 5.05(2H, s), 7.00-7.02(2H, m), 7.20(1H, t, J=8.6 Hz), 7.38-7.40(2H, m), 7.51(1H, s), 7.73-7.75(1H, m), 7.80-7.84(1H, m), 7.85-7.88(1H, m), 8.06(1H, s), 8.24(1H, d, J=8.1 Hz), 8.67(1H, s), 8.71(1H, d, J=4.8 Hz). MS found 484.1 [M+H]<sup>+</sup>.

## Example 561

N-(3-(2-Fluoro-4-cyanophenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline



10

Prepare by the method of Example 342 using N-(3-hydroxyphenyl)-3-pyridylmethylaniline and 3,4-difluoro-4-cyanobenzene gave N-(3-(2-fluoro-4-cyanophenoxy)phenyl)pyrid-3-ylmethylaniline which was purified by recrystallization from ethyl acetate/hexanes. <sup>1</sup>H NMR(400 MHz): 4.25(1H, s), 4.36(2H, d, J=5.5 Hz), 6.30(1H, t, J=2.0 Hz), 6.38(1H, m), 6.48(1H, m), 6.96(1H, t, J=8.2 Hz), 7.17(1H, t, J=8.20 Hz), 7.28(1H, m), 7.33(1H, d, J=8.6 Hz), 7.44(1H, m), 7.67(1H, d, J=7.8 Hz), 8.55(1H, s), 8.61(1H, s). MS Found 320.2 [M+H]<sup>+</sup>.

20

Sulfonation by the method of Example 526 in 1.8:1 dichloroethane/pyridine (0.25M) and raising the temperature after addition of to 45°C for 1 hour and 15 minutes after addition of the 2,2,2-trifluoroethanesulfonyl chloride. Next, 50 mg of anhydrous K<sub>2</sub>CO<sub>3</sub> was added to the reaction mixture and it was allowed to stir for 45 minutes. The mixture was next filtered through Celite®(521) and concentrated in vacuo. The crude material was


25

-151-

chromatographed via normal phase radial chromatography eluting with 1:1 to 0:1 hexanes/THF. The material was chromatographed a second time via normal phase radial chromatography eluting with 30:70 acetone/hexanes to give the title compound. Anal Calcd for  $C_{21}H_{15}F_4N_3O_3S$ : C, 54.19; H, 3.25; N, 9.03. Found: C, 54.28; H, 3.30; N, 8.97. MS found 466.1  $[M+H]^+$ .

## Example 562

10 N-(3-(2,4-Difluoro-4-cyanophenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylamine



Prepare by the method of Example 351 using N-(3-hydroxyphenyl)-3-pyridylmethylamine and 3,4,5-difluoro-4-cyanobenzene gave N-(3-(2,4-difluoro-4-cyanophenoxy)phenyl)pyrid-3-ylmethylamine which was purified by recrystallization from ethyl acetate/hexanes.  $^1H$  NMR(400 MHz, DMSO): 4.24(2H, d,  $J=5.9$  Hz), 6.11(1H, m), 6.15(1H, t,  $J=2.2$  Hz), 6.35(1H, m), 6.50(1H, t,  $J=5.9$  Hz), 7.00(1H, t,  $J=8.1$  Hz), 7.32(1H, m), 7.67(1H, m), 8.00(2H, m), 8.43(1H, d,  $J=3.4$  Hz), 8.51(1H, s). MS Found 338.2  $[M+H]^+$ .


The title compound was prepared following the procedure Example 526 using N-(3-(2,5-difluoro-4-cyanophenoxy)phenyl)-3-pyridylmethylamine and 2,2,2-trifluoroethanesulfonyl chloride in 2:1 dichloroethane/pyridine (0.23M) and the temperature was raised to 40°C for 4 hours after addition of 2,2,2-trifluoroethanesulfonyl chloride. Next, 50 mg of anhydrous  $K_2CO_3$  was added to the reaction mixture and it was

-152-

allowed to stir for 40 minutes. The mixture was next filtered through Celite®(521) and concentrated in vacuo. The crude material was chromatographed via normal phase radial chromatography eluting with 20:80 ethyl acetate/dichloromethane gave the title compound. Anal Calcd for  $C_{21}H_{14}F_5N_3O_3S$ : C, 52.18; H, 2.92; N, 8.69. Found: C, 52.12; H, 2.84; N, 8.61. MS found 484.0  $[M+H]^+$ .

Example 563

10 N-(4-(2-Fluoro-4-cyanophenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline




Prepare by the method of Example 351 using N-(4-hydroxyphenyl)-N-(ethanesulfonyl)-3- and 3,4-difluorobenzonitrile and purifying via radial chromatography eluting with 50:50 to 60:40 THF/hexanes. Anal Calcd for  $C_{21}H_{18}FN_3O_3S$ : C, 61.30; H, 4.41; N, 10.21. Found: C, 61.32; H, 4.50; N, 10.07.

-153-

Example 564


N-(4-(2-Fluoro-4-amidophenoxy)phenyl)-N-  
(ethanesulfonyl)pyrid-3-ylmethylamine



- 5        Prepare by the method of Example 560 using N-(4-(2-fluoro-4-cyanophenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine and 85% aqueous H<sub>2</sub>SO<sub>4</sub> (0.1M). The resultant solution was stirred for 3 days. (The reaction does not take 3 days and can be monitored by TLC). The pH of the  
10    reaction was adjusted to 10 with 5 N NaOH followed by saturated aqueous NaHCO<sub>3</sub> and extracted twice with ethyl acetate. The combined extracts were washed with brine, dried over sodium sulfate, filtered, and concentrated in vacuo. The crude product was purified via reverse-phase  
15    chromatography affording the HCl salt. Anal Calcd for C<sub>21</sub>H<sub>20</sub>FN<sub>3</sub>O<sub>4</sub>S•0.9HCl: C, 54.56; H, 4.56; N, 9.09. Found: C, 54.72; H, 4.55; N, 8.95. MS found 430.4 [M+H]<sup>+</sup>

Example 565

- 20    N-(3-(Bromomethyl)phenyl)-N-(ethanesulfonyl)pyrid-3-  
ylmethylamine




-154-

Resin-supported triphenylphosphine (1.6 g, 3 mmol per gram resin) was dispersed on dichloromethane. At 0 °C, bromine (0.850 g, 5.3 mmol) solution was added dropwise to the resin. And the mixture was stirred for 15 min, followed by dropping a solution of N-(3-hydroxymethylphenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylamine (0.930 g, 3 mmol). After 1 h, the resin was filtered and washed with CHCl<sub>3</sub>. Evaporation afforded the title compound. <sup>1</sup>H NMR: , 8.81 (s, 1H), 8.68 (d, 1H, J=5.4Hz), 8.47 (d, 1H, J=8.3Hz), 7.91 (m, 1H), 7.26-7.39 (m, 1H), 5.16 (s, 2H), 4.40 (s, 2H), 3.12 (q, 2H, J=7.3Hz), 1.40 (t, 3H, J=7.3Hz). MS calcd. 370.0; MS (M+1) 371.1.

Example 566


15 N-(3-(2-Methoxybenzyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine



-155-

The flask was charged with N-(3-bromomethylphenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylamine (0.180 g, 0.500 mmol), 2-methoxyphenylboronic acid (0.113 g, 0.75 mmol), 4.5 ml of DME, 1.5 ml of 2 M Na<sub>2</sub>CO<sub>3</sub>, and tetrakis(triphenylphosphine) palladium(0) (0.018 g, 0.016 mmol). After N<sub>2</sub>-flushing, the mixture was heated at 50 °C overnight. The reaction mixture was concentrated, diluted with water, and extracted with EtOAc. The organic layer was washed with brine and concentrated. The residue was chromatographed on a silica-gel column with 1:1 to 1:3 toluene/EtOAc to afford the title compound. <sup>1</sup>H NMR: , 8.37 (d, 1H, J=3.9Hz), 8.24 (s, 1H), 7.55 (d, 1H, J=7.8Hz), 6.74-7.17 (m, 9H), 4.76 (s, 2H), 3.80 (s, 2H), 3.69 (s, 2H), 2.94 (q, 2H, J=7.3Hz), 1.32 (t, 3H, J=7.3Hz). MS calcd. 396.2; MS (M+1) 397.2.

By a method similar to the method of Example 566 the following compounds were prepared:



No.	R <sub>2</sub>	Data
567	3-(2-trifluoromethyl benzyl)	<sup>1</sup> H NMR: 8.47 (d, 1H, J=4.9Hz), 8.31 (br s, 1H), 7.65 (m, 2H), 7.44 (m, 1H), 7.33 (m, 1H), 7.25 (d, 1H, J=7.8Hz), 7.21 (dd, 1H, J=7.8, 4.9Hz), 7.11 (br dd, 2H, J=7.8Hz), 7.06 (br d, 1H, J=7.3Hz), 7.01 (br d, 1H, J=7.8Hz), 6.95 (br s, 1H), 4.12 (s, 2H), 3.06 (q, 2H, J=7.3Hz), 1.40 (t, 3H, J=7.3Hz). MS calcd 434.1; MS (M+1) 435.2
568	3-(4-cyanobenzyl)	<sup>1</sup> H NMR: 8.48 (d, 1H, J=3.9Hz), 8.31 (s, 1H), 7.66 (d, 1H, J=7.8Hz), 7.55 (d, 2H, J=7.8Hz), 7.27 (m, 2H), 7.20 (m, 1H), 7.13 (m, 2H), 7.04 (d, 1H, J=7.3Hz), 6.97 (s, 1H), 4.85 (s, 2H), 3.95 (s, 2H), 3.08 (q, 2H, J=7.3Hz), 1.41 (t, 3H, J=7.3Hz). MS calcd 391.1; MS (M+1) 392.3
569	3-(2-chlorobenzyl)	<sup>1</sup> H NMR: 8.48 (br s, 1H), 8.33 (br s, 1H), 7.66 (br d, 1H, J=7.8Hz), 7.36 (m, 1H), 7.28-7.17 (m, 4H), 7.09 (br d, 2H, J=7.8Hz), 7.02 (m, 2H), 4.04 (d, 2H), 3.07



-156-

		(q,2H,J=7.3Hz), 1.40 (m,3H,J=7.3Hz). MS calcd 400.1; MS (M+1) 401.0
570	3-(4-fluorobenzyl)	<sup>1</sup> H NMR: 8.56 (d,1H,J=4.4Hz), 8.42 (s,1H), 7.95 (t,1H,J=7.4Hz), 7.72 (d,1H,J=7.6Hz), 7.27 (m,1H), 7.21 (t,1H,J=7.6Hz), 7.05 (m, 3H), 6.94 (m,3H), 4.86 (s,2H), 3.84 (s,2H), 3.06 (q,2H,J=7.3Hz), 1.40 (t,3H,J=7.3Hz). MS calcd 384.1; MS (M+1) 385.2.
571	3-(thien-2-yl methyl)	<sup>1</sup> H NMR: 8.47 (d,1H,J=4.9Hz), 8.34 (d,1H,J=1.5Hz), 7.66 (d,1H,J=8.3Hz), 7.25 (t,1H,J=7.3Hz), 7.20 (dd,1H,J=7.8,4.9Hz), 7.17-6.17 (m,4H), 6.91 (dd,1H,J=4.9,3.4Hz), 6.69 (d,1H,J=2.5Hz), 4.86 (s,2H), 4.10 (s,2H), 3.08 (q,2H,J=7.3Hz), 1.43 (t,3H,J=7.3Hz). MS calcd 372.1; MS (M+1) 373.1
572	3-(thien-3-yl methyl)	<sup>1</sup> H NMR: 8.48 (d,1H,J=3.9Hz), 8.33 (s,1H), 7.67 (d,1H,J=7.8Hz), 7.26-7.20 (m,3H), 7.10 (d,2H,J=7.3Hz), 7.02 (s,1H), 6.78 (m,2H), 3.91 (s,2H), 3.08 (q,2H,J=7.3Hz), 1.41 (t,3H,J=7.3Hz). MS calcd 372.1; MS (M+1) 373.1
573	3-(2-bromobenzyl)	<sup>1</sup> H NMR: 8.38 (d,1H,J=3.4Hz), 8.24 (d,1H,J=1.5Hz), 7.56 (d,1H,J=8.3Hz), 7.46 (d,1H,J=7.3Hz), 7.12 (m,3H), 7.01 (m,3H), 6.93 (m,2H), 4.75 (s,2H), 3.95 (s,2H), 2.98 (q,2H,J=7.3Hz), 1.30 (t,3H,J=7.3Hz). MS calcd 446.1; MS (M+1) 447.1
574	3-(5-methyl-thiophen-2-yl)methyl	<sup>1</sup> H NMR: 8.47 (dd,1H,J=4.9,1.6Hz), 8.35 (d,1H,J=1.6Hz), 7.66 (dt,1H,J=7.8,2.0Hz), 7.26-7.18 (m,2H), 7.15-7.07 (m,3H), 6.54 (m,1H), 6.46 (d,1H,J=3.1Hz), 4.84 (s,2H), 4.00 (s,2H), 3.07 (q,2H,J=7.3Hz), 2.41 (s,3H), 1.41 (t,3H,J=7.3Hz). MS calcd 386.1; MS (M+1) 387.1
575	3-(2,4-difluorobenzyl)	<sup>1</sup> H NMR: 8.8.49 (dd,1H,J=4.7,1.1Hz), 8.34 (d,1H,J=1.5Hz), 7.67 (d,1H,J=7.8Hz), 7.22 (m,2H), 7.08 (m,2H), 7.01 (s,1H), 6.94 (m,1H), 6.78 (m,2H), 4.84 (s,2H), 3.85 (s,2H), 3.08 (q,2H,J=7.3Hz), 1.41 (t,3H,J=7.3Hz). MS calcd 402.1; MS (M+1) 403.1
576	3-(2,4-dichloro benzyl)	

### Preparation 13

#### N-(4-(4-Cyanophenyl)phenyl)-N-ethylsulfonylamine

4-Amino-4'-cyanobiphenyl (1.03g, 5.3mmol) (TCI) was dissolved in a mixture of 8ml dichloromethane / 2ml pyridine was treated with ethanesulfonyl chloride (0.61g, 5.3mmol) and stirred for 18h at room temperature. The solution was diluted with dichloromethane and washed twice with 1N HCl. The organic phase was separated, dried over sodium sulfate,

-157-

filtered, and concentrated *in vacuo* to afford the title compound.  $^1\text{H}$  NMR: 7.75 (d, 2H, J=8Hz), 7.67 (d, 2H, J=8Hz), 7.60 (d, 2H, J=8Hz), 7.37 (d, 2H, J=8Hz), 6.90 (s, 1H), 3.22 (q, 2H, J=8Hz), 1.45 (t, 3H, J=8Hz).

5

#### Preparation 14

##### N-(2-Methoxy-5-phenyl)phenyl)-N-ethanesulfonylamine

The title compound was prepared using a method similar to the method of Preparation 13 using 5-phenyl-O-anisidine.

10 MS calcd. 291.3; MS (M+ -1) 290.2.

#### Preparation 15

##### N-(1-(4-Bromophenyl)ethyl)pyrid-3-ylmethylaniline

A solution of 4-bromo- $\alpha$ -methylbenzylamine (1.5g, 7.7mmol), 3-pyridinecarboxaldehyde (0.8g, 7.5mmol), and sodium triacetoxyborohydride (2.25g, 10.5mmol) in 60ml of 1,2-dichloroethane was stirred at room temperature for 18h. The solution was diluted with dichloromethane and washed twice with a saturated sodium bicarbonate solution, once with water, and once with brine. The organic layer was separated, dried over sodium sulfate, filtered, and concentrated *in vacuo* to afford the title compound.  $^1\text{H}$  NMR: 8.70 (m, 2H), 7.85 (m, 1H), 7.67 (d, 2H, J=8Hz), 7.45 (d, 2H, J=8Hz), 7.43 (m, 1H), 3.95 (q, 1H, J=8Hz), 3.83 (q, ABX, 2H), 1.57 (d, 3H, J=8Hz). MS calcd. 291.2; MS (M+1) 292.1.

#### Preparation 16

##### N-(1-(4-Bromophenyl)ethyl)-N-ethylsulfonylamine


30 4-Bromo- $\alpha$ -methylbenzylamine (8.15g, 40.73mmol) and (4.15g, 41.00mmol) triethylamine dissolved in 150ml dichloromethane was treated slowly with ethanesulfonyl chloride (5.27g, 41.00mmol) and stirred for 15 minutes. The reaction mixture was extracted with 50ml 1N HCl / 15ml

-158-

brine. The organic layer was separated, dried sodium sulfate, filtered, and concentrated *in vacuo* to afford the title compound.  $^1\text{H}$  NMR: 7.52 (d, 2H,  $J=12\text{Hz}$ ), 7.25 (d, 2H,  $J=12\text{Hz}$ ), 4.77 (m, 1H), 4.63 (m, 1H), 2.77 (m, 2H), 1.56 (d, 3H,  $J=8\text{Hz}$ ), 1.27 (t, 3H,  $J=8\text{Hz}$ ). MS calcd. 292.2; MS ( $M+ -1$ ) 291.1.

Example 577

10 N-(1-(4-Bromophenyl)ethyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline



N-(1-(4-Bromophenyl)ethyl)-N-ethylsulfonylamine (1.25g, 4.28mmol), and (0.75g, 4.6mmol) of 3-picolyl chloride hydrochloride, and (5.1g, 37mmol) of potassium carbonate were dissolved in 40ml DMF. Additional 3-picolyl chloride hydrochloride (0.75g, 4.6mmol) was added after 30 minutes and the mixture stirred 20h at room temperature. The mixture was diluted with 120ml ethyl acetate and 60ml of water and transferred to a separatory funnel. The organic layer was separated and washed twice with water. The organic layer was again separated and dried over sodium sulfate, filtered, and concentrated *in vacuo* to afford a dark oil. The oil was purified by radial chromatography on a 6 micron silica plate eluting with 1 to 4% methanol / dichloromethane to afford after concentration of fractions the title compound. MS calcd. 383.2; MS ( $M+1$ ) 384.1.

-159-

Preparation 17N-(Ethylsulfonyl)pyrid-3-ylmethanamine

3-Aminomethyl pyridine (2.15g, 19.88mmol) was dissolved in 40ml dichloromethane. Ethanesulfonyl chloride (2.56g, 19.9mmol) was added to the mixture and stirred for 18h. To the resultant precipitate was added 40ml diethyl ether and decanted from the solid. The solid was partitioned between a mixture of 100ml dichloromethane / 20ml isopropyl alcohol and 20ml of a saturated sodium bicarbonate solution. The organic layer was separated and the aqueous phase extracted twice with the 80/20 mixture of dichloromethane / isopropyl alcohol. The organic layers were combined, dried over sodium sulfate, filtered, and concentrated *in vacuo* to afford the title compound. <sup>1</sup>H NMR: 8.67 (s,1H), 8.60 (d,1H,J=4Hz), 7.90 (m,1H), 7.42 (m,1H), 5.77 (t,1H,J=6Hz), 4.38 (d,2H,J=6Hz), 3.07 (t,2H,J=8Hz), 1.38 (t,3H,J=8Hz). MS calcd. 200.2; MS (M+1) 201.1.

Preparation 184-Cyanophenylboronic acid.

A solution of 4-bromobenzonitrile (91 g, 0.50 mole) in THF (1.1 L) was dried in the presence of activated 3 Å molecular sieves at room temp. This solution was filtered and cooled to -100°C. A 1.6 M solution of *n*-butyllithium in hexanes (355 mL, 0.567 mol) was added to the cold solution over 15 min while maintaining the internal temperature between -105 and -93°C. To the resulting orange reaction mixture was added trimethylborate (81 g, 0.78 mol) over 3 min, briefly increasing the reaction temperature to -72°C. The reaction mixture was recooled to -100°C over 5 minutes and then was allowed to warm slowly to room temperature over 2.3 h. The reaction mixture was acidified with 4N HCl to pH 2.2, and was diluted with CH<sub>2</sub>Cl<sub>2</sub> (200 mL). The aqueous layer was separated and the organic layer was washed with brine (2

-160-

x 200 mL), dried over anhydrous  $\text{MgSO}_4$ , filtered, and the solvent removed under reduced pressure to a pale yellow solid. This solid was additionally purified by dissolution in 1N NaOH and extraction into  $\text{CH}_2\text{Cl}_2$  / THF (1:1, 2 x 200 mL). The aqueous phase was acidified with 4N HCl to pH 2.2 and was extracted into  $\text{CH}_2\text{Cl}_2$  / THF (1:1, 500 mL). The combined organic extracts were concentrated to a crude solid (64.6 g) that was triturated with diethyl ether (160 mL) and dried under vacuum to afford the intermediate title compound (44.0 g, 59.9%) as a white powder.  $^1\text{H}$  NMR ( $\text{d}_6$ -acetone, 300 MHz):  $\delta$  8.03 (d, 2H,  $J = 8.1$ ), 7.75 (d, 2H,  $J = 8.4$ ), 7.54 (s, 2H).

#### Preparation 19

##### 15 N-(3-(4-Cyanophenyl)phenyl)pyrid-3-ylmethylanine

A solution of N-(3-bromophenyl)pyridin-3-ylmethylanine (364mg, 1.38mmol), 4-cyanophenylboronic acid (305mg, 2.08mmol), sodium carbonate (2.15ml of a 2M aqueous solution, 4.3mmol), and tetrakis(triphenylphosphine)palladium(0) (70mg, 0.061mmol), under a nitrogen atmosphere was refluxed in dioxane (15ml) for 6h, then cooled to room temperature. The mixture was diluted with dichloromethane (70ml) and water (15ml). The organics were separated, dried over sodium sulfate, filtered, and concentrated to a brown solid. The solid was purified by radial chromatography eluting with 2 to 4% methanol / dichloromethane to afforded the title compound. MS calcd. 285.3; MS (M+1) 286.3.

#### Preparation 20

##### 30 N-(3-(4-(N'-Methanesulfonyl-N'-t-butoxycarbonylaminoethyl)phenyl)pyrid-3-ylmethylanine

To a 10 °C solution of phenethylamine (12.1 g, 0.100 mol) and triethylamine (11.1 g, 0.110 mol) in  $\text{CH}_2\text{Cl}_2$  (50 mL)

-161-

was added methanesulfonyl chloride (12.6 g, 0.110 mol) dropwise over 10 min. The solution was stirred at room temperature for 1.5 h and was then washed with 1 N HCl (5 x 20 mL). The organic phase was directly concentrated to provide the intermediate title compound, N-(methanesulfonyl)(2-phenylethyl)amine, (21.2 g, 93.3%) as an oil. <sup>1</sup>H NMR (300 MHz) 7.32 (m, 2H), 7.23 (m, 3H), 4.30 (br s, 1H), 3.40 (t, 2H, J=3.9), 2.88 (t, 2H, J=4.2), 2.81 (s, 3H).

To a stirring room temperature solution of N-(methanesulfonyl)-N-(2-phenylethyl)amine (205 g, 1.03 moles), water (200 mL), 95% sulfuric acid (111 g, 1.08 moles) in acetic acid (1 L), was added iodine (111 g, 0.438 mol) and periodic acid (45.6 g, 0.206 mol). The reaction mixture was warmed to 70-75 °C for 3 h. The heat was removed and the dark violet reaction mixture was allowed to proceed overnight at room temperature. Potassium hydroxide pellets (85%, 143 g, 2.16 moles) were added to neutralized the sulfuric acid and then enough saturated aqueous sodium sulfite was added to decolorize the mixture to afford a white suspension. The suspension was cooled to 15 °C and filtered. The filter cake was triturated thoroughly with water and was then dissolved in CH<sub>2</sub>Cl<sub>2</sub> (1 L) and extracted with additional water (2 x 200 mL). The organic phase was concentrated under reduced pressure to provide N-(2-(4-iodophenyl)ethyl)-N-(methanesulfonyl)amine, (201 g, 60.2%) as a white powder. <sup>1</sup>H NMR (300 MHz) 7.64 (d, 2H, J=4.8), 6.97 (d, 2H, J=5.1), 4.37 (br t, 1H, J=4), 3.36 (app. q, 2H, J=3.9), 2.85 (s, 3H), 2.82 (t, 2H, J=3.9).

A room temperature solution of N-(2-(4-iodophenyl)ethyl)-N-(methanesulfonyl)amine (201 g, 0.618 mol), 4-dimethylaminopyridine (3.8 g, 0.031 mol) and di-*t*-butyl dicarbonate (162 g, 0.744 mol) in DCM (1 L) was allowed to stir overnight. The reaction mixture was washed

-162-

with water (2 x 400 mL) and the organic phase was concentrated to about 600 mL and hexanes (400 mL) was added. This combined solution was washed again with water (400 mL) and was concentrated to a solid that was suspended in hexanes (600 mL) and filtered. The collected solids were dried under reduced pressure to afford the intermediate title compound, N-(t-butoxycarbonyl)-N-(2-(4-iodophenyl)ethyl)-N-(methylsulfonyl)amine (241.5 g, 91.5%) as a white solid. <sup>1</sup>H NMR (300 MHz) 7.63 (d, 2H, J=7.8), 6.98 (d, 2H, J=7.8), 3.88 (t, 2H, J=6.9), 3.10 (s, 3H), 2.88 (t, 2H, J=6.9), 1.51 (s, 9H).

To a degassed solution of N-(t-butoxycarbonyl)-N-(2-(4-iodophenyl)ethyl)-N-(methylsulfonyl)amine (128 g, 0.300 mol), triethylamine (91.1 g, 0.900 mol), and 1,1'-bis(diphenylphosphino)ferrocenedichloropalladium (II)-CH<sub>2</sub>Cl<sub>2</sub> complex (2.9 g, 0.0035 mol) in acetonitrile (600 mL) was added pinacolborane (50 g, 0.391 mol) dropwise. The mixture was stirred at 70-74 °C for 8 h and then was cooled to room temperature. The reaction mixture was concentrated to a fluid oil that was partitioned between MTBE (500 mL) and water (500 mL). The organic phase was separated and washed with water (2 x 200 mL) and concentrated to a residue that was partially dissolved with heptane (1 L). The heptane soluble fraction was filtered through Celite<sup>®</sup> 521 and concentrated to an oil (95 g). The residue was dissolved in acetone (600 mL) and heptane (600 mL) and filtered through Celite<sup>®</sup> 521. The combined filtrates were concentrated to 95 g of a mixture of a 3:1 molar ratio (<sup>1</sup>H NMR, 81.0% by weight) of intermediate title compound, N-(t-butoxycarbonyl)-N-(methylsulfonyl)-N-(2-(4-(4,4,5,5-tetramethyl(1,3,2-dioxaborolan-2-yl))phenyl)ethyl)amine and its protio derivative. <sup>1</sup>H NMR (300 MHz) 7.75 (d, 2H, J=7.8), 7.23 (d, 2H, J=8.1), 3.87 (t, 2H, J=8.1), 2.99 (s, 3H), 2.90 (t, 2H, J=7.5), 1.53 (s, 9H), 1.33 (s, 6H), 1.27 (s, 6H).

-163-

To a room temperature solution of N-(t-butoxycarbonyl)-N-(methylsulfonyl)-N-(2-(4-(4,4,5,5-tetramethyl(1,3,2-dioxaborolan-2-yl))phenyl)ethyl)amine (81.0% potent, 95 g, 0.18 mol) in acetone (2 L) was added 1N ammonium acetate (1L) and sodium periodate (145 g, 0.678 mol) with stirring. The reaction was allowed to proceed overnight. The reaction mixture was concentrated to remove the acetone, and the aqueous phase was decanted away from the oily product. The aqueous phase was extracted with DCM (100 mL) and MTBE (2 x 100 mL). The combined oily product and organic phases were adjusted to pH 12.5 with the addition of 1 N NaOH. The phases were separated, and the organic phase was extracted with 1 N NaOH (100 mL) and water (2 x 100 mL). HPLC analysis (60% CH<sub>3</sub>CN / 40% H<sub>2</sub>O, 2 mL / min, Zorbax C-18, 205 nm) of the organic phase indicated that the product had been removed from this phase. The aqueous phases (containing product) were finally combined and washed with CH<sub>2</sub>Cl<sub>2</sub> (100 mL) and MTBE (2 x 100 mL). The aqueous phase was added to CH<sub>2</sub>Cl<sub>2</sub> (450 mL) and 1 N H<sub>2</sub>SO<sub>4</sub> was added until the aqueous phase was at pH 3.05. The phases were separated and the aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub> (100 mL). The combined organic extracts (containing product) were concentrated to an oil (58.5 g) that crystallized overnight. The resulting solid mass was triturated with 10% MTBE in heptane (100 mL) to afford, after filtration and drying under reduced pressure, N-(t-butoxycarbonyl)-N-(methylsulfonyl)-N-(2-(4-boronic acid)phenyl)ethyl)amine (47.7 g, 77.2%) as a white powder. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO, 300 MHz) δ 7.83 (d, 2H, J= 4.8), 7.24 (d, 2H, J=5.1), 7.12 (s, 2H), 3.90 (t, 2H, J=3.9), 3.12 (s, 3H), 2.95 (t, 2H, J=4.5), 1.52 (s, 9H).

A solution of N-(3-bromophenyl)pyridin-3-ylmethylamine (180mg, 0.68mmol), N-(t-butoxycarbonyl)-N-(methylsulfonyl)-N-(2-(4-boronic acid)phenyl)ethyl)amine (350mg, 1.0mmol),



-164-

and dioxane (12ml) stirring under nitrogen was treated with 2M sodium carbonate (1.36ml, 2.7mmol) and tetrakis(triphenylphosphine) palladium(0) (35mg, 0.030mmol). This was heated to reflux for four hours, after which the reaction was concentrated *in vacuo*. The concentrated reaction mixture was then purified on a Chromatotron using a 4000 micron plate and 3% methanol/dichloromethane as the solvent system. The product containing fractions were collected and concentrated *in vacuo* to yield an oil, which was dissolved into ether/hexanes and concentrated *in vacuo* to give the title compound as an off-white foam. <sup>1</sup>H NMR: 8.60-8.70 (s, 1H), 8.50-8.60 (s, 1H), 7.75-7.80 (d, 1H, J=12Hz), 7.60-7.70 (m, 1H), 7.45-7.55 (m, 2H), 7.20-7.35 (m, 3H), 7.00-7.10 (m, 1H), 6.75-6.80 (d, 2H, 10Hz), 4.40-4.45 (s, 1H), 3.85-4.00 (m, 2H), 3.40-3.50 (q, 2H, 16Hz), 2.90-3.00 (m, 3H), 1.50-1.60 (s, 9H), 1.20-1.30 (m, 2H). MS calcd. 481.6; MS (M+1) 482.1.

#### Preparation 21

#### N-Methyl-4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide

4-Carboxyphenylboronic acid (10.0 g, 60.3 mmol) was suspended in 100 mL of toluene and pinacol added (7.1 g, 60.3 mmol). The reaction was heated to reflux with a Dean-Stark trap attached. The reaction became homogenous upon heating. After 2 h reflux the reaction was allowed to stand overnight and white needles were collected by filtration, rinsed with toluene, and vacuum dried to 4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzoic acid 12.2 g (81%). MS (FD MS): M = 248. Analysis calculated for C<sub>13</sub>H<sub>17</sub>BO<sub>4</sub>: C, 62.94; H, 6.91 Found: C, 62.71; H, 6.91.

4-(4,4,5,5-Tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzoic acid (6.0 g, 24.2 mmol) was suspended in 20 mL of thionyl chloride, 3 drops of DMF added and the reaction

-165-


heated to reflux under nitrogen. The solid dissolved upon warming. After refluxing 2 h the reaction was concentrated *in vacuo* and dried to a white solid under vacuum. The acid chloride was dissolved in 30 mL of THF and cooled to 0°C.

5 40% aqueous methylamine (10.4 mL) was added dropwise. After the addition was complete the ice bath was removed and stirring continued for 30 min. The reaction was concentrated *in vacuo* (3 x 1:1 EtOAc:toluene) then extracted with ether / water. The organic layer was dried over sodium  
10 sulfate, filtered and concentrated to 5.2 g (83%) of the title compound as a white solid. MS (FD MS): M = 261. Analysis calculated for C<sub>14</sub>H<sub>20</sub>BN<sub>3</sub>O<sub>3</sub>: C, 64.40; H, 7.72; N, 5.36. Found: C, 64.09; H, 7.70; N, 5.28.

15

Example 578

N-(4-(4-Cyanophenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine




A solution of N-(4-(4-cyanophenyl)phenyl)-N-  
20 ethylsulfonylamine (200mg, 0.7mmol), 3-picolyl chloride hydrochloride (123mg, 0.75mmol), and cesium carbonate (2.3g, 7.0mmol) was dissolved in 10ml DMF and stirred for 18h at room temperature. The mixture was diluted with 80ml ethyl acetate, 20ml water. The organic phase was separated and  
25 washed twice with 25ml of water. The organic phase was then dried over sodium sulfate, filtered, and concentrated *in vacuo* to an oily residue. The residue was flushed through a

-166-

20g mega bond elut silica column with 0 to 5% methanol / ethyl acetate. The desired fractions were concentrated *in vacuo* to afford the title compound.  $^1\text{H}$  NMR: 8.55 (bs, 1H), 8.41 (bs, 1H), 7.78 (d, 1H,  $J=8\text{Hz}$ ), 7.72 (d, 2H,  $J=12\text{Hz}$ ), 7.6 (d, 2H,  $J=12\text{Hz}$ ), 7.52 (d, 2H,  $J=12\text{Hz}$ ), 7.35 (d, 2H,  $J=12\text{Hz}$ ), 7.29 (m, 1H), 4.95 (s, 1H), 3.15 (q, 2H,  $J=8\text{Hz}$ ), 1.45 (t, 3H,  $J=8\text{Hz}$ ). MS calcd. 377.5; MS (M+1) 378.2.

Example 579

10 N-((2-Methoxy-5-phenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline




The title compound was prepared using a method similar to the method of Example 545 using N-(2-methoxy-5-phenyl)phenyl)-N-ethanesulfonylamine.  $^1\text{H}$  NMR: 8.5 (d, 1H,  $J=4\text{Hz}$ ), 8.41 (s, 1H), 7.88 (m, 1H), 7.51 (dd, 1H,  $J=4, 8\text{Hz}$ ), 7.28-7.45 (m, 7H), 7.0 (d, 1H,  $J=8\text{Hz}$ ), 4.9 (s, 2H), 3.98 (s, 1H), 3.16 (q, 2H,  $J=8\text{Hz}$ ), 1.45 (t, 3H,  $J=8\text{Hz}$ ). MS calcd. 382.5; MS (M+1) 383.1.

20

Example 580

N-(1-(4-Bromophenyl)ethyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline




-167-

A solution of N-(1-(4-bromophenyl)ethyl)pyrid-3-ylmethylaniline (630mg, 2.16mmol) and 4-(dimethylamino)pyridine (263mg, 2.16mmol) in 18ml of dichloromethane was treated with 2,2,2-trifluoroethylsulfonyl chloride (420mg, 2.3mmol) with stirring for 15 minutes at room temperature. The solution was concentrated to approximately 2ml and directly purified by radial chromatography on a 1 micron silica plate, eluting with 3% methanol / dichloromethane. The fractions were concentrated to afford the title compound. <sup>1</sup>H NMR: 8.52 (bs, 1H), 8.42 (bs, 1H), (d, 1H, J=8Hz), 7.45 (d, 2H, J=8Hz), 7.22 (d, 2H, J=8Hz), 5.25 (q, 1H, J=8Hz), 4.38 (q, 2H, ABX), 3.6 (q, 2H, J=12Hz), 1.68 (d, 3H, J=8Hz). MS calcd. 437.3; MS (M+1), (M+2), 438.0, 439.0.

#### Example 581

N-(3-(4-Cyanophenyl)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline




A solution of N-(3-(4-cyanophenyl)phenyl)pyrid-3-ylmethylaniline (255mg, 0.894mmol) in 8ml dichloromethane / 2ml pyridine was treated with 2,2,2-trifluoroethylsulfonyl chloride in 3 portions of 171mg (0.94mmol) over 1h at 45°C. The solution was cooled and diluted with 60ml diethyl ether and washed successively with water and saturated sodium bicarbonate solution. The organics were separated, dried over sodium sulfate, filtered, and concentrated to an oil. The oil was purified by radial chromatography on a 2 micron silica plate eluting with 2 to 4% methanol / dichloromethane

-168-

to afford the title compound.  $^1\text{H}$  NMR: 8.65 (bs, 1H), 8.42 (bs, 1H), 7.85 (m, 1H), 7.75 (d, 2H,  $J=8\text{Hz}$ ), 7.58 (d, 2H,  $J=8\text{Hz}$ ), 7.30-7.55 (m, 5H), 5.02 (s, 2H), 3.78 (q, 2H,  $J=8\text{Hz}$ ). MS calcd. 431.44; MS (M+1) 432.1.

5


Example 582N-((3-Iodophenyl)methyl)-N-(ethanesulfonyl)pyrid-3-ylamine

A solution of 3-aminopyridine (2.15g, 22.84mmol) and  
10 ethanesulfonyl chloride (2.18ml, 22.84mmol) in  
dichloromethane (150ml) was stirred for 20h at room  
temperature. The mixture was washed with 30ml saturated  
sodium bicarbonate solution and extracted with 15%  
isopropanol / dichloromethane. The organics were combined,  
15 dried over sodium sulfate, filtered, and concentrated *in*  
*vacuo* to give N-(ethanesulfonyl)-3-aminopyridine (2.15g, as  
a crude solid. The solid was dissolved in N,N-  
dimethylformamide (50ml) and treated with potassium  
carbonate (6.37g, 22.8mmol) and 3-iodobenzyl bromide (6.80g,  
20 22.8mmol) with stirring at room temperature for 3h. The  
mixture was diluted with ethyl acetate (150ml) and washed  
four times with water (100ml). The organics were separated,  
dried over sodium sulfate, filtered, and concentrated *in*  
*vacuo* to afford a crude residue. The residue was purified  
25 on a 200g pad of silica eluting with 3% methanol /  
dichloromethane to afford the title compound. MS calcd.  
402.3; MS (M+1) 403.1.

-169-

Example 583

N-((3-Iodophenyl)methyl)-N-(ethylsulfonyl)pyrid-3-ylmethylamine




5

A solution of N-(ethylsulfonyl)pyrid-3-ylmethylamine (1.1g, 5.5mmol), cesium carbonate (1.79g, 5.5mmol), and 3-iodobenzyl bromide (1.64g, 5.5mmol) was stirred in N,N-dimethylformamide (25ml) for 5h at room temperature. The mixture was diluted with ethyl acetate (100ml) and washed four times with water (50ml). The organics were separated, dried over sodium sulfate, filtered, and concentrated in vacuo to a crude residue. The residue was purified by radial chromatography on a 4 micron silica plate eluting with 3% methanol / dichloromethane to afford the title compound. <sup>1</sup>H NMR: 8.6 (d, 1H, J=4Hz), 8.41 (s, 1H), 7.75 (m, 1H), 7.65 (m, 1H), 7.60 (s, 1H), 7.25-7.40 (m, 2H), 7.08 (m, 1H), 4.43 (s, 2H), 4.33 (s, 2H), 3.08 (q, 2H, J=8Hz), 1.43 (t, 3H, J=8Hz). MS calcd. 416.3; MS (M+1) 417.1.

20

Example 584

N-(1-(4-(2,4-Difluorophenyl)phenyl)ethyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine




25

A solution of N-(1-(4-bromophenyl)ethyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine (200mg, 0.52mmol),

-170-

2,4-difluorophenylboronic acid (124mg, 0.78mmol), sodium carbonate (0.78ml of a 2M aqueous solution, 1.56mmol), and tetrakis(triphenylphosphine)palladium(0) (26mg, 0.023mmol), under a nitrogen atmosphere was refluxed in dioxane (8ml) for 6h, then cooled to room temperature. The mixture was concentrated to a tan residue and resuspended in 2ml dichloromethane. The suspension was subjected to purification by radial chromatography on a 2 micron silica plate eluting with 2 to 4% methanol / dichloromethane to afford the title compound. <sup>1</sup>H NMR: 8.45 (bs,1H), 8.37 (bs,1H), 7.62 (d,1H,J=8Hz), 7.45 (m,5H), 7.15 (m,1H), 6.95 (m,2H), 5.30 (q,1H,J=8Hz), 4.35 (s,2H), 2.92 (q,2H,J=8Hz), 1.67 (d,3H,J=8Hz), 1.37 (t,3H,J=8Hz). MS calcd. 416.5; MS (M+1) 417.1.

By a method similar to the method of Example 584 the following compounds were prepared:




No.	Y	R <sub>2</sub>	Data
585	CH(CH <sub>3</sub> )	4-(4-cyano phenyl)	<sup>1</sup> H NMR: 8.42 (bs,1H), 8.35 (bs,1H), 7.45-7.80 (m,9H), 7.21 (m,1H), 5.33 (q,1H,J=8Hz), 4.37 (s,2H), 2.95 (q,2H,J=8Hz), 1.70 (d,3H,J=8Hz), 1.40 (t,3H,J=8Hz). MS calcd. 405.5; MS (M+1) 406.2
586	CH <sub>2</sub>	3-(2,4-difluoro phenyl)	<sup>1</sup> H NMR : 8.57 (bs,1H), 8.45 (bs,1H), 7.58 (d,1H,J=8Hz), 7.15-7.35 (m,6H), 6.75-6.85 (m,2H), 4.87 (s,2H), 3.07 (q,2H,J=8Hz), 1.36 (t,3H,J=8Hz). MS calcd. 388.4; MS (M+1) 389.2
587	CH <sub>2</sub>	3-(4-cyano phenyl)	<sup>1</sup> H NMR: 8.53 (bs,1H), 8.42 (bs,1H), 7.78 (m,1H), 7.75 (d,2H,J=8Hz), 7.63 (d,2H,J=8Hz), 7.3-7.55 (m,5H), 4.49 (s,3H), 4.47 (s,3H), 3.12 (q,2H,J=8Hz), 1.45 (t,3H,J=8Hz). MS calcd. 391.5; MS (M+1) 392.2
588	CH <sub>2</sub>	3-(3-methane sulfonyl aminophenyl)	<sup>1</sup> H NMR: 8.65 (bs,1H), 8.58 (bs,1H), 7.9 (d,1H,J=8Hz), 7.70 (m,1H), 7.30-7.5 (m,9H), 4.54 (s,2H), 4.48 (s,2H), 3.13 (q,2H,J=8Hz), 3.09 (s,3H), 1.45 (t,3H,J=8Hz). MS calcd. 459.6; MS

- 171 -

			(M+1) 460.2
589	CH <sub>2</sub>	3-(4-N-methylamido phenyl)	<sup>1</sup> H NMR: 8.48 (bs,2H), 7.88 (d,2H,J=8Hz), 7.78 (m,1H), 7.60 (d,2H,J=8Hz), 7.54 (m,1H), 7.25-7.45 (m,4H), 6.35 (m,1H), 4.48 (s,2H), 3.10 (q,2H,J=8Hz), 3.07 (d,3H,J=4Hz), 1.45 (t,3H,J=8Hz). MS calcd. 423.5; MS (M+1) 423.2

By a method similar to the method of Example 584 the following compounds were prepared:



No.	R <sub>2</sub>	Data
590	3-(4-cyano phenyl)	<sup>1</sup> H NMR: 8.45 (bs,2H), 7.90 (d,1H,J=8Hz), 7.58 (d,2H,J=8Hz), 7.42 (d,2H,J=8Hz), 7.28-7.4 (m,5H), 4.90 (s,2H), 2.95 (q,2H,J=8Hz), 1.32 (t,3H,J=8Hz). MS calcd. 377.5; MS (M+1) 378.2
591	3-(2,4-difluoro phenyl)	<sup>1</sup> H NMR: 8.65 (bs,1H), 8.47 (bs,1H), 7.95 (m,1H), 7.25-7.45 (m,6H), 6.85-7.0 (m,2H), 5.0 (s,2H), 3.15 (q,2H,J=8Hz), 1.47 (t,3H,J=8Hz). MS calcd. 388.4; MS (M+1) 389.1
592	3-(3-methane sulfonyl aminophenyl)	<sup>1</sup> H NMR: 8.63 (s,1H), 8.58 (bs,1H), 7.9 (m,1H), 7.56 (s,1H), 7.25-7.5 (m,9H), 5.05 (s,2H), 3.3 (q,2H,J=8Hz), 3.08 (s,3H), 1.45 (t,3H,J=8Hz). MS calcd. 445.6; MS (M+1) 446.2
593	3-((3,5-bis-trifluoromethyl) phenyl)	<sup>1</sup> H NMR: 8.60 (bs,1H), 8.45 (bs,1H), 7.90 (m,3H), 7.35-7.55 (m,5H), 5.05 (s,2H), 3.20 (q,2H,J=8Hz), 1.52 (t,3H,J=8Hz). MS calcd. 488.5; MS (M+1) 489.1
594	3-phenyl	<sup>1</sup> H NMR: 8.49 (bs,1H), 8.41 (bs,1H), 7.90 (m,1H), 7.80 (m,1H), 7.15-7.45 (m,9H), 4.90 (s,2H), 3.07 (q,2H,J=8Hz), 1.38 (t,3H,J=8Hz). MS calcd. 352.5; MS (M+1) 353.2
595	3-((4-(2-trifluoroacetyl amino) ethyl)phenyl)	<sup>1</sup> H NMR: 8.65 (bs,1H), 8.57 (bs,1H), 7.92 (m,1H), 7.40-7.55 (m,4H), 7.20-7.30 (m,5H), 6.35 (m,1H), 5.15 (s,2H), 3.7 (m,2H), 3.20 (q,2H,J=8Hz), 2.95 (m,2H), 1.50 (t,3H,J=8Hz). MS calcd. 491.5; MS (M+1) 492.1
596	3-(4-acetylaminophenyl)	<sup>1</sup> H NMR: 8.55 (bs,1H), 8.50 (bs,1H), 7.95 (m,1H), 7.60 (m,3H), 7.35-7.55 (m,6H), 7.22 (m,1H), 5.03 (s,2H), 3.17 (q,2H,J=8Hz), 2.25 (s,3H), 1.45




		(t,3H,J=8Hz). MS calcd. 409.5; MS (M+1) 410.1
597	3-(4-methanesulfonylphenyl)	<sup>1</sup> H NMR: 8.57 (bs,1H), 8.43 (bs,1H), 8.03 (d,2H,J=16Hz), 7.85 (m,1H), 7.70 (d,2H,J=16Hz), 7.45-7.55 (m,3H), 7.35 (m,2H), 5.0 (s,2H), 3.18 (q,2H,J=8Hz), 3.12 (s,3H), 1.50 (t,3H,J=8Hz). MS calcd. 430.6; MS (M+1) 431.1
598	3-(4-trifluoromethoxyphenyl)	<sup>1</sup> H NMR: 8.57 (bs,1H), 8.45 (bs,1H), 7.85 (m,1H), 7.40-7.55 (m,5H), 7.35 (m,1H), 7.25-7.30 (m,3H), 5.0 (s,2H), 3.15 (q,2H,J=8Hz), 1.50 (t,3H,J=8Hz). MS calcd. 436.5; MS (M+1) 437.0
599	3-(pyrid-4-yl)	<sup>1</sup> H NMR: 8.68 (bs,2H), 8.55 (bs,1H), 8.45 (bs,1H), 7.82 (m,1H), 7.40-7.60 (m,6H), 7.32 (m,1H), 5.0 (s,2H), 3.17 (q,2H,J=8Hz), 1.50 (t,3H,J=8Hz). MS calcd. 353.4; MS (M+1) 354.2
600	3-(3-trifluoromethylphenyl)	<sup>1</sup> H NMR: 8.55 (bs,1H), 8.47 (bs,1H), 7.92 (m,1H), 7.45-7.70 (m,7H), 7.40 (m,1H), 7.32 (m,1H), 5.03 (s,2H), 3.17 (q,2H,J=8Hz), 1.50 (t,3H,J=8Hz). MS calcd. 420.5; MS (M+1) 421.1
601	3-(4-carboxyphenyl)	<sup>1</sup> H NMR: 8.70 (bs,1H), 8.62 (bs,1H), 8.15 (m,3H), 7.20-7.70 (m,7H), 5.12 (s,2H), 3.20 (q,2H,J=8Hz), 1.52 (t,3H,J=8Hz). MS calcd. 396.5; MS (M+1) 397.1
602	3-(4-(N'-methanesulfonyl-N'-t-butoxycarbonyl aminoethyl)phenyl)	MS calcd. 573.7; MS (M+1) 574.2
603	3-(2-methylphenyl)	<sup>1</sup> H NMR (Methanol d <sub>4</sub> ) : 8.40-8.60 (br s, 2H), 8.00-8.10 (m, 1H), 7.50-7.65 (br s, 1H), 7.35-7.50 (br s, 2H), 7.15-7.30 (m, 5H), 7.05-7.15 (br s, 1H), 5.00-5.15 (s, 2H), 3.20-3.35 (m, 2H), 2.05-2.20 (s, 3H), 1.40-1.50 (m, 3H). MS calcd. 480.51; MS (M+1) 367.2 (isolated as the trifluoroacetic acid salt)
604	3-(2-methoxyphenyl)	<sup>1</sup> H NMR (Methanol d <sub>4</sub> ): 8.40-8.50 (br s, 2H), 7.90-8.00 (d, 1H, J=10Hz), 7.50-7.55 (s, 1H), 7.40-7.50 (m, 1H), 7.25-7.40 (m, 4H), 7.20-7.25 (d, 1H, J=16Hz), 6.95-7.10 (m, 2H), 5.00-5.05 (s, 2H), 3.70-3.80 (s, 3H), 3.20-3.30 (m, 2H), 1.40-1.50 (m, 3H). MS calcd. 496.5; MS (M+1) 383.2 (isolated as the trifluoroacetic acid salt)
605	3-(2-trifluoromethylphenyl)	<sup>1</sup> H NMR (Methanol d <sub>4</sub> ) : 8.50-8.60 (br s, 2H), 8.05-8.15 (d, 1H, J=8Hz), 7.70-7.80 (d, 1H, J=16Hz), 7.50-7.70 (m, 3H), 7.40-7.50 (m, 2H), 7.30-7.35 (m, 2H), 7.20-7.30 (m, 1H), 5.05-5.15 (s, 2H), 3.20-3.35 (m, 2H), 1.35-1.45 (m, 3H). MS calcd. 534.5; MS (M+1) 421.2 (isolated as the trifluoroacetic acid salt)

-173-

606	3-(2-chlorophenyl)	<sup>1</sup> H NMR: 8.45-8.55 (s, 1H), 8.35-8.45 (s, 1H), 7.70-7.80 (d, 1H, J=10Hz), 7.40-7.50 (m, 1H), 7.35-7.40 (m, 2H), 7.25-7.35 (m, 3H), 7.20-7.35 (m, 3H), 4.90-4.95 (s, 2H), 3.10-3.20 (m, 2H), 1.40-1.50 (t, 3H, J=8Hz). MS calcd. 386.9; MS (M+1) 387.2
607	3-(2-fluorophenyl)	<sup>1</sup> H NMR (Methanol d <sub>4</sub> ) : 8.40-8.50 (br s, 2H), 7.85-7.95 (d, 1H, J=10Hz), 7.50-7.55 (s, 1H), 7.30-7.50 (m, 6H), 7.15-7.30 (m, 2H), 5.00-5.10 (s, 2H), 3.20-3.35 (m, 2H), 1.40-1.50 (t, 3H, J=8Hz). MS calcd. 484.5; MS (M+1) 371.1 (isolated as the trifluoroacetic acid salt)
608	3-(3,5-difluorophenyl)	<sup>1</sup> H NMR (Methanol d <sub>4</sub> ) : 8.55-8.40 (br s, 2H), 7.90-8.00 (d, 1H, J=10Hz), 7.50-7.60 (m, 2H), 7.40-7.50 (m, 3H), 7.15-7.25 (d, 2H, J=16Hz), 6.90-7.00 (m, 1H), 5.00-5.10 (s, 2H), 3.20-3.35 (m, 2H), 1.40-1.50 (m, 3H). MS calcd. 502.4; MS (M+1) 389.2 (isolated as the trifluoroacetic acid salt)
609	3-(3-acetylaminophenyl)	<sup>1</sup> H NMR: 8.45-8.50 (d, 1H, J=8Hz), 8.40-8.45 (s, 1H), 7.70-7.75 (d, 1H, J=16Hz), 7.50-7.60 (m, 2H), 7.45-7.50 (d, 1H, J=16Hz), 7.40-7.45 (s, 1H), 7.35-7.40 (m, 2H), 7.20-7.30 (m, 3H), 4.90-4.95 (s, 2H), 3.45-3.50 (q, 1H, 12Hz), 3.10-3.20 (q, 2H, J=16Hz), 2.20-2.25 (s, 3H), 1.40-1.50 (t, 3H, J=8Hz). MS calcd. 446.0; MS (M+1) 410.2 (isolated as the hydrochloric acid salt)

Example 610

N-((4-Methyl-3-(4-cyanophenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine



5

A solution of N-(3-iodo-4-methylphenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylamine (103mg, 0.247mmol) and 4-cyanophenylboronic acid (55mg, 0.371mmol) in 4ml of 1,4-dioxane stirring under nitrogen was treated with 2M sodium


-174-

carbonate (0.37mL, 0.741mmol) and  
tetrakis(triphenylphosphine)palladium (0) (12.6mg,  
0.011mmol). This was refluxed for three hours and then  
allowed to stir at room temperature for 16 hours. The  
5 reaction had not completed after this time so the reaction  
was treated with an additional 5.7mg(0.005mmol) of  
tetrakis(triphenylphosphine)palladium (0) and refluxed for  
two hours. The reaction was then cooled to room temperature  
and concentrated *in vacuo*. This was then dissolved in a  
10 minimal amount of dichloromethane and purified on a  
Chromatotron (2000micron plate) with 3%  
methanol/dichloromethane to give a residue which was  
recrystallized using ether/hexanes to give the title  
compound. <sup>1</sup>H NMR: 8.45-8.55 (d, 1H, J=4Hz), 8.35-8.40 (s,  
15 1H), 7.75-7.80 (d, 1H, J=12Hz), 7.65-7.70 (d, 1H, J=12Hz),  
7.20-7.35 (m, 5H), 7.15-7.20 (m, 1H), 7.00-7.05 (s, 1H),  
4.85-4.95 (s, 2H), 3.05-3.15 (q, 2H, J=10Hz), 2.15-2.20 (s,  
3H), 1.40-1.50 (t, 3H, J=8Hz) ; MS calcd. 391.5; MS (M+1) =  
392.1.

20

Example 611

N-(3-(4-(2-Methanesulfonylaminoethyl)phenyl)phenyl)-N-  
(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylamine



25

A solution of N-(3-(4-(N'-methanesulfonyl-N'-t-butoxycarbonylaminoethyl)phenyl)pyrid-3-ylmethylamine and  
2,2,2-trifluoroethanesulfonyl chloride (134mg, 0.735mmol) in  
3:1 dichloromethane/pyridine (4mL) was stirred for 16 hours  
after which the reaction had not completed yet. Another


-175-

67mg (0.368mmol) of 2,2,2-trifluoroethanesulfonyl chloride was added and the reaction was heated to 45°C for 30 minutes. The reaction was then allowed to cool to room temperature. Ether was used to extract the organic material, which was washed four times with deionized water. The organic layer was then dried with sodium carbonate, filtered, and concentrated *in vacuo*. The concentrated mixture was then dissolved in a minimal amount of dichloromethane and purified on a Chromatotron (6000micron plate) using 3% methanol/dichloromethane. The proper fractions were collected and concentrated *in vacuo* and treated with 3mL of 2:1 trifluoroacetic acid:dichloromethane and stirred for 1 hour to take off the Boc protecting group. Acetonitrile was then added and the reaction was concentrated *in vacuo* in order to remove any water. This step was repeated two additional times, which left an oil. The oil was treated with 0.64mL (0.64mmol) of 1M hydrochloric acid, 5mL of ether, and 5mL hexanes, which caused the title compound to precipitate out of solution. The solid was filtered and collected. This yielded the title compound as the HCl salt. <sup>1</sup>H NMR (Methanol d<sub>4</sub>) : 8.35-8.45 (br s, 2H), 7.80-7.90 (d, 1H, J=12Hz), 7.55-7.60 (d, 1H, J=16Hz), 7.40-7.55 (m, 4H), 7.30-7.40 (m, 4H), 5.00-5.05 (s, 2H), 4.30-4.40 (q, 4H, J=16Hz), 3.25-3.35 (s, 4H), 2.80-2.90 (m, 4H). MS calcd. 564.0; MS (M+1) = 528.3.

-176-

Example 612

N-((3-(4-Amidophenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine



5


N-((3-(4-Cyanophenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine (147mg, 0.36mmol) was treated with concentrated sulfuric acid (2.5ml) for 22h at room temperature. The reaction was neutralized with 50 weight % aqueous sodium hydroxide to pH 7.0. This milky solution was extracted three times with 15% isopropanol / dichloromethane and the organics were combined and concentrated in vacuo to a tan solid. The solid was purified by radial chromatography on a 1 micron silica plate eluting with 3 to 10% methanol / dichloromethane to afford the title compound (49mg, 35%). <sup>1</sup>H NMR: 8.56 (s, 1H), 8.5 (s, 1H), 7.95 (m, 1H), 7.90 (d, 2H, J=8Hz), 7.72 (d, 2H, J=8Hz), 7.3-7.58 (m, 5H), 6.20 (bs, 1H), 5.75 (bs, 1H), 5.05 (s, 2H), 3.17 (q, 2H, J=8Hz), 1.5 (t, 3H, J=8Hz). MS calcd. 395.5; MS (M+1) 396.1.

20

Example 613

N-(3-(4-(2-Methanesulfonylaminoethyl)phenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine

-177-




A solution of N-(3-(4-(N'-(methanesulfonyl)-N'-t-butoxycarbonylaminoethyl)phenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine (78mg, 0.136mmol) in  
 5 2ml dichloromethane / 3ml trifluoroacetic acid was stirred for 40 minutes at room temperature. The mixture was diluted with acetonitrile and concentrated to an oil. The oil was dissolved in 0.5ml dichloromethane and treated with ethereal HCl to afford the title compound as the HCl salt. <sup>1</sup>H NMR (d<sub>4</sub>  
 10 methanol): 8.80 (bs,1H), 8.73 (bs,1H), 8.60 (m,1H), 8.0 (m,1H), 7.66 (m,1H), 7.57 (m,1H), 7.52 (d,2H,J=8Hz), 7.43 (m,2H), 7.35 (d,2H,J=8Hz), 5.23 (s,2H), 3.32 (m,4H), 2.87 (t,2H,J=8Hz), 2.82 (s,3H), 1.40 (t,3H,J=8Hz). MS calcd. 510.1; MS (M+1-HCl) 474.1.

15

Example 614

N-((3-(2-Methyl-4-cyanophenyl)phenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine



20 In a one pot reaction, a solution of 4-bromo-3-methylbenzonitrile (166mg, 0.845mmol), bis(pinacolato)diboron (236mg, 0.93mmol), potassium acetate (249mg, 2.5mmol), and DMSO (7mL) was stirred under nitrogen until all the materials present were dissolved. Then 55mg

-178-

(0.068mmol) palladium(II) chloride (dppf) complex with dichloromethane (1:1) was introduced to the solution, which was then heated to 80°C for three hours. During this time the solution turned a dark brown color.

5       The reaction was allowed to cool to room temperature after the three hours had passed N-(3-bromophenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline (300mg, 0.845mmol), 2M sodium carbonate (1.25mL, 2.5mmol) and another equivalent of palladium(II) chloride (dppf) complex with dichloromethane  
10   (1:1) (55mg, 0.068mmol) were added to the reaction. The reaction was then stirred under nitrogen while being heated to 120°C for 16 hours.

After 16 hours, the reaction was allowed to cool again. Dichloromethane was used for the extraction and the crude  
15   was washed with water and brine. After the washing, the organic layer was dried with sodium sulfate, filtered, and concentrated *in vacuo*. This was then purified using a Chromatotron (4000 micron plate) and 2% methanol/dichloromethane. The fractions containing the  
20   product were collected and concentrated *in vacuo* and recrystallized from ether/hexanes to give the title compound. <sup>1</sup>H NMR: 8.45-8.55 (d, 1H, J=8Hz), 8.30-8.40 (s, 1H), 7.70-7.80 (d, 1H, J=16Hz), 7.45-7.55 (m, 1H), 7.40-7.50 (t, 2H, J=16Hz), 7.25-7.35 (m, 3H), 7.15-7.20 (m, 1H), 7.10-  
25   7.15 (s, 1H), 4.90-5.00 (s, 2H), 3.05-3.10 (q, 2H, J=10Hz), 2.05-2.10 (s, 3H), 1.4-1.5 (t, 3H, J=16Hz). MS calcd. 391.5; MS (M+1) 392.2.

#### Preparation 22

30   N-(4-(2,4-Difluorophenoxy)phenyl)-N-ethylsulfonylamine

Ethanesulfonyl chloride (9.4 mL, 99 mmol) was added dropwise over 5 min to a dark gray solution of 4-(2,4-difluorophenoxy)aniline (20.0 g, 90 mmol) and 4-dimethylaminopyridine (16.6 g, 136 mmol) in methylene

-179-

chloride (180 mL) at room temperature under nitrogen, after which the mixture was stirred at room temperature for 26 h. The mixture was washed with saturated aqueous sodium bicarbonate solution (200 mL), brine (100 mL), dried over  
5 MgSO<sub>4</sub>, filtered and the solvent was removed under reduced pressure to afford the crude product as an amber oil. This residue was triturated with diethyl ether (300 mL) and filtered to provide an amber solid. This material was further purified by flash column chromatography on silica  
10 gel, eluting with ethyl acetate/chloroform (1:9), to provide the title compound as an off-white powder: <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) 9.69 (s, 1H), 7.52-7.44 (m, 1H), 7.28-7.09 (m, 2H), 7.20 (d, J = 8.8 Hz, 2H), 6.95 (d, J = 8.8 Hz, 2H), 3.03 (q, J = 7.6 Hz, 2H), 2.50 (t, J = 1.6 Hz, 3H).

#### Preparation 23

N-(4-(2,4-Difluorophenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)amine

2,2,2-Trifluoroethanesulfonyl chloride (10.0 g, 54.8  
20 mmol) was added dropwise to a solution of 4-(2,4-difluorophenoxy)aniline (11.0 g, 50.0 mmol) and 4-dimethylaminopyridine (9.1 g, 74.6 mmol) in methylene chloride (100 mL) at 0 °C under nitrogen. The mixture was slowly warmed to room temperature, stirring for a total of  
25 72 h. The mixture was washed with saturated aqueous sodium bicarbonate solution (200 mL), dried over MgSO<sub>4</sub>, filtered and the solvent was removed under reduced pressure to afford the crude product as an orange oil. This residue was purified by flash column chromatography on silica gel,  
30 eluting with ethyl acetate/chloroform (1:4), to provide the title compound as a yellow solid: mp 105-108 °C; TLC R<sub>f</sub> (1:9 ethyl acetate/chloroform) = 0.75; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) 7.23 (d, J = 8.9 Hz, 2H), 7.14-6.87 (m, 3H), 6.93



-180-

(d,  $J = 8.9$  Hz, 2H), 3.76 (q,  $J = 8.8$  Hz, 2H) ppm; APCI MS  $m/z$  366  $[C_{14}H_{10}F_5NO_3S-H]^-$ .

#### Preparation 24

##### 5 (2-Fluoropyridin-3-yl)methanol

Thionyl chloride (0.43 mL, 5.87 mmol) was added to a slurry of 2-fluoronicotinic acid (9, 0.69 g, 4.89 mmol) and N,N-dimethylformamide (0.1 mL, catalytic amount) in benzene (60 mL) at room temperature under nitrogen, after which the  
10 mixture was heated at reflux for 3 h. The solvent was removed under reduced pressure to provide an amber oil that was dissolved in 1,4-dioxane (40 mL) under nitrogen. The solution was treated with sodium borohydride (0.38 g, 10.0 mmol) and the mixture was stirred at room temperature for 24  
15 h. The suspension was diluted with saturated  $NaHCO_3$  solution (200 mL) and extracted with diethyl ether (3 x 200 mL). The combined organic extracts were dried over  $Na_2SO_4$ , filtered and the solvents were removed under reduced pressure to provide the title compound:  $^1H$  NMR (300 MHz)  
20 8.13 (d,  $J = 3.7$  Hz, 1H), 7.93-7.88 (m, 1H), 7.24-7.19 (m, 1H), 4.78 (s, 2H), 2.14 (bs, 1H) ppm.

#### Preparation 25

##### (2-Methylpyridin-3-yl)methanol

25 To a slurry of lithium aluminum hydride (500 mg, 13.2 mmol) in anhydrous tetrahydrofuran (45 mL) at 0 °C under nitrogen was added dropwise a solution of methyl 2-methylnicotinate (17, 1.0 g, 6.6 mmol) in anhydrous tetrahydrofuran (5 mL). After 1 h the suspension was  
30 diluted with water (1 mL) and 6 N NaOH solution (5 mL). The precipitate was removed by vacuum filtration and the filtrate was dried over  $Na_2SO_4$ , filtered, and the solvent was removed under reduced pressure to provide the title compound as a light yellow oil:  $^1H$  NMR (300 MHz) 8.37 (d,

-181-

J = 4.9 Hz, 1H), 7.72 (d, J = 7.6 Hz, 1H), 7.14 (dd, J = 7.6 Hz, 4.9 Hz, 1H), 4.72 (s, 2H), 2.95-2.74 (bs, 1H), 2.53 (s, 3H) ppm.

5

Preparation 26(6-Methylpyridin-3-yl)methanol

To a slurry of lithium aluminum hydride (500 mg, 13.2 mmol) in anhydrous tetrahydrofuran (45 mL) at 0 °C under nitrogen was added dropwise a solution of methyl 4-methylnicotinate (15, 1.0 g, 6.6 mmol) in anhydrous tetrahydrofuran (5 mL). After 1.5 h the suspension was diluted with water (1 mL) and 6 N NaOH solution (5 mL). The precipitate was removed by vacuum filtration and the filtrate was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and the solvent was removed under reduced pressure to provide the title compound as a light yellow oil: <sup>1</sup>H NMR (300 MHz) 8.34 (s, 1H), 7.60 (d, J = 7.9 Hz, 1H), 7.12 (d, J = 7.9 Hz, 1H), 4.65 (s, 2H), 4.18-4.10 (bs, 1H), 2.51 (s, 3H) ppm.

20

Preparation 27(6-Fluoropyridin-3-yl)methanol

Thionyl chloride (0.29 mL, 4.08 mmol) was added to a slurry of 6-fluoronicotinic acid (13, 0.48 g, 3.40 mmol) and N,N-dimethylformamide (0.1 mL, catalytic amount) in benzene (40 mL) at room temperature under nitrogen, after which the mixture was heated at reflux for 3 h. The solvent was removed under reduced pressure to provide an amber oil that was dissolved in 1,4-dioxane (25 mL) under nitrogen. The solution was treated with sodium borohydride (0.26 g, 6.80 mmol) and the mixture was stirred at room temperature for 24 h. The suspension was diluted with sat. NaHCO<sub>3</sub> solution (200 mL) and extracted with diethyl ether (3 x 200 mL). The combined organic extracts were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and the solvents were removed under reduced pressure to


-182-

provide the title compound as a colorless oil:  $^1\text{H}$  NMR (300 MHz) 8.20 (s, 1H), 7.87-7.81 (dt,  $J = 8.0, 2.3$  Hz, 1H), 6.97-6.93 (dd,  $J = 8.4, 2.8$  Hz, 1H), 4.73 (s, 2H), 2.04 (bs, 1H) ppm.

5

Example 615

N-(4-(2,4-Difluorophenoxy)phenyl)-N-(ethanesulfonyl)-2-fluoropyrid-3-ylmethylaniline




10 Di-tert-butyl azodicarboxylate (0.36 g, 1.57 mmol) was added to a suspension of (2-fluoropyridin-3-yl)methanol (10, 0.20 g, 1.57 mmol), N-(4-(2,4-difluorophenoxy)phenyl)-N-ethylsulfonylamine (0.49 g, 1.56 mmol) and polystyrene resin-bound triphenylphosphine (0.52 g, 1.57 mmol, ~3  
15 mmol/g) in anhydrous tetrahydrofuran (5 mL) at 0 °C under nitrogen. The resulting yellow suspension was warmed to room temperature and stirred for 19 h, after which additional polystyrene resin bound triphenylphosphine (0.52 g, 1.57 mmol, ~3mmol/g) and di-tert-butyl azodicarboxylate  
20 (0.36 g, 1.57 mmol) were added. This solution was stirred for 24 h after which the suspension was diluted with anhydrous tetrahydrofuran (100 mL) and the solids were removed by vacuum filtration. The filtrate solvent was removed under reduced pressure to provide the crude product  
25 as a yellow oil. This oil was purified by medium pressure liquid chromatography on silica gel, eluting with hexanes/ethyl acetate (7:3), to provide a colorless oil. This oil was triturated with hexanes/ethyl acetate (4-5 mL) to give the title compound as a white powder (0.28 g, 53%):

-183-

mp 110-112 °C; TLC  $R_f$  (3:2 hexanes/ethyl acetate) = 0.44;  $^1\text{H}$  NMR (300 MHz) 8.11 (m, 1H), 7.90 (m, 1H), 7.24-7.15 (m, 3H), 7.11-7.03 (m, 1H), 6.98-6.86 (m, 2H), 6.84 (d,  $J$  = 6.8 Hz, 2H), 4.91 (s, 2H), 3.15-3.07 (q,  $J$  = 7.4 Hz, 2H), 1.43 (t,  $J$  = 7.4 Hz, 1H) ppm;  $^{13}\text{C}$  NMR (75 MHz) 163.3, 161.3, 160.1, 157.8, 156.5, 153.1, 147.5 (d,  $J$  = 14.9 Hz), 141.7, 139.1, 133.6, 130.5, 123.8 (d,  $J$  = 9.7 Hz), 122.1, 119.1 (d,  $J$  = 29.2 Hz), 117.5, 112.0 (d,  $J$  = 22.9 Hz), 106.0 (t,  $J$  = 24.3 Hz), 49.1, 46.0, 8.44 ppm; APCI MS  $m/z$  423  $[\text{C}_{20}\text{H}_{17}\text{F}_3\text{N}_2\text{O}_3\text{S}+\text{H}]^+$ .  
 5 = 7.4 Hz, 1H) ppm;  $^{13}\text{C}$  NMR (75 MHz) 163.3, 161.3, 160.1, 157.8, 156.5, 153.1, 147.5 (d,  $J$  = 14.9 Hz), 141.7, 139.1, 133.6, 130.5, 123.8 (d,  $J$  = 9.7 Hz), 122.1, 119.1 (d,  $J$  = 29.2 Hz), 117.5, 112.0 (d,  $J$  = 22.9 Hz), 106.0 (t,  $J$  = 24.3 Hz), 49.1, 46.0, 8.44 ppm; APCI MS  $m/z$  423  $[\text{C}_{20}\text{H}_{17}\text{F}_3\text{N}_2\text{O}_3\text{S}+\text{H}]^+$ .  
 10 Anal. Calcd. for  $\text{C}_{20}\text{H}_{17}\text{F}_3\text{N}_2\text{O}_3\text{S}$ : C, 56.87; H, 4.06; N, 6.63. Found: C, 56.90; H, 4.10; N, 6.45.

By a method similar to the method of Example 615 the following compounds were prepared:



15

No.	$R_1$	$R_6$	Data
616	$\text{SO}_2\text{CH}_2\text{CF}_3$	2-fluoro	mp 114-115 °C; TLC $R_f$ (7:3 hexanes/ethyl acetate) = 0.45; $^1\text{H}$ NMR (300 MHz) 8.13 (m, 1H), 7.90 (m, 1H), 7.22-7.11 (m, 3H), 7.10-7.05 (m, 1H), 6.99-6.90 (m, 2H), 6.87 (d, $J$ = 8.9 Hz, 2H), 4.92 (s, 2H), 3.86-3.77 (q, $J$ = 9.1 Hz, 2H) ppm; $^{13}\text{C}$ NMR (75 MHz) 163.3, 161.4, 160.1, 158.5, 158.1, 156.5, 153.2, 147.9 (d, $J$ = 14.9 Hz), 141.7, 138.9, 131.9, 130.8, 123.9 (d, $J$ = 9.7 Hz), 122.2, 118.3 (d, $J$ = 29.3 Hz), 117.7, 112.2 (d, $J$ = 22.9 Hz), 106.0 (t, $J$ = 21.9 Hz), 52.6 (q, $J$ = 31.9 Hz), 49.8 ppm; APCI MS $m/z$ 477 $[\text{C}_{20}\text{H}_{14}\text{F}_6\text{N}_2\text{O}_3\text{S}+\text{H}]^+$ . Anal. Calcd. for $\text{C}_{20}\text{H}_{14}\text{F}_6\text{N}_2\text{O}_3\text{S}$ : C, 50.42; H, 2.96; N, 5.88. Found: C, 50.05; H, 2.79; N, 5.72
617	$\text{SO}_2\text{CH}_2\text{CH}_3$	6-fluoro	mp 103-105 °C; TLC $R_f$ (1:1 hexanes/ethyl acetate) = 0.38; $^1\text{H}$ NMR (300 MHz) 7.92 (d, $J$ = 2.2 Hz, 1H), 7.86-7.80 (dt, $J$ = 8.0, 2.4 Hz, 1H), 7.13 (d, $J$ = 9.0 Hz, 2H), 7.10-7.04 (m, 1H), 6.98-6.87 (m, 2H), 6.84 (d, $J$ = 8.9 Hz, 2H), 4.81 (s, 2H), 3.12-3.05 (q, $J$ = 7.4 Hz, 2H), 1.43 (t, $J$ = 7.4 Hz, 1H) ppm; $^{13}\text{C}$ NMR (75 MHz)

			165.3, 162.1, 161.3 (d, $J = 10.2$ Hz), 158.0, 156.5 (d, $J = 12.5$ Hz), 153.2 (d, $J = 12.2$ Hz), 147.9 (d, $J = 15.0$ Hz), 142.2, 139.1 (d, $J = 12.1$ Hz), 133.3, 130.9, 130.3, 124.0 (d, $J = 9.8$ Hz), 117.5, 112.0 (d, $J = 22.9$ Hz), 110.1 (d, $J = 37.5$ Hz), 106.0 (t, $J = 22.0$ Hz), 52.6, 46.1, 8.50 ppm; APCI MS $m/z$ 423 [C <sub>20</sub> H <sub>17</sub> F <sub>3</sub> N <sub>2</sub> O <sub>3</sub> S+H] <sup>+</sup> . Anal. Calcd. for C <sub>20</sub> H <sub>17</sub> F <sub>3</sub> N <sub>2</sub> O <sub>3</sub> S: C, 56.87; H, 4.06; N, 6.63. Found: C, 56.78; H, 3.97; N, 6.53
618	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	6-fluoro	mp 107-109 °C; TLC $R_f$ (3:2 hexanes/ethyl acetate) = 0.54; <sup>1</sup> H NMR (300 MHz) 7.90 (d, $J = 2.3$ Hz, 1H), 7.84-7.78 (dt, $J = 2.4, 8.0$ Hz, 1H), 7.11 (d, $J = 9.0$ Hz, 2H), 7.15-7.06 (m, 2H), 7.00-6.90 (m, 2H), 6.87 (d, $J = 8.9$ Hz, 2H), 4.83 (s, 2H), 3.83-3.76 (q, $J = 9.1$ Hz, 2H) ppm; <sup>13</sup> C NMR (75 MHz) 165.5, 162.3, 161.4 (d, $J = 10.3$ Hz), 158.6, 156.6 (d, $J = 12.4$ Hz), 153.2 (d, $J = 12.2$ Hz), 148.1 (d, $J = 15.1$ Hz), 142.1 (d, $J = 8.2$ Hz), 138.8, 131.5, 131.0, 129.5, 124.1 (d, $J = 9.7$ Hz), 120.2, 117.7, 112.3 (d, $J = 23.0$ Hz), 110.3 (d, $J = 37.6$ Hz), 106.0 (t, $J = 21.9$ Hz), 53.4, 52.7 (q, $J = 31.7$ Hz) ppm; APCI MS $m/z$ 477 [C <sub>20</sub> H <sub>14</sub> F <sub>6</sub> N <sub>2</sub> O <sub>3</sub> S+H] <sup>+</sup> . Anal. Calcd. for C <sub>20</sub> H <sub>14</sub> F <sub>6</sub> N <sub>2</sub> O <sub>3</sub> S: C, 50.42; H, 2.96; N, 5.88. Found: C, 50.40; H, 2.87; N, 5.77
619	SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	6-methyl	mp 108-111 °C; TLC $R_f$ (9:1 methylene chloride/methanol) = 0.52; <sup>1</sup> H NMR (300 MHz) 8.20 (d, $J = 2.0$ Hz, 1H), 7.59 (dd, $J = 2.3$ Hz, 8.0 Hz, 1H), 7.13 (d, $J = 9.0$ Hz, 1H), 7.15-7.06 (m, 3H), 6.86-6.83 (m, 3H), 6.84 (d, $J = 8.9$ Hz, 1H), 4.79 (s, 2H), 3.11-3.04 (q, $J = 7.4$ Hz, 2H), 2.51 (s, 3H), 1.72 (s, H), 1.43 (t, $J = 7.4$ Hz, 1H) ppm; <sup>13</sup> C NMR (75 MHz) 161.2 (d, $J = 10.2$ Hz), 158.6, 158.0 (d, $J = 10.2$ Hz), 157.8, 156.5 (d, $J = 12.2$ Hz), 153.2 (d, $J = 12.2$ Hz), 149.5, 139.2 (d, $J = 11.9$ Hz), 137.2, 133.6, 131.0, 129.3, 123.9 (d, $J = 9.7$ Hz), 123.6, 117.5, 112.0 (d, $J = 22.9$ Hz), 106.0 (t, $J = 22.0$ Hz), 53.3, 46.1, 24.5, 8.5 ppm; APCI MS $m/z$ 419 [C <sub>21</sub> H <sub>20</sub> F <sub>2</sub> N <sub>2</sub> O <sub>3</sub> S+H] <sup>+</sup> . Anal. Calcd. for C <sub>21</sub> H <sub>20</sub> F <sub>2</sub> N <sub>2</sub> O <sub>3</sub> S: C, 60.27; H, 4.82; N, 6.69. Found: C, 59.94; H, 4.76; N, 6.58
620	SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	6-methyl	mp 94-96 °C; TLC $R_f$ (4:6 hexanes/ethyl acetate) = 0.58; <sup>1</sup> H NMR (300 MHz) 8.16 (d, $J = 2.1$ Hz, 1H), 7.59 (dd, $J = 2.3$ Hz, 7.9 Hz, 2H), 7.13 (d, $J = 9.0$ Hz, 1H), 7.09-7.06 (m, 2H), 6.99-6.86 (m, 3H), 6.86 (d, $J = 8.9$ Hz, 1H), 4.81 (s, 2H), 3.85-3.76 (q, $J = 9.1$ Hz, 2H), 2.52 (s, 3H), 1.61 (s, 2H) ppm; <sup>13</sup> C NMR (75 MHz) 161.4 (d, $J = 10.2$ Hz), 159.0, 158.4, 158.1 (d, $J = 10.2$ Hz), 157.8, 156.6 (d, $J = 12.2$ Hz), 153.2 (d, $J = 12.2$ Hz), 149.5, 139.0 (d, $J = 12.0$ Hz), 137.1, 131.9, 131.1, 128.4, 124.0 (d, $J = 9.8$ Hz), 123.7, 120.3,

-185-

			117.6, 112.1 (d, $J = 22.9$ Hz), 106.0 (t, $J = 22.0$ Hz), 54.1, 53.4-52.1 (q, $J = 31.7$ Hz), 24.6 ppm; APCI MS $m/z$ 473 $[C_{21}H_{17}F_5N_2O_3S+H]^+$ . Anal. Calcd. for $C_{21}H_{17}F_5N_2O_3S$ : C, 53.39; H, 3.63; N, 5.93. Found: C, 53.34; H, 3.44; N, 5.82
621	$SO_2CH_2CH_3$	2-methyl	TLC $R_f$ (9:1 methylene chloride/methanol) = 0.47; $^1H$ NMR (300 MHz) 8.37 (d, $J = 4.9$ Hz, 1H), 7.55 (d, $J = 7.7$ Hz, 1H), 7.17 (d, $J = 9.2$ Hz, 2H), 7.10-7.02 (m, 3H), 6.98-6.88 (m, 3H), 6.83 (d, $J = 8.9$ Hz, 2H), 4.88 (s, 2H), 3.14-3.07 (q, $J = 7.4$ Hz, 2H), 2.46 (s, 3H), 1.45 (t, $J = 7.3$ Hz, 1H) ppm; $^{13}C$ NMR (75 MHz) 161.4, 157.9, 157.3, 156.8, 154.1, 148.5, 139.4, 137.5, 134.2, 131.0, 130.0, 124.3, 122.2, 117.5, 112.3 (d, $J = 22.9$ Hz), 106.3 (t, $J = 22.0$ Hz), 52.5, 45.6, 28.8, 23.7, 8.7 ppm; APCI MS $m/z$ 419 $[C_{21}H_{20}F_2N_2O_3S+H]^+$ . Anal. Calcd. for $C_{21}H_{20}F_2N_2O_3S \cdot 0.10H_2O$ : C, 60.02; H, 4.84; N, 6.67. Found: C, 59.86; H, 4.91; N, 6.74
622	$SO_2CH_2CF_3$	2-methyl	TLC $R_f$ (7:3 ethyl acetate/hexanes) = 0.46; $^1H$ NMR (300 MHz) 8.38 (d, $J = 4.8$ Hz, 1H), 7.54 (d, $J = 7.7$ Hz, 1H), 7.15 (d, $J = 9.0$ Hz, 1H), 7.12-7.04 (m, 3H), 6.99-6.88 (m, 3H), 6.85 (d, $J = 9.0$ Hz, 1H), 4.90 (s, 2H), 3.85-3.76 (q, $J = 9.1$ Hz, 2H), 2.41 (s, 3H) ppm; $^{13}C$ NMR (75 MHz) 161.4 (d, $J = 10.3$ Hz), 158.5, 158.0 (d, $J = 9.9$ Hz), 157.4, 156.6 (d, $J = 10.2$ Hz), 153.2 (d, $J = 10.2$ Hz), 149.2, 138.9 (d, $J = 10.3$ Hz), 137.8, 132.0, 130.9, 129.3, 123.9 (t, $J = 3.9$ Hz), 121.8, 120.3, 117.6, 112.1 (d, $J = 22.9$ Hz), 106.0 (t, $J = 22.0$ Hz), 53.5, 53.1-52.0 (q, $J = 31.8$ Hz), 22.3 ppm; APCI MS $m/z$ 473 $[C_{21}H_{17}F_5N_2O_3S+H]^+$ . Anal. Calcd. for $C_{21}H_{17}F_5N_2O_3S$ : C, 53.39; H, 3.63; N, 5.93. Found: C, 53.73; H, 3.76; N, 6.07

### Preparation 28

#### N-(Ethylsulfonyl)-N-(3-(N'-(2-chlorophenyl)-N'-methylsulfonyl)aniline

5 To a solution of 2-chloro-N-methylaniline (4.96 mmol, 1.1 eq., 702 mg) and pyridine (5.86 mmol, 464 mg) in 20 mL of methylene chloride was added 3-nitrobenzenesulfonyl chloride (4.51 mmol, 1.0 g) in one portion. The reaction mixture was stirred 16 hr and diluted with methylene

10 chloride and water. The aqueous layer was separated and extracted once with methylene chloride. The combined organic layers were extracted once with a saturated solution

-186-

of aqueous sodium carbonate, once with water, and then dried over sodium sulfate. The crude product was purified using flash chromatography (10-60% ethyl acetate / hexane) to give N-(2-chlorophenyl)-N-methyl-(3-nitrophenyl)sulfonamide which solidifies on standing. <sup>1</sup>H NMR (300 MHz) 3.29 (s, 3H), 7.25-7.45 (m, 4H), 7.71 (t, J = 7.82 Hz, 1H), 8.06 (d, J = 7.82 Hz, 1H), 8.40-8.50 (m, 1H), 8.60-8.65 (m, 1H). MS (ESI): m/e 327.13, 329.13 (M+1).

A solution of N-(2-chlorophenyl)-N-methyl-(3-

nitrophenyl)sulfonamide (2.6 mmol, 848 mg) and tin (II) chloride dihydrate (10.4 mmol, 2.35 g) were heated for 4 hr at reflux in 15 mL of methanol. The reaction mixture was cooled to room temperature and diluted with 20 mL of ethyl acetate and 10 mL of a saturated solution of aqueous sodium carbonate. Heavy gas evolution was observed upon addition of aqueous sodium carbonate to the reaction mixture. The reaction mixture was stirred 1 hr and then filtered through a plug of Celite®. The layers were separated and the organic layer was extracted once with a saturated solution of aqueous sodium carbonate, once with brine, and then dried with sodium sulfate. The crude product was purified using flash chromatography (0-5% ethyl acetate / methylene chloride) to give N-(2-chlorophenyl)-N-methyl-(3-aminophenyl)sulfonamide as an off white foam. <sup>1</sup>H NMR (300 MHz) 3.13 (s, 3H), 3.80 (br s, 2H), 6.80-6.87 (m, 1H), 7.02-7.25 (m, 6H), 7.30-7.38 (m, 1 H). MS (ESI): m/e 297.2, 299.2 (M+1).

To a solution of

N-(2-Chlorophenyl)-N-methyl-(3-aminophenyl)sulfonamide (3.38 mmol, 1.0 g), DMAP (0.83 mmol, 120 mg) and pyridine (16.9 mol, 1.33 mL) in 20 mL methylene chloride was added ethanesulfonyl chloride (12.2 mmol, 3.6 eq., 1.56 g) in two portions. The reaction mixture was stirred 10 hr and diluted with then methylene chloride and water. The layers


-187-

were separated and the aqueous layer was extracted once with methylene chloride. The combined organic layers were extracted once with a saturated solution of aqueous sodium carbonate and then dried with sodium sulfate. The crude product was purified using flash chromatography (0-30% ethyl acetate / methylene chloride) to give the title compound. <sup>1</sup>H NMR (300 MHz) 1.36 (t, J = 5.87 Hz, 3H), 3.10-3.16 (m, 2H), 3.23 (s, 3H), 7.04 (s, 1H), 7.22-7.60 (m, 8 H). MS (ESI): m/e 389.01, 390.88 (M+1).

10

Example 623

N-(3-(N'-Methyl-N'-(2-chlorophenylsulfonamido)phenyl))-N-(ethanesulfonyl)pyrid-3-ylmethylaniline



To a solution of (0.48 mmol, 186.6 mg) dissolved in 5 mL of anhydrous DMF was added a 1.0 M solution of lithium bis(trimethylsilyl)amide in hexane (1.2 mmol, 2.5 eq., 1.2 mL). The solution was stirred 5 min. and 3-picolyl chloride hydrochloride (0.50 mmol, 82 mg) was added in one portion. After overnight stirring, the reaction mixture was concentrated to dryness under vacuum. The residue was dissolved in methanol and loaded onto a 2 g SCX ion exchange cartridge (Varian Sample Preparation, 0.75 meq / g loading). After washing with additional methanol the crude product was eluted from the column using 10 mL of a 2.0 M solution of ammonia in methanol. The crude produce was purified using mass guided reverse phase semi-preparative chromatography (ESI detection; ACN with 0.1% TFA / water with 0.1% TFA;



-188-

Flow rate 25 mL/min.; 10% organic for 1 min. then a gradient over 12 min. to 95% organic then 95% organic for 1 min.) to give the TFA salt of the product as an oil after solvent evaporation. The purified sample was desalted using  
5 a 1 g SCX ion exchange column as previously described (methanol washing solvent and 2.0 M ammonia in methanol the eluting solvent) to give the title compound. <sup>1</sup>H NMR (300 MHz) 1.40 (t, J = 7.43 Hz, 3H), 3.04 (q, J = 7.43 Hz, 2H), 3.18 (s, 3H), 4.89 (s, 2H), 7.20-7.30 (m, 3H), 7.32-7.37 (m,  
10 1H), 7.44-7.52 (m, 3H), 7.65-7.75 (m, 3H), 8.39 (br s, 1H), 8.51 (br s, 1H). MS (ESI): m/e 480.2, 482.2 (M+1).

The compounds of the present invention can be  
15 administered alone or in the form of a pharmaceutical composition, that is, combined with pharmaceutically acceptable carriers or excipients, the proportion and nature of which are determined by the solubility and chemical properties of the compound selected, the chosen route of  
20 administration, and standard pharmaceutical practice. The compounds of the present invention, while effective themselves, may be formulated and administered in the form of their pharmaceutically acceptable salts, for purposes of stability, convenience of crystallization, increased  
25 solubility, and the like.

In practice, the compounds of formula I are usually administered in the form of pharmaceutical compositions, that is, in admixture with pharmaceutically acceptable carriers or diluents, the proportion and nature of which are  
30 determined by the chemical properties of the selected compound of formula I, the chosen route of administration, and standard pharmaceutical practice.

Thus, the present invention provides pharmaceutical compositions comprising a compound of the formula I and a  
35 pharmaceutically acceptable diluent.

-189-

The compounds of formula I can be administered by a variety of routes. In effecting treatment of a patient afflicted with disorders described above, a compound of formula I can be administered in any form or mode which  
5 makes the compound bioavailable in an effective amount, including oral and parenteral routes. For example, compounds of formula I can be administered orally, by inhalation, subcutaneously, intramuscularly, intravenously, transdermally, intranasally, rectally, occularly, topically,  
10 sublingually, buccally, and the like. Oral administration is generally preferred for treatment of the neurological and psychiatric disorders described herein.

One skilled in the art of preparing formulations can readily select the proper form and mode of administration  
15 depending upon the particular characteristics of the compound selected, the disorder or condition to be treated, the stage of the disorder or condition, and other relevant circumstances. (*Remington's Pharmaceutical Sciences*, 18th Edition, Mack Publishing Co. (1990)).

20 The pharmaceutical compositions of the present invention are prepared in a manner well known in the pharmaceutical art. The carrier or excipient may be a solid, semi-solid, or liquid material which can serve as a vehicle or medium for the active ingredient. Suitable carriers or  
25 excipients are well known in the art. The pharmaceutical composition may be adapted for oral, inhalation, parenteral, or topical use and may be administered to the patient in the form of tablets, capsules, aerosols, inhalants, suppositories, solution, suspensions, or the like.

30 The compounds of the present invention may be administered orally, for example, with an inert diluent or capsules or compressed into tablets. For the purpose of oral therapeutic administration, the compounds may be  
35 incorporated with excipients and used in the form of tablets, troches, capsules, elixirs, suspensions, syrups, wafers, chewing gums and the like. These preparations should contain at least 4% of the compound of the present

-190-

invention, the active ingredient, but may be varied depending upon the particular form and may conveniently be between 4% to about 70% of the weight of the unit. The amount of the compound present in compositions is such that  
5 a suitable dosage will be obtained. Preferred compositions and preparations according to the present invention may be determined by a person skilled in the art.

The tablets, pills, capsules, troches, and the like may also contain one or more of the following adjuvants: binders  
10 such as microcrystalline cellulose, gum tragacanth or gelatin; excipients such as starch or lactose, disintegrating agents such as alginic acid, Primogel, corn starch and the like; lubricants such as magnesium stearate or Sterotex; glidants such as colloidal silicon dioxide; and  
15 sweetening agents such as sucrose or saccharin may be added or a flavoring agent such as peppermint, methyl salicylate or orange flavoring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier such as polyethylene glycol or a fatty oil.  
20 Other dosage unit forms may contain other various materials which modify the physical form of the dosage unit, for example, as coatings. Thus, tablets or pills may be coated with sugar, shellac, or other coating agents. A syrup may contain, in addition to the present compounds, sucrose as a  
25 sweetening agent and certain preservatives, dyes and colorings and flavors. Materials used in preparing these various compositions should be pharmaceutically pure and non-toxic in the amounts used.

For the purpose of parenteral therapeutic  
30 administration, the compounds of the present invention may be incorporated into a solution or suspension. These preparations typically contain at least 0.1% of a compound of the invention, but may be varied to be between 0.1 and about 90% of the weight thereof. The amount of the compound  
35 of formula I present in such compositions is such that a suitable dosage will be obtained. The solutions or suspensions may also include one or more of the following

-191-

adjuvants: sterile diluents such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl paraben; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylene diaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic. Preferred compositions and preparations are able to be determined by one skilled in the art.

The compounds of the present invention may also be administered topically, and when done so the carrier may suitably comprise a solution, ointment, or gel base. The base, for example, may comprise one or more of the following: petrolatum, lanolin, polyethylene glycols, bees wax, mineral oil, diluents such as water and alcohol, and emulsifiers, and stabilizers. Topical formulations may contain a concentration of the formula I or its pharmaceutical salt from about 0.1 to about 10% w/v (weight per unit volume).

In order to more fully illustrate the operation of this invention, typical pharmaceutical compositions are described below. The examples are illustrative only, and are not intended to limit the scope of the invention in any way.

-192-

FORMULATION 1

A typical tablet, appropriate for use in this method, may be prepared using conventional techniques and may contain:

5		Amount per Tablet	Concentration by Weight (%)
	Compound of		
	formula I	5.0 mg	4.7
	Lactosum	67.8 mg Ph. Eur.	64.2
10	Avicel®	31.4 mg	29.8
	Amberlite®	1.0 mg	1.0
	magnesium stearate	0.25 mg Ph. Eur.	0.3
	TOTAL	105.45 mg	100

15 FORMULATION 2

Hard gelatin capsules may be prepared using the following ingredients:

		Amount per Tablet	Concentration by Weight (%)
20	Compound of		
	formula I	0.1 mg	0.05
	starch dried	200 mg	95.2
	magnesium stearate	10 mg	4.8
	TOTAL	210.1 mg	100

25 The above ingredients are mixed and filled into hard gelatin capsules in 210.1 mg quantities.

FORMULATION 3

30 Suspensions each containing 1 mg of medicament per 5 mL dose are as follows:

	Amount per 5mL of suspension
Compound of formula I	1 mg
sodium carboxymethyl cellulose	50 mg
35 syrup	1.25 mL
benzoic acid solution	0.10 mL
flavor	q.v.

-193-

color	q.v.
water	q.s. to 5 mL

The medicament is passed through a No. 45 mesh U.S. sieve and mixed with the sodium carboxymethyl cellulose and syrup to form a smooth paste. The benzoic acid solution, flavor and color is diluted with some of the water and added to the paste with stirring. Sufficient water is then added to produce the required volume.

The compounds of formula I are potentiators of metabotropic glutamate (mGlu) receptor function, in particular they are potentiators of mGlu<sub>2</sub> and/or mGlu<sub>3</sub> receptors. That is, the compounds of formula I do not appear to bind at the glutamate recognition site on the mGlu receptor, but in the presence of glutamate or a glutamate agonist, the compounds of formula I increase mGlu receptor response. The present potentiators are expected to have their effect at mGlu receptors by virtue of their ability to increase the response of such receptors to glutamate or glutamate agonists, enhancing the function of the receptors. The behavior of the potentiators of formula I at mGlu<sub>2</sub> receptors is shown in Example A which is suitable to identify potentiators useful for carrying out the present invention.

#### Example A

##### [<sup>3</sup>H]-2S-2-Amino-2-(1S,2S-2-carboxycyclopropan-1-yl)-3-(xanth-9-yl) propionic acid Receptor Binding

[<sup>3</sup>H]-2S-2-Amino-2-(1S,2S-2-carboxycyclopropan-1-yl)-3-(xanth-9-yl) propionic acid (17.5 Ci/mmol) was tritiated by Chemsyn Laboratories. Membranes from mGlu<sub>2</sub> receptor expressing RGT cells were prepared by harvesting adherent cells from confluent T-150 flasks with a cell scraper. Cells and media were then placed in 50 ml conical tubes and centrifuged at 4° C at 1000g for 5 minutes. The supernatant was removed and the pellet was frozen at -10° C until use. Washed cell membranes were prepared by adding 20 ml of 10 mM

-194-

potassium phosphate buffer pH 7.6 @ 5° C + 100 mM potassium bromide and homogenizing with an Ultra-Turrax tissuemizer for 15 seconds at 90% output. The homogenate was centrifuged at 48,000 x g for 10 minutes at 5°C. This last step was repeated two times, and the final pellet was resuspended in 10 ml of the same buffer and reserved on ice until initiating the binding assay.

[<sup>3</sup>H]-2S-2-Amino-2-(1S,2S-2-carboxycyclopropan-1-yl)-3-(xanth-9-yl) propionic acid can be assayed by the method described by Johnson et al. *Neuropharmacology* 38: 1519-1529 (1999). Accordingly, [<sup>3</sup>H]-2S-2-amino-2-(1S,2S-2-carboxycyclopropan-1-yl)-3-(xanth-9-yl) propionic acid binding was assayed in a reaction mixture containing 10 mM potassium phosphate (pH 7.6), 100 mM potassium bromide and generally 1 nM [<sup>3</sup>H]-2S-2-amino-2-(1S,2S-2-carboxycyclopropan-1-yl)-3-(xanth-9-yl) propionic acid, (final volume 500 µl). The incubation was initiated by the addition of the membrane suspension, (about 15 µg membrane protein), and allowed to continue on ice for 30 min.

Incubation was terminated by rapid filtration with a Brandel cell harvester through glass fiber Whatman GF/B filters pre-wet with the same potassium phosphate buffer assay buffer (at 4° C). The filters were washed five times with 1 ml of buffer. Filter sections were transferred to minivials and 5 ml of Fisher Scientific ScintiSafe liquid scintillation cocktail was added to each vial. Vials were allowed to set for several hours prior to counting on a Beckman LS6000 liquid scintillation counter.

Displacement of 1 nM [<sup>3</sup>H]-2S-2-amino-2-(1S,2S-2-carboxycyclopropan-1-yl)-3-(xanth-9-yl) propionic acid binding was assayed in homogenates derived from a mGlu<sub>2</sub> receptor expressing cell line. At concentrations up to 100 µM, the potentiator of formula I did not significantly displace [<sup>3</sup>H]-2S-2-amino-2-(1S,2S-2-carboxycyclopropan-1-yl)-3-(xanth-9-yl) propionic acid binding. Displacement of 1 nM [<sup>3</sup>H]-2S-2-Amino-2-(1S,2S-2-carboxycyclopropan-1-yl)-3-(xanth-9-yl)propionic acid is presented in the table below.

-195-

Treatment	K <sub>T</sub> (μM)	(95% Confidence Interval)
Glutamate	7.8	(1.1-15.7)
Compound of Example 16	>100	

Potentiation of Glutamate-Induced Increase in Intracellular Calcium with a mGluR<sub>2</sub> Expressing Cell Line.

5 Cell lines expressing human mGlu<sub>2</sub> receptors were derived as previously described (Desai, Burnett, Mayne, Schoepp, *Mol. Pharmacol.* 48, 648-657, 1995) and cultured in DMEM with 5% dialyzed fetal bovine serum, 1 mM glutamine, 1 mM sodium pyruvate, 50 μg/ml Geneticin G418, and 0.2 mg/ml  
10 hygromycin B. Confluent cultures were passaged weekly. These cells are referred to as RGT cells for Rat Glutamate Transporter, and have been co-transfected with the glutamate/aspartate transporter GLAST. The RGT cell line expressing the mGluR<sub>2</sub> receptors was stably transfected with  
15 the promiscuous G-protein, Galpha15 to change the signaling pathway to the mGlu<sub>2</sub> receptor to one that could be easily measured through release of intracellular calcium. Thus, intracellular calcium levels were monitored before and after the addition of drugs on a Fluorometric Imaging Plate Reader  
20 (i.e. FLIPR, Molecular Devices). The following buffer was used throughout as an assay buffer: 10 mM KCl, 138 mM NaCl, 5 mM Ca Cl<sub>2</sub>, 1 mM MgCl<sub>2</sub>, 4 mM NaH<sub>2</sub>PO<sub>4</sub>, 10 mM Glucose, 10 mM HEPES, pH 7.4. Cells that had been plated 48 h prior at a density of 30-40,000 cells per well in a 96-well plate were  
25 loaded with a calcium-sensitive dye for 90 min at 25°C. Fluo-3 (2 mM in DMSO, Molecular Probes) were mixed with a equal volume of 10% pluronic acid in DMSO, and diluted to 8 μM into the buffer described above containing 10% fetal bovine serum to make the loading buffer. Following loading  
30 of the cells, the loading buffer was removed and replaced with assay buffer prior to drug addition and monitoring on



-196-

the FLIPR. Results are analyzed by taking the difference of the peak height minus the background fluorescence and expressing the results as a percent of the maximal glutamate response (30  $\mu$ M glutamate, typically about 30-50,000

5 Relative Fluorescent Units).

Data from such an experiment using a representative compound of the present invention is presented in the table below.

Treatment	% Maximal Glutamate Response (as defined by 30 $\mu$ M Glutamate) mean $\pm$ SD
20 $\mu$ M Compound of Example 1	2.3 $\pm$ 0.8 %
0.5 $\mu$ M L-glutamate	1.9 $\pm$ 1.0 %
20 $\mu$ M Compound of Example 1 + 0.5 $\mu$ M L-glutamate	85.5 $\pm$ 5.8 %*

10 \* p < 0.05, ANOVA followed by contrast analysis

As seen from the results above the compounds of formula I potentiate the response of the mGlu<sub>2</sub> receptor to glutamate. This response can be blocked by pretreatment of  
15 the cells with the mGlu<sub>2/3</sub> antagonists 2S-2-amino-2-(1S,2S-2-carboxycyclopropan-1-yl)-3-(xanth-9-yl) propionic acid. Taken together with the inability of these compounds to displace [<sup>3</sup>H]-2S-2-amino-2-(1S,2S-2-carboxycyclopropan-1-yl)-3-(xanth-9-yl) propionic acid-labeled receptors, these  
20 results show that the compounds of formula I act at a site other than the glutamate recognition site to potentiate the effects of glutamate at mGlu receptors.

Using the procedure of Example A, the effect of representative compounds of the present invention on the  
25 response of different mGluR<sub>2</sub> glutamate-site agonists (such as glutamate, 2R,4R-4-aminopyrrolidine-2,4-dicarboxylic acid (APDC), and 2S,2'R,3'R)-2-(2',3'-dicarboxycyclopropyl)

-197-

glycine (DCG IV), (Schoepp, D.D., Jane, D.E., Monn, J.A., *Neuropharmacology* 38: 1431-1476, (1999)) is presented in the tables below.

5

	Buffer	3 $\mu$ M Compound of Example 22
	% Maximal Response (mean $\pm$ SD, n=2)	
Buffer	0.0 $\pm$ 3.2%	4.7 $\pm$ 0.4%
30 $\mu$ M APDC	100.7 $\pm$ 1.6%	120.3 $\pm$ 10.0%
10 $\mu$ M APDC	107.2 $\pm$ 3.6%	119.1 $\pm$ 4.0%
3 $\mu$ M APDC	80.3 $\pm$ 9.3%	114.0 $\pm$ 0.8%*
1 $\mu$ M APDC	16.8 $\pm$ 1.5%	92.0 $\pm$ 3.4%*
300 nM APDC	1.7 $\pm$ 0.4%	28.4 $\pm$ 1.7%*
100 nM APDC	-0.8 $\pm$ 0.5%	0.2 $\pm$ 0.6%
30 nM APDC	1.3 $\pm$ 0.1%	0.5 $\pm$ 1.5%
1 nM APDC	0.3 $\pm$ 1.2%	0.1 $\pm$ 0.7%

\* p < 0.05, t-test vs. agonists alone

	Buffer	3 $\mu$ M Compound of Example 22
	% Maximal Response (mean $\pm$ SD, n=2)	
Buffer	0.0 $\pm$ 0.7%	3.7 $\pm$ 1.4%
30 $\mu$ M DCG IV	68.5 $\pm$ 0.14%	138.2 $\pm$ 7.1%

-198-

10 $\mu$ M DCG IV	79.3 $\pm$ 10.8%	147.9 $\pm$ 1.5%
3 $\mu$ M DCG IV	68.3 $\pm$ 5.1%	153.0 $\pm$ 2.8%
1 $\mu$ M DCG IV	67.9 $\pm$ 15.0%	141.1 $\pm$ 5.9%
300 nM DCG IV	34.3 $\pm$ 6.3%	132.9 $\pm$ 5.5%
100 nM DCG IV	4.2 $\pm$ 0.8%	83.3 $\pm$ 10.5%
30 nM DCG IV	4.5 $\pm$ 4.3%	11.1 $\pm$ 6.1%
1 nM DCG IV	4.3 $\pm$ 3.5%	-0.1 $\pm$ 0.6%

	Buffer	1 $\mu$ M Compound of Example 16
	% Maximal Response (mean $\pm$ SD, n=2)	
Buffer	0.0 $\pm$ 0.3%	1.6 $\pm$ 1.0%
24 $\mu$ M Glutamate	100.0 $\pm$ 5.8%	95.4 $\pm$ 2.4%
8 $\mu$ M Glutamate	96.7 $\pm$ 3.2%	98.4 $\pm$ 6.4%
2.4 $\mu$ M Glutamate	83.5 $\pm$ 4.6%	92.5 $\pm$ 4.4%
800 nM Glutamate	16.9 $\pm$ 8.2%	62.9 $\pm$ 0.1%
240 nM Glutamate	-0.1 $\pm$ 0.4%	17.4 $\pm$ 2.8%
80 nM Glutamate	0.1 $\pm$ 0.1%	0.3 $\pm$ 1.1%
24 nM Glutamate	0.2 $\pm$ 0.2%	0.1 $\pm$ 0.8%
8 nM Glutamate	-0.7 $\pm$ 0.3%	2.8 $\pm$ 1.9%

-199-

	Buffer +	1 $\mu$ M Compound of Example 1	1 $\mu$ M Compound of Example 9
	% Maximal Response (mean $\pm$ SD, n=2)		
	0.0 $\pm$ 2.6%	0.1 $\pm$ 0.1%	4.2 $\pm$ 2.7%
30 $\mu$ M Glutamate	107.2 $\pm$ 11.7%	116.7 $\pm$ 4.3%	108.0 $\pm$ 12.3%
10 $\mu$ M Glutamate	108.2 $\pm$ 4.8%	107.6 $\pm$ 10.7%	102.9 $\pm$ 6.2%
3 $\mu$ M Glutamate	84.2 $\pm$ 4.3%	112.1 $\pm$ 10.3%	101.6 $\pm$ 8.1%
1 $\mu$ M Glutamate	22.6% (N=1)	79.2 $\pm$ 14.6%	93.1 $\pm$ 2.5%
300 nM Glutamate	-1.1 $\pm$ 0.6%	42.5 $\pm$ 4.3%	60.0 $\pm$ 10.6%
100 nM Glutamate	-1.0 $\pm$ 1.2%	1.5 $\pm$ 1.2%	20.5 $\pm$ 4.3%
30 nM Glutamate	-1.6 $\pm$ 0.4%	2.2 $\pm$ 1.3%	4.7 $\pm$ 0.4%
10 nM Glutamate	-1.1 $\pm$ 0.6%	-0.3 $\pm$ 0.2%	2.4 $\pm$ 0.3%

The results in the tables above demonstrate the ability of compounds of the present invention to potentiate the effects of mGlu<sub>2</sub> receptor agonists such as glutamate, APDC, and DCG IV. It is recognized that the compounds of the present invention would be expected to increase the effectiveness of glutamate and glutamate agonists of the mGlu<sub>2</sub> receptor. Thus, the potentiators of the present invention are expected to be useful in the treatment of various neurological and psychiatric disorders associated with glutamate dysfunction described to be treated herein and others that can be treated by such potentiators as are appreciated by those skilled in the art.

In one embodiment of the present invention provides methods of treating neurological and psychiatric disorders

-200-

associated with glutamate dysfunction, comprising:  
administering to a patient in need thereof an effective  
amount of a potentiator of metabotropic glutamate receptors.

Specifically, the present invention provides a method  
5 of treating neurological and psychiatric disorders  
associated with glutamate dysfunction, comprising:  
administering to a patient in need thereof an effective  
amount of a potentiator of mGlu<sub>2</sub> and/or mGlu<sub>3</sub> receptors,  
that is, the present invention provides methods using an  
10 effective amount of a potentiator of mGlu<sub>2</sub> receptors, an  
effective amount of a potentiator of mGlu<sub>3</sub> receptor, or an  
effective amount of a potentiator of mGlu<sub>2</sub> and mGlu<sub>3</sub>  
receptors.

In a preferred embodiment the present invention  
15 provides a method for treating migraine, comprising:  
administering to a patient in need thereof an effective  
amount of a metabotropic glutamate potentiator, in  
particular a potentiator of mGlu<sub>2</sub> and/or mGlu<sub>3</sub> receptors.

In another preferred embodiment the present invention  
20 provides a method for treating anxiety, comprising:  
administering to a patient in need thereof an effective  
amount of a metabotropic glutamate potentiator, in  
particular a potentiator of mGlu<sub>2</sub> and/or mGlu<sub>3</sub> receptors.

Particularly preferred anxiety disorders are  
25 generalized anxiety disorder, panic disorder, and obsessive  
compulsive disorder.

In another preferred embodiment the present invention  
provides a method for treating schizophrenia, comprising:  
administering to a patient in need thereof an effective  
30 amount of a metabotropic glutamate potentiator, in  
particular a potentiator of mGlu<sub>2</sub> and/or mGlu<sub>3</sub> receptors.

In yet another preferred embodiment the present  
invention provides a method for treating epilepsy,  
comprising: administering to a patient in need thereof an  
35 effective amount of a metabotropic glutamate potentiator, in  
particular a potentiator of mGlu<sub>2</sub> and/or mGlu<sub>3</sub> receptors.

-201-

Because the compounds of formula I enhance the normal physiological function of the mGlu receptors, the compounds of formula I are useful for the treatment of a variety of neurological and psychiatric disorders associated with

5 glutamate dysfunction, including: acute neurological and psychiatric disorders such as cerebral deficits subsequent to cardiac bypass surgery and grafting, stroke, cerebral ischemia, spinal cord trauma, head trauma, perinatal hypoxia, cardiac arrest, hypoglycemic neuronal damage,

10 dementia (including AIDS-induced dementia), Alzheimer's disease, Huntington's Chorea, amyotrophic lateral sclerosis, ocular damage, retinopathy, cognitive disorders, idiopathic and drug-induced Parkinson's disease, muscular spasms and disorders associated with muscular spasticity (including

15 tremors) seizures, epilepsy, convulsions, migraine (including migraine headache), urinary incontinence, substance tolerance, substance withdrawal (including, substances such as opiates, nicotine, tobacco products, alcohol, benzodiazepines, cocaine, sedatives, hypnotics,

20 etc.), psychosis, schizophrenia, anxiety (including generalized anxiety disorder, panic disorder, and obsessive compulsive disorder), mood disorders (including depression, mania, bipolar disorders), trigeminal neuralgia, hearing loss, tinnitus, macular degeneration of the eye, emesis,

25 brain edema, pain (including acute and chronic pain states, severe pain, intractable pain, neuropathic pain, and post-traumatic pain), tardive dyskinesia, sleep disorders (including narcolepsy), attention deficit/hyperactivity disorder, and conduct disorder.

30 At present, the fourth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV™) (1994, American Psychiatric Association, Washington, D.C.), provides a diagnostic tool for identifying many of the disorders described herein. The skilled artisan will

35 recognize that there are alternative nomenclatures, nosologies, and classification systems for neurological and

-202-

psychiatric disorders described herein and that these systems evolve with medical scientific progress.

Because the compounds of formula I potentiate mGlu receptor response, in particular mGlu<sub>2</sub> and/or mGlu<sub>3</sub> receptor  
5 response, to glutamate and glutamate agonists. Such agonists are easily recognized and some are available in the art. Schoepp, D.D., Jane, D.E., Monn, J.A., *Neuropharmacology* 38: 1431-1476, (1999).

Thus, a more particular embodiment, it is understood  
10 that the present invention extends to a method of potentiating the action of a glutamate receptor agonist at the Group II mGlu receptors, comprising administering to a patient in need thereof an effective amount of a mGluR<sub>2/3</sub> potentiator, in particular a compound of formula I, in  
15 combination with a potentiated amount of an mGlu receptor agonist. Such a combination may be advantageous in that it may augment the activity and selectivity of mGlu agonist.

As used herein, the term "patient" refers to a warm blooded animal such as a mammal which is afflicted with one  
20 or more neurological and psychiatric disorders associated with glutamate dysfunction. It is understood that guinea pigs, dogs, cats, rats, mice, horses, cattle, sheep, and humans are examples of animals within the scope of the meaning of the term. It is also understood that this  
25 invention relates specifically to the potentiation of mammalian metabotropic glutamate receptors.

It is also recognized that one skilled in the art may affect the neurological and psychiatric disorders by treating a patient presently afflicted with the disorders or  
30 by prophylactically treating a patient afflicted with the disorders with an effective amount of the compound of formula I. Thus, the terms "treatment" and "treating" are intended to refer to all processes wherein there may be a slowing, interrupting, arresting, controlling, or stopping  
35 of the progression of the neurological and psychiatric disorders described herein, but does not necessarily indicate a total elimination of all disorder symptoms, and

-203-

is intended to include prophylactic treatment of such neurological and psychiatric disorders.

As used herein, the term "effective amount" of a compound of formula I refers to an amount, that is, the dosage which is effective in treating the neurological and psychiatric disorders described herein.

An effective amount can be readily determined by the attending diagnostician, as one skilled in the art, by the use of conventional techniques and by observing results obtained under analogous circumstances. In determining an effective amount, the dose of a compound of formula I, a number of factors are considered by the attending diagnostician, including, but not limited to: the compound of formula I to be administered; the co-administration of an mGlu agonist, if used; the species of mammal; its size, age, and general health; the specific disorder involved; the degree of involvement or the severity of the disorder; the response of the individual patient; the mode of administration; the bioavailability characteristics of the preparation administered; the dose regimen selected; the use of other concomitant medication; and other relevant circumstances.

An effective amount of a compound of formula I is expected to vary from about 0.1 milligram per kilogram of body weight per day (mg/kg/day) to about 100 mg/kg/day. Preferred amounts are able to be determined by one skilled in the art.

As used herein, the term "potentiated amount" refers to an amount of an mGlu agonist, that is, the dosage of agonist which is effective in treating the neurological and psychiatric disorders described herein when administered in combination with an effective amount of a compound of formula I. A potentiated amount is expected to be less than the amount that is required to provided the same effect when the mGlu agonist is administered without an effective amount of a compound of formula I.



-204-

A potentiated amount can be readily determined by the attending diagnostician, as one skilled in the art, by the use of conventional techniques and by observing results obtained under analogous circumstances. In determining a  
5 potentiated amount, the dose of an mGlu agonist to be administered in combination with a compound of formula I, a number of factors are considered by the attending diagnostician, including, but not limited to: the mGlu agonist selected to be administered, including its potency  
10 and selectivity; the compound of formula I to be co-administered; the species of mammal; its size, age, and general health; the specific disorder involved; the degree of involvement or the severity of the disorder; the response of the individual patient; the modes of administration; the  
15 bioavailability characteristics of the preparations administered; the dose regimens selected; the use of other concomitant medication; and other relevant circumstances.

A potentiated amount of an mGlu agonist to be administered in combination with an effective amount of a  
20 compound of formula I is expected to vary from about 0.1 milligram per kilogram of body weight per day (mg/kg/day) to about 100 mg/kg/day and is expected to be less than the amount that is required to provided the same effect when administered without an effective amount of a compound of  
25 formula I. Preferred amounts of a co-administered mGlu agonist are able to be determined by one skilled in the art.

Of the neurological and psychiatric disorders associated with glutamate dysfunction which are treated according to the present invention, the treatment of  
30 migraine, anxiety, schizophrenia, and epilepsy are particularly preferred. Particularly preferred anxiety disorders are generalized anxiety disorder, panic disorder, and obsessive compulsive disorder.

Thus, in a preferred embodiment the present invention  
35 provides a method for treating migraine, comprising: administering to a patient in need thereof an effective

-205-

amount of a compound of formula I or a pharmaceutical composition thereof.

In one of the available sources of diagnostic tools, *Dorland's Medical Dictionary* (23<sup>rd</sup> Ed., 1982, W. B. Saunders Company, Philadelphia, PA), migraine is defined as a symptom complex of periodic headaches, usually temporal and unilateral, often with irritability, nausea, vomiting, constipation or diarrhea, and photophobia. As used herein the term "migraine" includes these periodic headaches, both temporal and unilateral, the associated irritability, nausea, vomiting, constipation or diarrhea, photophobia, and other associated symptoms. The skilled artisan will recognize that there are alternative nomenclatures, nosologies, and classification systems for neurological and psychiatric disorders, including migraine, and that these systems evolve with medical scientific progress.

In another preferred embodiment the present invention provides a method for treating anxiety, comprising: administering to a patient in need thereof an effective amount of a compound of formula I or a pharmaceutical composition thereof.

At present, the fourth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV<sup>TM</sup>) (1994, American Psychiatric Association, Washington, D.C.), provides a diagnostic tool including anxiety and related disorders. These include: panic disorder with or without agoraphobia, agoraphobia without history of panic disorder, specific phobia, social phobia, obsessive-compulsive disorder, post-traumatic stress disorder, acute stress disorder, generalized anxiety disorder, anxiety disorder due to a general medical condition, substance-induced anxiety disorder and anxiety disorder not otherwise specified. As used herein the term "anxiety" includes treatment of those anxiety disorders and related disorder as described in the DSM-IV. The skilled artisan will recognize that there are alternative nomenclatures, nosologies, and classification systems for neurological and psychiatric disorders, and

-206-

particular anxiety, and that these systems evolve with medical scientific progress. Thus, the term "anxiety" is intended to include like disorders that are described in other diagnostic sources.

5       A number of preclinical laboratory animal models for migraine and anxiety have been described. One commonly used model of migraine is the dural extravasation model that has been described by Phebus et al., Life Sci., 61(21), 2117-2126 (1997) which can be used to evaluate the present  
10    compounds.

#### Example B

##### Animal Model of Dural Protein Extravasation

Harlan Sprague-Dawley rats (250-350 g) were  
15    anesthetized with sodium pentobarbital intraperitoneally (65 mg/kg) and placed in a stereotaxic frame (David Kopf Instruments) with the incisor bar set at -3.5 mm. Following a midline sagittal scalp incision, two pairs of bilateral holes were drilled through the skull (6  
20    mm posteriorly, 2.0 and 4.0 mm laterally, all coordinates referenced to bregma). Pairs of stainless steel stimulating electrodes, insulated except at the tips (Rhodes Medical Systems, Inc.), were lowered through the holes in both hemispheres to a depth of 9 mm.

25       The femoral vein was exposed and a dose of the test compound was injected intravenously (i.v.) at a dosing volume of 1 ml/kg or, in the alternative, test compound was administered orally (p.o.) via gavage at a volume of 2 ml/kg. Approximately 8 minutes post i.v. injection, a 50  
30    mg/kg dose of Evans Blue, a fluorescent dye, was also injected intravenously. The Evans Blue complexed with proteins in the blood and functioned as a marker for protein extravasation. Exactly 10 minutes post-injection of the test compound, the left trigeminal ganglion was stimulated  
35    for 3 minutes at a current intensity of 1.0 mA (5 Hz, 4 msec duration) with a Model 273 potentiostat/ galvanostat (EG&G Princeton Applied Research).

-207-

Fifteen minutes following stimulation, the animals were killed and exsanguinated with 40 mL of saline. The top of the skull was removed to facilitate the collection of the dural membranes. The membrane samples were removed from both hemispheres, rinsed with water, and spread flat on microscopic slides. Once dried, the tissues were coverslipped with a 70% glycerol/water solution.

A fluorescence microscope (Zeiss) equipped with a grating monochromator and a spectrophotometer was used to quantify the amount of Evans Blue dye in each sample. An excitation wavelength of approximately 535 nm was utilized and the emission intensity at 600 nm was determined. The microscope was equipped with a motorized stage and also interfaced with a personal computer. This facilitated the computer-controlled movement of the stage with fluorescence measurements at 25 points (500 nm steps) on each dural sample. The mean and standard deviation of the measurements were determined by the computer.

The extravasation induced by the electrical stimulation of the trigeminal ganglion was an ipsilateral effect (i.e. occurs only on the side of the dura in which the trigeminal ganglion was stimulated). This allows the other (unstimulated) half of the dura to be used as a control. The ratio of the amount of extravasation in the dura from the stimulated side, over the amount of extravasation in the unstimulated side, was calculated. Control animals dosed only with saline, yielded a ratio of approximately 1.9. In contrast, a compound which effectively prevented the extravasation in the dura from the stimulated side would yield a ratio of approximately 1.0.

Data from such an experiment using a representative compound of the present invention is presented in the table below.

#### Inhibition of Dural Protein Extravasation

Compound	Dose	Route of Admin	Extrav. Ratio	n
----------	------	----------------	---------------	---

-208-

			$\pm$ s.e.m.	
vehicle	1 mL/kg	i.v.	1.89 $\pm$ 0.08	9
Example 16	10 pg/kg	i.v.	1.87 $\pm$ 0.07	4
Example 16	100 pg/kg	i.v.	1.37 $\pm$ 0.07	3
Example 16	1 ng/kg	i.v.	1.04 $\pm$ 0.05	3
Example 16	100 ng/kg	i.v.	0.99 $\pm$ 0.004	3

The fear potentiated startle response model has been extensively used as a model of anxiety and can be used to evaluate the present compounds. Davis, *Psychopharmacol.*, **62**, 1 (1979); Davis, *Behav. Neurosci.*, **100**, 814 (1986); Davis, *Tr. Pharmacol. Sci.*, **13**, 35 (1992).

#### Example C

##### Fear Potentiated Startle Paradigm

Male Sprague-Dawley rats weighing 325-400 g were purchased from Harlan Sprague-Dawley, Inc. (Cumberland, IN) and given a one week acclimation period before testing. Rats were individually housed with food and water ad libitum in an animal room on a 12-hour light/dark cycle with lights on between 6:00 A.M. and 6:00 P.M. The compound of Example 16 was prepared in a suspension of 5% ethanol, 0.5% CMC, 0.5% Tween 80 and 99% water. 2S-2-amino-2-(1S,2S-2-carboxycyclopropan-1-yl)-3-(xanth-9-yl) propionic acid was prepared in sterile water. Control rats were given the respective vehicle.

The fear potentiated startle paradigm is conducted over three consecutive days. All three days begin with a 5-minute adaptation period before the trial starts. On day one (baseline startle) after the adaptation period, the animal receives 30 trials of 120dB auditory noise. The mean startle amplitude ( $V_{\max}$ ) is used to assign animals to groups with similar means before conditioning begins. Day two consists of conditioning the animals. Each animal receives 0.5 mA of shock for 500 msec preceded by a 5 second presentation of light which remains on for the duration of the shock. Ten presentations of the light and shock are administered. Day three is the testing trial where drug administration occurs

-209-

prior to testing. Twenty-four hours after conditioning, startle testing sessions are conducted. Ten trials of acoustic startle (120 dB), non-light paired, are presented at the beginning of the session. This is followed by 20  
 5 random trials of the noise alone and 20 random trials of noise preceded by light. Excluding the first 10 trials, the startle response amplitudes for each trial type are averaged for each animal. Data is presented as the difference between light + noise and noise-alone. Differences in startle  
 10 response amplitudes were analyzed by JMP statistical software using a One-way Anova (analysis of variance, t-test). Group differences were considered to be significant at  $p < 0.05$ .

Data from such an experiment using a representative  
 15 compound of the present invention is presented in the table below. In the experiment shown below the compound of Example 16 was given i.p. 30 minutes prior to testing.

## Dose Response Experiment

Compound of Example 16 (mg/kg)	Difference in Startle Amplitude ( $V_{\max}$ ) + S.E.
Exp. 1 Control	188 + 47
0.03	108 + 41
0.3	154 + 38
3.0	21 + 24*
Exp. 2 Control	455 + 106
1.0	120 + 87*
3.0	148 + 86*
10.0	105 + 91*

\* $P < 0.05$ 

In another preferred embodiment the present invention provides a method for treating epilepsy, comprising:  
 25 administering to a patient in need thereof an effective amount of a compound of formula I or a pharmaceutical composition thereof.

At present, there are several types and subtypes of seizures associated with epilepsy, including idiopathic,

-210-

symptomatic, and cryptogenic. These epileptic seizures can be focal (partial) or generalized. They can also be simple or complex. Epilepsy is described in the art, such as Epilepsy: A comprehensive textbook. Ed. by Jerome Engel, Jr. and Timothy A. Pedley. (Lippincott-Raven, Philadelphia, 1997). At present, the International Classification of Diseases, Ninth Revision, (ICD-9) provides a diagnostic tool including epilepsy and related disorders. These include: generalized nonconvulsive epilepsy, generalized convulsive epilepsy, petit mal status epilepticus, grand mal status epilepticus, partial epilepsy with impairment of consciousness, partial epilepsy without impairment of consciousness, infantile spasms, epilepsy partialis continua, other forms of epilepsy, epilepsy, unspecified, NOS. As used herein the term "epilepsy" includes these all types and subtypes. The skilled artisan will recognize that there are alternative nomenclatures, nosologies, and classification systems for neurological and psychiatric disorders, including epilepsy, and that these systems evolve with medical scientific progress.

Various electroshock-induces models has been extensively used as a model of seizure disorders.

#### Example D

##### Electroshock-induced seizures

Application of electrical stimulation by corneal electrodes to mice can induce tonic hindlimb-extensor seizures. Blockade of tonic extensor seizures induced by electroshock is considered predictive for drugs which block seizure propagation and may be effective in preventing various seizures in humans, including epileptic seizures.

Vehicle or a dose of a test drug are administered to groups of 5 to 10 mice each. Thirty minutes later, electroshock (10 mA, 0.2 sec duration) is administered by transcorneal electrodes. The number of mice exhibiting tonic extensor seizures in each group is recorded. The data

-211-

are reported as the percentage of mice which are protected from seizures.

Data from such an experiment using a representative compound of the present invention is presented in the table  
5 below.

Compound of Example 590 (mg/kg, ip)	Percent Protection
vehicle	0%
30	60 %
100	100 %


10



-212-

WHAT IS CLAIMED IS:

1. A compound of the formula



wherein

$R_1$  is selected from the group consisting of  $-C(O)R_3$ ,  $-C(O)OR_4$ , and  $-SO_2R_5$

wherein  $R_3$  is selected from the group consisting of alkyl and cycloalkyl,  $R_4$  is selected from the group consisting of alkyl and cycloalkyl,  $R_5$  is selected from the group consisting of alkyl, cycloalkyl, and fluorinated alkyl;

$R_2$  is from 1 to 3 substituents independently selected from the group consisting of hydrogen, hydroxy, trisubstituted silyloxy, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkoxy, substituted alkoxy, cycloalkoxy, substituted cycloalkoxy, cycloalkyl, substituted cycloalkyl, halogen, cyano, nitro, phenyl, substituted phenyl, pyridyloxy, thiophenoxy, substituted thiophenoxy, phenylsulfinyl, substituted phenylsulfinyl, phenylsulfonyl, substituted phenylsulfonyl, benzoyl, substituted benzoyl, phenoxy, and substituted phenoxy;

or

two  $R_2$  substituents are taken together, on adjacent positions, to form a fused cycloalkyl or a methylenedioxy ring, and one  $R_2$  substituent is selected

-213-

from the group consisting of hydrogen, hydroxy,  
trisubstituted silyloxy, alkyl, substituted alkyl,  
alkenyl, substituted alkenyl, alkoxy, substituted  
alkoxy, cycloalkoxy, substituted cycloalkoxy,  
5 cycloalkyl, substituted cycloalkyl, halogen, cyano,  
nitro, phenyl, substituted phenyl, pyridyloxy,  
thiophenoxy, substituted thiophenoxy, phenylsulfinyl,  
substituted phenylsulfinyl, phenylsulfonyl, substituted  
phenylsulfonyl, benzoyl, substituted benzoyl, phenoxy,  
10 and substituted phenoxy;

R<sub>6</sub> is from 1 to 2 substituents independently selected  
from the group consisting of hydrogen, alkyl, alkoxy,  
trifluoromethyl, halogen, phenoxy, and substituted  
15 phenoxy;

X is selected from the group consisting of a bond,  
-CH<sub>2</sub>-, -CHR<sub>7</sub>-, and -CH<sub>2</sub>CH<sub>2</sub>- wherein R<sub>7</sub> is lower alkyl;

20 Y is selected from the group consisting of a bond,  
-CH<sub>2</sub>-, -CHR<sub>8</sub>-, -CH<sub>2</sub>CH<sub>2</sub>-, -CHR<sub>9</sub>CH<sub>2</sub>-, and -CH<sub>2</sub>CHR<sub>9</sub> wherein  
R<sub>8</sub> is lower alkyl and R<sub>9</sub> is lower alkyl;

25 and the pharmaceutically acceptable salts thereof and  
the pyridyl N-oxides thereof.

2. A compound according to Claim 1 wherein the  
pyridyl moiety is attached to X at the 3-position.

3. A compound according to Claim 2 wherein R<sub>1</sub> is  
-SO<sub>2</sub>R<sub>5</sub>.

30 4. A compound according to Claim 1 wherein the  
pyridyl moiety is attached to X at the 4-position.

5. A compound according to Claim 2 wherein R<sub>1</sub> is  
-C(O)OR<sub>4</sub>.

-214-

6. A compound according to Claim 3 or 5 wherein  $R_2$  is from 1 to 3 substituents independently selected from the group consisting of hydrogen, hydroxy, trisubstituted silyloxy, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkoxy, substituted alkoxy, cycloalkoxy, substituted cycloalkoxy, cycloalkyl, substituted cycloalkyl, halogen, cyano, nitro, phenyl, substituted phenyl, pyridyloxy, thiophenoxy, substituted thiophenoxy, phenylsulfinyl, substituted phenylsulfinyl, phenylsulfonyl, substituted phenylsulfonyl, benzoyl, substituted benzoyl, phenoxy, and substituted phenoxy.

7. A compound according to Claim 6 wherein  $R_2$  is selected from group consisting of substituted alkyl, phenyl, substituted phenyl, thiophenoxy, substituted thiophenoxy, phenoxy, and substituted phenoxy.

8. A compound according to Claim 7 wherein the substituted alkyl is selected from the group consisting of benzyl and substituted benzyl

9. A compound according to Claim 7 wherein  $R_2$  is selected from group consisting of phenyl and substituted phenyl.

10. A compound according to Claim 7 wherein  $R_2$  is selected from group consisting of phenoxy and substituted phenoxy.

11. A compound according to any of Claims 1-10 wherein at least one of the  $R_2$  substituents is attached in the 3-position.

12. A compound according to any of Claims 1-10 wherein at least one of the  $R_2$  substituents is attached in the 4-position.

13. A compound according to Claim 1 wherein at least 1  $R_2$  is substituted benzyl, substituted phenyl, or substituted phenoxy and one of the substituents is

-215-

selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, halogen, cyano, amido, and substituted amido.

14. A compound according to any one of Claims 1-13 wherein X is a -CH<sub>2</sub>-.

15. A compound according to any one of Claims 1-13 wherein Y is a bond.

16. A compound according to any one of Claims 1-13 wherein X is a -CH<sub>2</sub>- and Y is bond.

17. A compound of Claim 1 selected from the group consisting of N-(3-(2-Methoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylamine, N-(3-(2-Benzyloxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylamine, N-(4-Phenoxy)phenyl)-N-(ethylsulfonyl)-2-(pyrid-3-yl)ethylamine, N-(4-Phenoxy)phenylethyl)-N-(ethylsulfonyl)pyrid-3-ylmethylamine, N-(4-Phenoxy)phenylmethyl)-N-(ethylsulfonyl)pyrid-3-ylmethylamine, N-(4-(Phenoxy)phenyl)-N-(isopropylsulfonyl)pyrid-3-ylmethylamine, N-(2-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylamine, N-(4-(Phenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylamine, N-(4-(2-Methoxyphenoxy)-3-chlorophenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylamine, N-(4-(2,6-Dimethoxyphenoxy)-3-chlorophenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylamine, N-(4-(2-Benzyloxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylamine, N-(4-(2-Methoxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylamine, N-(4-(2-Methoxyphenoxy)phenyl)-N-(cyclopropylsulfonyl)pyrid-3-ylmethylamine, N-(4-(2-Methoxyphenoxy)-3-chlorophenyl)-N-ethylsulfonyl)pyrid-3-ylmethylamine, N-(4-(2-Methoxyphenoxy)-3-

-216-

chlorophenyl)-N-cyclopropylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-Methoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-Ethoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-Ethoxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-Ethoxyphenoxy)phenyl)-N-(cyclopropylsulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-Methoxyphenoxy)phenyl)-N-(cyclopropylsulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-Methoxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-Hydroxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-Hydroxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(methylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(butylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(methoxycarbonyl)pyrid-2-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(methylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(butylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(methoxycarbonyl)pyrid-3-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(methylsulfonyl)pyrid-4-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(ethylsulfonyl)pyrid-4-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(butylsulfonyl)pyrid-4-ylmethylaniline, N-(4-

-217-

(Phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-4-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(methoxycarbonyl)pyrid-4-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(propionyl)pyrid-2-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(cyclopentylcarbonylpropionyl)pyrid-2-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(propionyl)pyrid-3-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(cyclopentylcarbonyl)pyrid-3-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(propionyl)pyrid-4-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(cyclopentylcarbonyl)pyrid-4-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(trifluoromethylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(trifluoromethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(trifluoromethylsulfonyl)pyrid-4-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(propylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(propylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(propylsulfonyl)pyrid-4-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(2-methylpropionyl)pyrid-2-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(2-methylpropionyl)pyrid-3-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(2-methylpropionyl)pyrid-4-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(acetyl)pyrid-2-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(acetyl)pyrid-3-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(acetyl)pyrid-4-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(cyclopropylcarbonyl)pyrid-2-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(cyclopropylcarbonyl)pyrid-3-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(cyclopropylcarbonyl)pyrid-4-ylmethylaniline, N-(4-(4-Benzyloxyphenoxy)phenyl)-N-(cyclopropylcarbonyl)pyrid-3-ylmethylaniline, N-(4-(4-

-218-

Cyclohexylphenoxy) phenyl) -N- (cyclopropylcarbonyl) pyrid-  
3-ylmethylaniline, N- (4- (4-Benzylphenoxy) phenyl) -N-  
(cyclopropylcarbonyl) pyrid-3-ylmethylaniline, N- (4- (4-  
Thiophenoxyphenoxy) phenyl) -N-  
5 (cyclopropylcarbonyl) pyrid-3-ylmethylaniline, N- (4- (4-  
Benzyloxyphenoxy) phenyl) -N- (methylsulfonyl) pyrid-3-  
ylmethylaniline, N- (4- (4-Cyclohexylphenoxy) phenyl) -N-  
(methylsulfonyl) pyrid-3-ylmethylaniline, N- (4- (4-  
Benzylphenoxy) phenyl) -N- (methylsulfonyl) pyrid-3-  
10 ylmethylaniline, N- (4- (4-Thiophenoxyphenoxy) phenyl) -N-  
(methylsulfonyl) pyrid-3-ylmethylaniline, N- (4- (4-  
Benzyloxyphenoxy) phenyl) -N- (ethylsulfonyl) pyrid-3-  
ylmethylaniline, N- (4- (4-Cyclohexylphenoxy) phenyl) -N-  
(ethylsulfonyl) pyrid-3-ylmethylaniline, N- (4- (4-  
15 Benzylphenoxy) phenyl) -N- (ethylsulfonyl) pyrid-3-  
ylmethylaniline, N- (4- (4-Thiophenoxyphenoxy) phenyl) -N-  
(ethylsulfonyl) pyrid-3-ylmethylaniline, N- (4- (4-  
Benzyloxyphenoxy) phenyl) -N- (2,2,2-  
trifluoroethylsulfonyl) pyrid-3-ylmethylaniline, N- (4- (4-  
20 Cyclohexylphenoxy) phenyl) -N- (2,2,2-  
trifluoroethylsulfonyl) pyrid-3-ylmethylaniline, N- (4- (4-  
Benzylphenoxy) phenyl) -N- (2,2,2-  
trifluoroethylsulfonyl) pyrid-3-ylmethylaniline, N- (4- (4-  
Thiophenoxyphenoxy) phenyl) -N- (2,2,2-  
25 trifluoroethylsulfonyl) pyrid-3-ylmethylaniline, N- (4- (4-  
Benzyloxyphenoxy) phenyl) -N-  
(trifluoromethylsulfonyl) pyrid-3-ylmethylaniline, N- (4-  
(4-Cyclohexylphenoxy) phenyl) -N-  
(trifluoromethylsulfonyl) pyrid-3-ylmethylaniline, N- (4-  
30 (4-Benzylphenoxy) phenyl) -N-  
(trifluoromethylsulfonyl) pyrid-3-ylmethylaniline, N- (4-  
(4-Thiophenoxyphenoxy) phenyl) -N-  
(trifluoromethylsulfonyl) pyrid-3-ylmethylaniline, N- (4-  
(4-Benzyloxyphenoxy) phenyl) -N- (propylsulfonyl) pyrid-3-

-219-

ylmethylanine, N-(4-(4-Cyclohexylphenoxy)phenyl)-N-(propylsulfonyl)pyrid-3-ylmethylanine, N-(4-(4-Benzylphenoxy)phenyl)-N-(propylsulfonyl)pyrid-3-ylmethylanine, N-(4-(4-Thiophenoxyphenoxy)phenyl)-N-(propylsulfonyl)pyrid-3-ylmethylanine, N-(4-(4-Benzylphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethylanine, N-(4-(4-Cyclohexylphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethylanine, N-(4-(4-Benzylphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethylanine, N-(4-(4-Thiophenoxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethylanine, N-(4-(4-Benzylphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethylanine, N-(4-(4-Cyclohexylphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethylanine, N-(4-(4-Benzylphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethylanine, N-(4-(4-Thiophenoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethylanine, N-(4-(4-(Pentoxo)phenyl)-N-(cyclopropylcarbonyl)pyrid-3-ylmethylanine, N-(4-(3,4-Methylenedioxyphenoxy)phenyl)-N-(cyclopropylcarbonyl)pyrid-3-ylmethylanine, N-(4-(Methoxy)phenyl)-N-(cyclopropylcarbonyl)pyrid-3-ylmethylanine, N-(4-(5,6,7,8-Tetrahydronaphth-2-yloxy)phenyl)-N-(cyclopropylcarbonyl)pyrid-3-ylmethylanine, N-(4-(4-Cyclohexyloxy)phenyl)-N-(cyclopropylcarbonyl)pyrid-3-ylmethylanine, N-(4-(4-Pyrid-2-yloxyphenoxy)phenyl)-N-(cyclopropylcarbonyl)pyrid-3-ylmethylanine, N-(4-(4-(Pentoxo)phenyl)-N-(methylsulfonyl)pyrid-3-ylmethylanine, N-(4-(3,4-Methylenedioxyphenoxy)phenyl)-N-(methylsulfonyl)pyrid-3-ylmethylanine, N-(4-(Methoxy)phenyl)-N-(methylsulfonyl)pyrid-3-ylmethylanine, N-(4-(5,6,7,8-Tetrahydronaphth-2-



-220-

yloxy)phenyl)-N-(methylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-Cyclohexyloxy)phenyl)-N-(methylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-Pyrid-2-yloxyphenoxy)phenyl)-N-(methylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(Pentoxo)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(3,4-Methylenedioxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(Methoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(5,6,7,8-Tetrahydronaphth-2-yloxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-Cyclohexyloxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-Pyrid-2-yloxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(Pentoxo)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(3,4-Methylenedioxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(Methoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(5,6,7,8-Tetrahydronaphth-2-yloxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-Cyclohexyloxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-Pyrid-2-yloxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(Pentoxo)phenyl)-N-(trifluoromethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(3,4-Methylenedioxyphenoxy)phenyl)-N-(trifluoromethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(Methoxy)phenyl)-N-(trifluoromethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(5,6,7,8-Tetrahydronaphth-2-yloxy)phenyl)-N-(trifluoromethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-Cyclohexyloxy)phenyl)-N-(trifluoromethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-Pyrid-2-yloxyphenoxy)phenyl)-N-

-221-

(trifluoromethylsulfonyl)pyrid-3-ylmethanamine, N-(4-(Pentoxy)phenyl)-N-(propylsulfonyl)pyrid-3-ylmethanamine, N-(4-(3,4-Methylenedioxyphenoxy)phenyl)-N-(propylsulfonyl)pyrid-3-ylmethanamine, N-(4-(Methoxy)phenyl)-N-(propylsulfonyl)pyrid-3-ylmethanamine, N-(4-(5,6,7,8-Tetrahydronaphth-2-yloxy)phenyl)-N-(propylsulfonyl)pyrid-3-ylmethanamine, N-(4-(4-Cyclohexyloxy)phenyl)-N-(propylsulfonyl)pyrid-3-ylmethanamine, N-(4-(4-Pyrid-2-yloxyphenoxy)phenyl)-N-(propylsulfonyl)pyrid-3-ylmethanamine, N-(4-(Pentoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethanamine, N-(4-(3,4-Methylenedioxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethanamine, N-(4-(Methoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethanamine, N-(4-(5,6,7,8-Tetrahydronaphth-2-yloxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethanamine, N-(4-(4-Cyclohexyloxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethanamine, N-(4-(4-Pyrid-2-yloxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethanamine, N-(4-(Pentoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethanamine, N-(4-(3,4-Methylenedioxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethanamine, N-(4-(Methoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethanamine, N-(4-(5,6,7,8-Tetrahydronaphth-2-yloxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethanamine, N-(4-(4-Cyclohexyloxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethanamine, N-(4-(4-Pyrid-2-yloxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethanamine, N-(3-(Phenoxy)phenyl)-N-(cyclopropylcarbonyl)pyrid-3-ylmethanamine, N-(2-(Phenoxy)phenyl)-N-(cyclopropylcarbonyl)pyrid-3-ylmethanamine, N-(4-(4-

-222-

Chlorophenoxy)phenyl)-N-(cyclopropylcarbonyl)pyrid-3-ylmethylaniline, N-(4-(3-Trifluoromethylphenoxy)phenyl)-N-(cyclopropylcarbonyl)pyrid-3-ylmethylaniline, N-(4-(2-Methoxyphenoxy)phenyl)-N-(cyclopropylcarbonyl)pyrid-3-ylmethylaniline, N-(4-(4-Cyanophenoxy)phenyl)-N-(cyclopropylcarbonyl)pyrid-3-ylmethylaniline, N-(4-(2,4-Difluorophenoxy)phenyl)-N-(cyclopropylcarbonyl)pyrid-3-ylmethylaniline, N-(3-(Phenoxy)phenyl)-N-(methylsulfonyl)pyrid-3-ylmethylaniline, N-(2-(Phenoxy)phenyl)-N-(methylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-Chlorophenoxy)phenyl)-N-(methylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(3-Trifluoromethylphenoxy)phenyl)-N-(methylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-Methoxyphenoxy)phenyl)-N-(methylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-Cyanophenoxy)phenyl)-N-(methylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2,4-Difluorophenoxy)phenyl)-N-(methylsulfonyl)pyrid-3-ylmethylaniline, N-(3-(Phenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethylaniline, N-(2-(Phenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(4-Chlorophenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(3-Trifluoromethylphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(2-Methoxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(4-Cyanophenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(2,4-Difluorophenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethylaniline, N-(3-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethylaniline, N-(2-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(4-Chlorophenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(3-Trifluoromethylphenoxy)phenyl)-N-(2,2,2-

-223-

trifluoroethylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(2-Methoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(4-Cyanophenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(2,4-Difluorophenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethylaniline, N-(3-(Phenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(2-(Phenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-Chlorophenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(3-Trifluoromethylphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-Methoxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-Cyanophenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2,4-Difluorophenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(3-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(2-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-Chlorophenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(3-Trifluoromethylphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-Methoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-Cyanophenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2,4-Difluorophenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(3-(Phenoxy)phenyl)-N-(trifluoromethylsulfonyl)pyrid-3-ylmethylaniline, N-(2-(Phenoxy)phenyl)-N-(trifluoromethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-Chlorophenoxy)phenyl)-N-

-224-

(trifluoromethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(3-Trifluoromethylphenoxy)phenyl)-N-(trifluoromethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-Methoxyphenoxy)phenyl)-N-(trifluoromethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-Cyanophenoxy)phenyl)-N-(trifluoromethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2,4-Difluorophenoxy)phenyl)-N-(trifluoromethylsulfonyl)pyrid-3-ylmethylaniline, N-(3-(Phenoxy)phenyl)-N-(propylsulfonyl)pyrid-3-ylmethylaniline, N-(2-(Phenoxy)phenyl)-N-(propylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-Chlorophenoxy)phenyl)-N-(propylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(3-Trifluoromethylphenoxy)phenyl)-N-(propylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-Methoxyphenoxy)phenyl)-N-(propylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-Cyanophenoxy)phenyl)-N-(propylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2,4-Difluorophenoxy)phenyl)-N-(propylsulfonyl)pyrid-3-ylmethylaniline, N-(2-Chloro-5-methyl-4-(phenoxy)phenyl)-N-(cyclopropylcarbonyl)pyrid-3-ylmethylaniline, N-(3-Chloro-4-(phenoxy)phenyl)-N-(cyclopropylcarbonyl)pyrid-3-ylmethylaniline, N-(2-Methyl-4-(phenoxy)phenyl)-N-(cyclopropylcarbonyl)pyrid-3-ylmethylaniline, N-(3-Trifluoromethyl-4-(phenoxy)phenyl)-N-(cyclopropylcarbonyl)pyrid-3-ylmethylaniline, N-(4-(4-Nitrophenoxy)phenyl)-N-(cyclopropylcarbonyl)pyrid-3-ylmethylaniline, N-(2-Chloro-5-methyl-4-(phenoxy)phenyl)-N-(methylsulfonyl)pyrid-3-ylmethylaniline, N-(3-Chloro-4-(phenoxy)phenyl)-N-(methylsulfonyl)pyrid-3-ylmethylaniline, N-(2-Methyl-4-(phenoxy)phenyl)-N-(methylsulfonyl)pyrid-3-ylmethylaniline, N-(3-Trifluoromethyl-4-(phenoxy)phenyl)-N-(methylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-

-225-

Nitrophenoxy)phenyl)-N-(methylsulfonyl)pyrid-3-ylmethylaniline, N-(2-Chloro-5-methyl-4-(phenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(3-Chloro-4-(phenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(2-Methyl-4-(phenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(3-Trifluoromethyl-4-(phenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-Nitrophenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(2-Chloro-5-methyl-4-(phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(3-Chloro-4-(phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(2-Methyl-4-(phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(3-Trifluoromethyl-4-(phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-Nitrophenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(2-Chloro-5-methyl-4-(phenoxy)phenyl)-N-(trifluoromethylsulfonyl)pyrid-3-ylmethylaniline, N-(3-Chloro-4-(phenoxy)phenyl)-N-(trifluoromethylsulfonyl)pyrid-3-ylmethylaniline, N-(2-Methyl-4-(phenoxy)phenyl)-N-(trifluoromethylsulfonyl)pyrid-3-ylmethylaniline, N-(3-Trifluoromethyl-4-(phenoxy)phenyl)-N-(trifluoromethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-Nitrophenoxy)phenyl)-N-(trifluoromethylsulfonyl)pyrid-3-ylmethylaniline, N-(2-Chloro-5-methyl-4-(phenoxy)phenyl)-N-(propylsulfonyl)pyrid-3-ylmethylaniline, N-(3-Chloro-4-(phenoxy)phenyl)-N-(propylsulfonyl)pyrid-3-ylmethylaniline, N-(2-Methyl-4-(phenoxy)phenyl)-N-(propylsulfonyl)pyrid-3-ylmethylaniline, N-(3-

-226-

Trifluoromethyl-4-(phenoxy)phenyl)-N-(propylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-Nitrophenoxy)phenyl)-N-(propylsulfonyl)pyrid-3-ylmethylaniline, N-(2-Chloro-5-methyl-4-(phenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethylaniline, N-(3-Chloro-4-(phenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethylaniline, N-(2-Methyl-4-(phenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethylaniline, N-(3-Trifluoromethyl-4-(phenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(4-Nitrophenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethylaniline, N-(2-Chloro-5-methyl-4-(phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethylaniline, N-(3-Chloro-4-(phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethylaniline, N-(2-Methyl-4-(phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethylaniline, N-(3-Trifluoromethyl-4-(phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(4-Nitrophenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)-6-methoxypyrid-3-ylmethylaniline, N-(3-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)-6-methoxypyrid-3-ylmethylaniline, N-(4-(2-Methoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)-6-methoxypyrid-3-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(ethylsulfonyl)-6-methoxypyrid-3-ylmethylaniline, N-(3-(Phenoxy)phenyl)-N-(ethylsulfonyl)-6-methoxypyrid-3-ylmethylaniline, N-(4-(2-Methoxyphenoxy)phenyl)-N-(ethylsulfonyl)-6-methoxypyrid-3-ylmethylaniline, N-(4-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)-6-methylpyrid-3-ylmethylaniline, N-(3-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)-6-methylpyrid-3-

-227-

ylmethyllamine, N-(4-(2-Methoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)-6-methylpyrid-3-ylmethyllamine, N-(4-(Phenoxy)phenyl)-N-(ethylsulfonyl)-6-methylpyrid-3-ylmethyllamine, N-(3-(Phenoxy)phenyl)-N-(ethylsulfonyl)-6-methylpyrid-3-ylmethyllamine, N-(4-(2-Methoxyphenoxy)phenyl)-N-(ethylsulfonyl)-6-methylpyrid-3-ylmethyllamine, N-(4-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)-6-(4-chlorophenoxy)pyrid-3-ylmethyllamine, N-(3-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)-6-(4-chlorophenoxy)pyrid-3-ylmethyllamine, N-(4-(2-Methoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)-6-(4-chlorophenoxy)pyrid-3-ylmethyllamine, N-(4-(Phenoxy)phenyl)-N-(ethylsulfonyl)-6-(4-chlorophenoxy)pyrid-3-ylmethyllamine, N-(3-(Phenoxy)phenyl)-N-(ethylsulfonyl)-6-(4-chlorophenoxy)pyrid-3-ylmethyllamine, N-(4-(2-Methoxyphenoxy)phenyl)-N-(ethylsulfonyl)-6-(4-chlorophenoxy)pyrid-3-ylmethyllamine, N-(4-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)-2-chloropyrid-3-ylmethyllamine, N-(3-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)-2-chloropyrid-3-ylmethyllamine, N-(4-(2-Methoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)-2-chloropyrid-3-ylmethyllamine, N-(4-(Phenoxy)phenyl)-N-(ethylsulfonyl)-2-chloropyrid-3-ylmethyllamine, N-(3-(Phenoxy)phenyl)-N-(ethylsulfonyl)-2-chloropyrid-3-ylmethyllamine, N-(4-(2-Methoxyphenoxy)phenyl)-N-(ethylsulfonyl)-2-chloropyrid-3-ylmethyllamine, N-(4-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)-2-phenoxypyrid-3-ylmethyllamine, N-(3-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)-2-phenoxypyrid-3-ylmethyllamine, N-(4-(2-Methoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)-2-phenoxypyrid-3-ylmethyllamine, N-(4-(Phenoxy)phenyl)-N-(ethylsulfonyl)-



-228-

2-phenoxy)pyrid-3-ylmethanamine, N-(3-(Phenoxy)phenyl)-  
N-(ethylsulfonyl)-2-phenoxy)pyrid-3-ylmethanamine, N-(4-  
(2-Methoxyphenoxy)phenyl)-N-(ethylsulfonyl)-2-  
phenoxy)pyrid-3-ylmethanamine, N-(4-(Phenoxy)phenyl)-N-  
5 (2,2,2-trifluoroethylsulfonyl)-6-chloropyrid-3-  
ylmethanamine, N-(3-(Phenoxy)phenyl)-N-(2,2,2-  
trifluoroethylsulfonyl)-6-chloropyrid-3-  
ylmethanamine, N-(4-(2-Methoxyphenoxy)phenyl)-N-(2,2,2-  
trifluoroethylsulfonyl)-6-chloropyrid-3-  
10 ylmethanamine, N-(4-(Phenoxy)phenyl)-N-(ethylsulfonyl)-  
6-chloropyrid-3-ylmethanamine, N-(3-(Phenoxy)phenyl)-N-  
(ethylsulfonyl)-6-chloropyrid-3-ylmethanamine, N-(4-(2-  
Methoxyphenoxy)phenyl)-N-(ethylsulfonyl)-6-chloropyrid-  
3-ylmethanamine, N-(4-(Phenoxy)phenyl)-N-(2,2,2-  
15 trifluoroethylsulfonyl)-2-chloro-6-methylpyrid-3-  
ylmethanamine, N-(3-(Phenoxy)phenyl)-N-(2,2,2-  
trifluoroethylsulfonyl)-2-chloro-6-methylpyrid-3-  
ylmethanamine, N-(4-(2-Methoxyphenoxy)phenyl)-N-(2,2,2-  
trifluoroethylsulfonyl)-2-chloro-6-methylpyrid-3-  
20 ylmethanamine, N-(4-(Phenoxy)phenyl)-N-(ethylsulfonyl)-  
2-chloro-6-methylpyrid-3-ylmethanamine, N-(3-  
(Phenoxy)phenyl)-N-(ethylsulfonyl)-2-chloro-6-  
methylpyrid-3-ylmethanamine, N-(4-(2-  
Methoxyphenoxy)phenyl)-N-(ethylsulfonyl)-2-chloro-6-  
25 methylpyrid-3-ylmethanamine, N-(4-(Phenoxy)phenyl)-N-  
(cyclopropylsulfonyl)pyrid-3-ylmethanamine, N-(4-(2-  
Methoxyphenoxy)phenyl)phenyl)-N-  
(cyclopropylsulfonyl)pyrid-3-ylmethanamine, N-(4-(2-  
Methoxyphenoxy)phenyl)-N-(2,2,2-  
30 trifluoroethylsulfonyl)pyrid-4-ylmethanamine, N-(4-(2-  
Methoxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-4-  
ylmethanamine, N-(4-(2,6-Dimethoxyphenoxy)phenyl)-N-  
(cyclopropylsulfonyl)pyrid-3-ylmethanamine, N-(4-(2,6-  
Dimethoxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-

-229-

ylmethylanine, N-(4-(2,6-Dimethoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylanine, N-(4-(2,6-Dimethoxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethylanine, N-(4-(2,6-Dimethoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethylanine, N-(4-(2,6-Dimethoxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-4-ylmethylanine, N-(4-(2,6-Dimethoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-4-ylmethylanine, N-(4-(2-Phenoxy-6-chloro)phenylmethyl)-N-(cyclopropylsulfonyl)pyrid-3-ylmethylanine, N-(4-(2-Phenoxy-6-chloro)phenylmethyl)-N-(ethylsulfonyl)pyrid-3-ylmethylanine, N-(4-(2-Phenoxy-6-chloro)phenylmethyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylanine, N-(4-(2-Phenoxy-6-chloro)phenylmethyl)-N-(ethylsulfonyl)pyrid-2-ylmethylanine, N-(4-(2-Phenoxy-6-chloro)phenylmethyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethylanine, N-(4-(2-Phenoxy-6-chloro)phenylmethyl)-N-(ethylsulfonyl)pyrid-4-ylmethylanine, N-(4-(2-Phenoxy-6-chloro)phenylmethyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-4-ylmethylanine, N-(4-(2,6-Dichlorophenoxy)phenyl)-N-(cyclopropylsulfonyl)pyrid-3-ylmethylanine, N-(4-(2,6-Dichlorophenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylanine, N-(4-(2,6-Dichlorophenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylanine, N-(4-(2,6-Dichlorophenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethylanine, N-(4-(2,6-Dichlorophenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethylanine, N-(4-(2,6-Dichlorophenoxy)phenyl)-N-(ethylsulfonyl)pyrid-4-ylmethylanine, N-(4-(2,6-Dichlorophenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-4-ylmethylanine, N-

-230-

(4-(2-Isopropoxyphenoxy)phenyl)-N-(cyclopropylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-Isopropoxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-Isopropoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-Isopropoxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(2-Isopropoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(2-Isopropoxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-4-ylmethylaniline, N-(4-(2-Isopropoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-4-ylmethylaniline, N-(4-(2-Fluoro-6-methoxyphenoxy)phenyl)-N-(cyclopropylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-Fluoro-6-methoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-Fluoro-6-methoxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-Fluoro-6-methoxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(2-Fluoro-6-methoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(2-Fluoro-6-methoxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-4-ylmethylaniline, N-(4-(2-Fluoro-6-methoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-4-ylmethylaniline, N-(4-(2-Trifluoromethylphenoxy)phenyl)-N-(cyclopropylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-Trifluoromethylphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-Trifluoromethylphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-Trifluoromethylphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(2-Trifluoromethylphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-2-ylmethylaniline, N-(4-(2-

-231-

Trifluoromethylphenoxy) phenyl) -N- (ethylsulfonyl) pyrid-  
4-ylmethylaniline, N- (4- (2-  
Trifluoromethylphenoxy) phenyl) -N- (2,2,2-  
trifluoroethylsulfonyl) pyrid-4-ylmethylaniline, N- (4- (2-  
5 Isopropyl-5-methylphenoxy) phenyl) -N-  
(cyclopropylsulfonyl) pyrid-3-ylmethylaniline, N- (4- (2-  
Isopropyl-5-methylphenoxy) phenyl) -N- (2,2,2-  
trifluoroethylsulfonyl) pyrid-3-ylmethylaniline, N- (4- (2-  
Isopropyl-5-methylphenoxy) phenyl) -N-  
10 (ethylsulfonyl) pyrid-3-ylmethylaniline, N- (4- (2-  
Isopropyl-5-methylphenoxy) phenyl) -N-  
(ethylsulfonyl) pyrid-2-ylmethylaniline, N- (4- (2-  
Isopropyl-5-methylphenoxy) phenyl) -N- (2,2,2-  
trifluoroethylsulfonyl) pyrid-2-ylmethylaniline, N- (4- (2-  
15 Isopropyl-5-methylphenoxy) phenyl) -N-  
(ethylsulfonyl) pyrid-4-ylmethylaniline, N- (4- (2-  
Isopropyl-5-methylphenoxy) phenyl) -N- (2,2,2-  
trifluoroethylsulfonyl) pyrid-4-ylmethylaniline, N- (4- (2-  
Cyclopentylphenoxy) phenyl) -N-  
20 (cyclopropylsulfonyl) pyrid-3-ylmethylaniline, N- (4- (2-  
Cyclopentylphenoxy) phenyl) -N- (2,2,2-  
trifluoroethylsulfonyl) pyrid-3-ylmethylaniline, N- (4- (2-  
Cyclopentylphenoxy) phenyl) -N- (ethylsulfonyl) pyrid-3-  
ylmethylaniline, N- (4- (2-Cyclopentylphenoxy) phenyl) -N-  
25 (ethylsulfonyl) pyrid-2-ylmethylaniline, N- (4- (2-  
Cyclopentylphenoxy) phenyl) -N- (2,2,2-  
trifluoroethylsulfonyl) pyrid-2-ylmethylaniline, N- (4- (2-  
Cyclopentylphenoxy) phenyl) -N- (ethylsulfonyl) pyrid-4-  
ylmethylaniline, N- (4- (2-Cyclopentylphenoxy) phenyl) -N-  
30 (2,2,2-trifluoroethylsulfonyl) pyrid-4-ylmethylaniline, N-  
(4- (Phenoxy) phenyl) -N- (2,2,2-trifluoroethylsulfonyl) -2-  
(2,2,2-trifluoroethoxy) pyrid-3-ylmethylaniline, N- (4- (2-  
Methoxyphenoxy) phenyl) -N- (2,2,2-  
trifluoroethylsulfonyl) -2- (2,2,2-trifluoroethoxy) pyrid-

-232-

3-ylmethylanine, N-(4-(2-Isopropoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)-2-(2,2,2-trifluoroethoxy)pyrid-3-ylmethylanine, N-(4-(2-Fluoro-6-methoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)-2-(2,2,2-trifluoroethoxy)pyrid-3-ylmethylanine, , N-(4-(4-Hydroxyphenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylanine, N-(3-(2-Methoxyphenylsulfinyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(2,4-Difluorophenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(2-Methoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(3-Benzyloxyphenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(3-Hydroxyphenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylanine, N-(4-t-Butyldimethylsilyloxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-Hydroxyphenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-t-Butyldimethylsilyloxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylanine, N-(4-Hydroxyphenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(Phenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)-1-oxypyrid-3-ylmethylanine, N-(4-(2-Cyanophenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(2-Cyanophenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(3-(2-Methoxyphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylanine, N-(3-(2-phenylphenoxy)phenyl)-N-(ethylsulfonyl)pyrid-3-ylmethylanine, N-(4-Bromophenyl)-N-(ethanesulfonyl)pyrid-3-

-233-

ylmethylanine, N-(4-(2-Fluorophenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(3-Fluorophenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(4-Fluorophenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(2,4-Difluorophenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(2-trifluoromethoxyphenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(4-fluoro-2-methylphenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(4-trifluoromethoxyphenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(2,4-dimethylphenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(4-methylphenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(4-isopropylphenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(4-t-butylphenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(2-methylphenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(2-Amidophenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(3-Methylsulfinylphenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(4-t-butyl-2-fluorophenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(4-fluoro-2-n-propylphenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(4-Carbonylmethoxyphenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(4-amidomethylphenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(3-t-butylphenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(2-methoxyphenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(3-methylphenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(2-

-234-

chlorophenoxy) phenyl) -N- (ethanesulfonyl) pyrid-3-ylmethylamine, N- (4- (4-Amidophenoxy) phenyl) -N- (ethanesulfonyl) pyrid-3-ylmethylamine, N- (4- (2-methylphenoxy) phenyl) -N- (ethanesulfonyl) pyrid-3-ylmethylamine, N- (4- (4-Methylsulfonylphenoxy) phenyl) -N- (ethanesulfonyl) pyrid-3-ylmethylamine, N- (4- (2- (prop-2-yl) phenoxy) phenyl) -N- (ethanesulfonyl) pyrid-3-ylmethylamine, N- (4- (2-ethylphenoxy) phenyl) -N- (ethanesulfonyl) pyrid-3-ylmethylamine, N- (4- (4-Sulfonylamidophenoxy) phenyl) -N- (ethanesulfonyl) pyrid-3-ylmethylamine, N- (4- (2,4-Difluorophenoxy) phenyl) -N- (ethanesulfonyl) -1-pyrid-3-ylethylamine, N- (3-Benzyloxyphenoxy) phenyl) -N- (ethanesulfonyl) pyrid-3-ylmethylamine, N- (3- (Bromophenyl) -N- (ethylsulfonyl) pyrid-3-ylmethylamine, N- (3- (2-fluorophenoxy) phenyl) -N- (ethylsulfonyl) pyrid-3-ylmethylamine, N- (3- (3-fluorophenoxy) phenyl) -N- (ethylsulfonyl) pyrid-3-ylmethylamine, N- (3- (4-fluorophenoxy) phenyl) -N- (ethylsulfonyl) pyrid-3-ylmethylamine, N- (3- (2, 4-difluorophenoxy) phenyl) -N- (ethylsulfonyl) pyrid-3-ylmethylamine, N- (3- (2-ethoxyphenoxy) phenyl) -N- (ethylsulfonyl) pyrid-3-ylmethylamine, N- (3- (2-phenyl phenoxy) phenyl) -N- (ethylsulfonyl) pyrid-3-ylmethylamine, N- (3- (4-trifluoromethoxyphenoxy) phenyl) -N- (ethylsulfonyl) pyrid-3-ylmethylamine, N- (3- (4-methylphenoxy) phenyl) -N- (ethylsulfonyl) pyrid-3-ylmethylamine, N- (3- (2-methyl-4-fluorophenoxy) phenyl) -N- (ethylsulfonyl) pyrid-3-ylmethylamine, N- (3- (2-trifluoromethoxyphenoxy) phenyl) -N- (ethylsulfonyl) pyrid-3-ylmethylamine, N- (3- (4-Amidomethylphenoxy) phenyl) -N- (ethylsulfonyl) pyrid-3-ylmethylamine, N- (3- (4-Methylsulfonylphenoxy) phenyl) -N- (ethylsulfonyl) pyrid-3-

-235-

ylmethylamine, N- (3- (4-Methylsulfinylphenoxy) phenyl) -N-  
(ethylsulfonyl)pyrid-3-ylmethylamine, N- (3- (4-t-  
butylphenoxy) phenyl) -N- (ethylsulfonyl)pyrid-3-  
ylmethylamine, N- (3- (3-methylphenoxy) phenyl) -N-  
5 (ethylsulfonyl)pyrid-3-ylmethylamine, N- (3- (2-n-propyl-  
4-fluorophenoxy) phenyl) -N- (ethylsulfonyl)pyrid-3-  
ylmethylamine, N- (3- (2-chlorophenoxy) phenyl) -N-  
(ethylsulfonyl)pyrid-3-ylmethylamine, N- (3- (2-  
methylphenoxy) phenyl) -N- (ethylsulfonyl)pyrid-3-  
10 ylmethylamine, N- (3- (phenoxy) phenyl) -N-  
(ethylsulfonyl)pyrid-3-ylmethylamine, N- (3- (4-  
carbonylmethoxyphenoxy) phenyl) -N- (ethylsulfonyl)pyrid-  
3-ylmethylamine, N- (3- (3-t-butylphenoxy) phenyl) -N-  
(ethylsulfonyl)pyrid-3-ylmethylamine, N- (3- (4-fluoro-2-  
15 (3-hydroxy-3-methylbutyl) phenoxy) phenyl) -N-  
(ethylsulfonyl)pyrid-3-ylmethylamine, N- (3- (2-fluoro-4-  
t-butylphenoxy) phenyl) -N- (ethylsulfonyl)pyrid-3-  
ylmethylamine, N- (3- (4-  
Cabonylmethoxymethylphenoxy) phenyl) -N-  
20 (ethylsulfonyl)pyrid-3-ylmethylamine, N- (3- (3-  
Amidophenoxy) phenyl) -N- (ethylsulfonyl)pyrid-3-  
ylmethylamine, N- (3- (2,4-difluorophenoxy) phenyl) -N-  
(ethylsulfonyl)pyrid-3-ylmethylamine, N- (3- (2,4,6-  
trifluorophenoxy) phenyl) -N- (ethylsulfonyl)pyrid-3-  
25 ylmethylamine, N- (3- (2- (prop-2-yl) phenoxy) phenyl) -N-  
(ethylsulfonyl)pyrid-3-ylmethylamine, N- (3- (2-  
ethylphenoxy) phenyl) -N- (ethylsulfonyl)pyrid-3-  
ylmethylamine, N- (3- (4-Sulfonylamidophenoxy) phenyl) -N-  
(ethylsulfonyl)pyrid-3-ylmethylamine, N- (3-  
30 (phenoxy) phenyl) -N- (ethylsulfonyl)pyrid-3-  
ylmethylamine, N- (4- (3-Methoxybenzyloxy) phenyl) -N-  
(ethanesulfonyl)pyrid-3-ylmethylamine, N- (3- (2-  
Fluorophenoxy) phenyl) -N- (2,2,2-  
trifluorethanesulfonyl)pyrid-3-ylmethylamine, N- (3- (2-



-236-

Methylphenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(4-Carboxymethylphenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-Fluoro-4-methylphenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-methoxy-phenylsulfanyl))phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-methoxy-phenylsulfanyl))phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-methoxy-phenoxy))phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-benzyloxy-phenoxy))phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(4-fluoro-2-methoxy-phenoxy))phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(4-fluoro-2-methoxy-phenoxy))phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-bromo -phenoxy))phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2,3-dimethoxy-phenoxy))phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-chloro-phenoxy))phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-chloro-6-methyl-phenoxy))phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-isopropoxy-phenoxy))phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-fluoro-5-(2-methoxy-phenoxy))phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-fluoro-5-(2-methoxy-phenoxy))phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-methoxy-phenoxy))phenyl)-N-(3,3,3-trifluoropropanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-bromo-phenoxy))phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2,3-dimethoxy-phenoxy))phenyl)-N-(2,2,2-

-237-

trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-chloro-phenoxy))phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-chloro-phenoxy))phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-chloro-6-methyl-phenoxy))phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-isopropoxy-phenoxy))phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-(Benzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-(Benzyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-Methoxycarbonylphenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-Methoxycarbonylphenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-(Hydoxymethyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(Hydroxymethyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-( $\alpha$ -Hydroxybenzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-( $\alpha$ -hydroxybenzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-( $\alpha$ -hydroxy-2-methylsulfanylbenzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-( $\alpha$ -hydroxy-4-methylsulfanylbenzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-( $\alpha$ -hydroxy-2-methoxybenzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-( $\alpha$ -hydroxy-2-methoxybenzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-( $\alpha$ -hydroxy-3,4-difluorobenzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-( $\alpha$ -

-238-

hydroxy-3,4-difluorobenzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-( $\alpha$ -hydroxy-2-phenylbenzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-( $\alpha$ -hydroxy-2-phenylbenzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-(1-hydroxy-1-(5-methylpyrid-2-yl)methyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-( $\alpha$ -hydroxy-2-trifluoromethylbenzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-( $\alpha$ -hydroxy-2-fluorobenzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-( $\alpha$ -hydroxy-2-fluorobenzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-( $\alpha$ -hydroxy-3-fluorobenzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(Benzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-methylsulfonyl benzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(4-methylsulfonylbenzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-(4-methoxybenzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(4-methoxybenzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-(3,4-difluorobenzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(3,4-difluorobenzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-phenylbenzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-phenylbenzyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-

-239-

Methoxycarbonylphenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(Hydroxymethyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-(Hydroxymethyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(Phenoxymethyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-(Phenoxymethyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-( $\alpha$ -hydroxybenzyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-( $\alpha$ -hydroxybenzyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-( $\alpha$ -hydroxy-2-fluorobenzyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-( $\alpha$ -Hydroxy-3-ethoxycarbonylbenzyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(4-ethoxycarbonylbenzyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-fluorobenzyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-( $\alpha$ -Fluorobenzyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-( $\alpha$ -Fluorobenzyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-Benzoylphenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-Benzoylphenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(1-Phenylvinyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-(1-Phenylvinyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-( $\alpha$ -Methylbenzyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-( $\alpha$ -Methylbenzyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-Iodo-4-methylphenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-( $\alpha$ -

-240-

Hydroxybenzyl)-4-methylphenyl)-N-ethanesulfonyl)pyrid-  
3-ylmethylaniline, N-(3-Benzyl-4-methylphenyl)-N-  
(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(4-t-  
butylphenoxy)-4-methylphenyl)-N-(ethanesulfonyl)pyrid-  
5 3-ylmethylaniline, N-(3-(2-Methoxyphenoxy)benzyl)-N-  
(2,2,2-trifluoroethanesulfonyl)pyrid-3-  
ylmethylaniline, N-(3-(2-Methoxyphenoxy)benzyl)-N-  
(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-  
Phenylvinyl)phenyl)-N-(ethanesulfonyl)pyrid-3-  
10 ylmethylaniline, N-(3-(2-Phenylethyl)phenyl)-N-  
(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(Prop-1-en-  
1-yl)phenyl)-N-(ethanesulfonyl)pyrid-3-  
ylmethylaniline, N-(3-(2-N,N-Diethylamidovinyl)phenyl)-N-  
(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-  
15 Ethoxycarbonylvinyl)phenyl)-N-(ethanesulfonyl)pyrid-3-  
ylmethylaniline, N-(3-Propylphenyl)-N-  
(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-  
Trifluoromethoxyphenyl)-N-(2,2,2-  
trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, , N-(3-  
20 trifluoromethoxyphenyl)-N-(ethanesulfonyl)pyrid-3-  
ylmethylaniline, N-(4-trifluoromethylphenyl)-N-(2,2,2-  
trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-  
trifluoromethoxyphenyl)-N-(ethanesulfonyl)pyrid-3-  
ylmethylaniline, N-(3-methoxyphenyl)-N-(2,2,2-  
25 trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-  
(prop-2-yl)phenyl)-N-(2,2,2-  
trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(2-  
trifluoromethoxyphenyl)-N-(2,2,2-  
trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-  
30 (prop-2-yloxy)phenyl)-N-(2,2,2-  
trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-  
Benzyl-4-chlorophenyl)-N-(2,2,2-  
trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-(3-  
Cyclobutyloxyphenyl)-N-(2,2,2-

-241-

trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-isobutyloxyphenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-sec-butoxyphenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-cyclopentyloxyphenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-cyclohexyloxyphenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(1-ethoxycarbonyl-1,1-dimethyl, methoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(1-ethoxycarbonyl-1-methylmethoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(1-(ethoxycarbonyl)-cyclobut-1-yloxyphenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-pent-3-yloxyphenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-pent-2-yloxyphenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2,2,2-trifluoroethoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-methyl-butyloxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(ethoxycarbonyl, methoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-Fluoro-4-cyanophenoxy)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-Fluoro-4-amidophenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-Fluoro-4-cyanophenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2,4-Difluoro-4-cyanophenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylaniline, N-(4-(2-Fluoro-4-cyanophenoxy)phenyl)-N-(ethanesulfonyl)pyrid-

-242-

3-ylmethylanine, N-(4-(2-Fluoro-4-amidophenoxy)phenyl)-  
N-(ethanesulfonyl)pyrid-3-ylmethylanine, N-(3-  
(Bromomethyl)phenyl)-N-(ethanesulfonyl)pyrid-3-  
ylmethylanine, N-(3-(2-Methoxybenzyl)phenyl)-N-  
5 (ethanesulfonyl)pyrid-3-ylmethylanine, N-(3-(2-  
trifluoromethyl, benzyl)phenyl)-N-  
(ethanesulfonyl)pyrid-3-ylmethylanine, N-(3-(4-  
cyanobenzyl)phenyl)-N-(ethanesulfonyl)pyrid-3-  
ylmethylanine, N-(3-(2-chlorobenzyl)phenyl)-N-  
10 (ethanesulfonyl)pyrid-3-ylmethylanine, N-(3-(4-  
fluorobenzyl)phenyl)-N-(ethanesulfonyl)pyrid-3-  
ylmethylanine, N-(3-(thien-2-yl methyl)phenyl)-N-  
(ethanesulfonyl)pyrid-3-ylmethylanine, N-(3-(thien-3-yl  
methyl)phenyl)-N-(ethanesulfonyl)pyrid-3-  
15 ylmethylanine, N-(3-(2-bromobenzyl)phenyl)-N-  
(ethanesulfonyl)pyrid-3-ylmethylanine, N-(3-(5-methyl-  
thiophen-2-yl)methyl phenyl)-N-(ethanesulfonyl)pyrid-3-  
ylmethylanine, N-(3-(2,4-difluorobenzyl)phenyl)-N-  
(ethanesulfonyl)pyrid-3-ylmethylanine, N-(3-(2,4-  
20 dichlorobenzyl)phenyl)-N-(ethanesulfonyl)pyrid-3-  
ylmethylanine, N-(1-(4-Bromophenyl)ethyl)-N-  
(ethanesulfonyl)pyrid-3-ylmethylanine, N-(4-(4-  
Cyanophenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-  
ylmethylanine, N-((2-Methoxy-5-phenyl)phenyl)-N-  
25 (ethanesulfonyl)pyrid-3-ylmethylanine, N-(1-(4-  
Bromophenyl)ethyl)-N-(2,2,2-  
trifluoroethanesulfonyl)pyrid-3-ylmethylanine, N-(3-(4-  
Cyanophenyl)phenyl)-N-(2,2,2-  
trifluoroethylsulfonyl)pyrid-3-ylmethylanine, N-((3-  
30 Iodophenyl)methyl)-N-(ethylsulfonyl)pyrid-3-ylamine, N-  
((3-Iodophenyl)methyl)-N-(ethylsulfonyl)pyrid-3-  
ylmethylanine, N-(1-(4-(2,4-  
Difluorophenyl)phenyl)ethyl)-N-(ethanesulfonyl)pyrid-3-  
ylmethylanine, N-(1-(4-(4-cyanophenyl)phenyl)ethyl)-N-

-243-

(ethanesulfonyl)pyrid-3-ylmethylaniline, N-((3-(2,4-difluorophenyl)methyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(3-(4-cyanophenyl)phenyl)methyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(3-(3-methanesulfonylaminophenyl)phenyl)methyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(3-(4-N-methylamidophenyl)phenyl)methyl)-N-(ethylsulfonyl)pyrid-3-ylmethylaniline, N-(1-(3-(4-cyanophenyl)phenyl)ethyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2,4-difluorophenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(4-N-methylamidophenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-((3,5-bis-trifluoromethyl)phenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-((3-phenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-((4-(2-trifluoroacetylamino), ethyl)phenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(4-acetylaminophenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(4-methanephensulfonyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(4-trifluoromethoxyphenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(pyrid-4-yl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(3-trifluoromethylphenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(4-carboxyphenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(4-(N'-methanesulfonyl-N'-t-butoxycarbonylaminoethyl)phenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-methylphenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-methoxyphenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-trifluoromethylphenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylaniline, N-(3-(2-



-244-

chlorophenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine, N-(3-(2-fluorophenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine, N-(3-(3,5-difluorophenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine, N-(3-(3-acetylaminophenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine, N-((4-Methyl-3-(4-cyanophenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine, N-(3-(4-(2-Methanesulfonylaminoethyl)phenyl)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)pyrid-3-ylmethylamine, N-((3-(4-Amidophenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine, N-(3-(4-(2-Methanesulfonylaminoethyl)phenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine, N-((3-(2-Methyl-4-cyanophenyl)phenyl)-N-(ethanesulfonyl)pyrid-3-ylmethylamine, N-(4-(2,4-Difluorophenoxy)phenyl)-N-(ethanesulfonyl)-2-fluororpyrid-3-ylmethylamine, N-(4-(2,4-Difluorophenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)-2-fluororpyrid-3-ylmethylamine, N-(4-(2,4-Difluorophenoxy)phenyl)-N-(ethanesulfonyl)-6-fluororpyrid-3-ylmethylamine, N-(4-(2,4-Difluorophenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)-6-fluororpyrid-3-ylmethylamine, N-(4-(2,4-Difluorophenoxy)phenyl)-N-(ethanesulfonyl)-2-methylrpyrid-3-ylmethylamine, N-(4-(2,4-Difluorophenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)-2-methylrpyrid-3-ylmethylamine, N-(4-(2,4-Difluorophenoxy)phenyl)-N-(ethanesulfonyl)-6-methylrpyrid-3-ylmethylamine, N-(4-(2,4-Difluorophenoxy)phenyl)-N-(2,2,2-trifluoroethanesulfonyl)-6-methylrpyrid-3-ylmethylamine, and N-(3-(N'-Methyl-N'-(2-chlorophenylsulfonamido)phenyl))-N-(ethanesulfonyl)pyrid-3-ylmethylamine.

-245-

18. A pharmaceutical composition comprising a compound of Claim 1 and a pharmaceutically acceptable diluent.

19. A method of treating neurological and psychiatric disorders associated with glutamate dysfunction, comprising: administering to a patient in need thereof an effective amount of a compound of Claim 1.

20. A method of treating anxiety, comprising: administering to a patient in need thereof an effective amount of a compound of Claim 1.

21. A method of treating migraine, comprising: administering to a patient in need thereof an effective amount of a compound of Claim 1.

22. A method of treating schizophrenia, comprising: administering to a patient in need thereof an effective amount of a compound of Claim 1.

23. A method of treating epilepsy, comprising: administering to a patient in need thereof an effective amount of a compound of Claim 1.

24. The use of a compound of Claim 1 as a medicament.

25. The use of a compound of Claim 1 for the manufacture of a medicament for treating neurological and psychiatric disorders associated with glutamate dysfunction.

26. The use of a compound of Claim 1 for the manufacture of a medicament for treating anxiety.

27. The use of a compound of Claim 1 for the manufacture of a medicament for treating migraine.

28. The use of a compound of Claim 1 for the manufacture of a medicament for treating schizophrenia.

29. The use of a compound of Claim 1 for the manufacture of a medicament for treating epilepsy.

30. A method of treating neurological and psychiatric disorders associated with glutamate dysfunction,

-246-

comprising: administering to a patient in need thereof an effective amount of an mGluR potentiator.

31. A method of treating anxiety, comprising:  
administering to a patient in need thereof an effective  
5 amount of a compound of an mGluR potentiator.

32. A method of treating migraine, comprising:  
administering to a patient in need thereof an effective  
amount of an mGluR potentiator.

33. A method of treating schizophrenia, comprising:  
10 administering to a patient in need thereof an effective  
amount of an mGluR potentiator.

34. A method of treating epilepsy, comprising:  
administering to a patient in need thereof an effective  
amount of an mGluR potentiator.

15 35. A method according to anyone of Claims 30-34  
wherein the mGluR potentiator is a mGluR<sub>2</sub> potentiator.

36. A method according to anyone of Claims 30-34  
wherein the mGluR potentiator is a mGluR<sub>3</sub> potentiator.

**DERWENT-ACC-NO:** 2001-570499

**DERWENT-WEEK:** 200465

*COPYRIGHT 2008 DERWENT INFORMATION LTD*

**TITLE:** Potentiators of metabotropic glutamate receptors  
useful for treating neurological and psychiatric  
disorders associated with glutamate dysfunction e.g.  
anxiety or migraine

**INVENTOR:** BARDA D A; BRITTON T C ; COLEMAN D S ; DRESSMAN B  
A ; FICHTNER M W ; HENRY S S ; HORNBACK W J ;  
JAGDMANN G E ; JAGDMANN G E J ; JOHNSON K W ;  
JOHNSON M P ; LARGE T H ; MONN J A ; SCHOEPP D D ;  
TIZZANO J P

**PATENT-ASSIGNEE:** BARDA D A[BARDI] , BRITTON T C[BRITI] ,  
COLEMAN D S[COLEI] , DRESSMAN B A[DRESI] ,  
FICHTNER M W[FICHI] , HENRY S S[HENRI] ,  
HORNBACK W J[HORNI] , JAGDMANN G E  
[JAGDI] , JOHNSON K W[JOHNI]

**PRIORITY-DATA:** 2000US-180089P (February 3, 2000) , 2000US-  
180047P (February 3, 2000) , 2002US-182961  
(November 20, 2002)

**PATENT-FAMILY:**

<b>PUB-NO</b>	<b>PUB-DATE</b>	<b>LANGUAGE</b>
WO 0156990 A2	August 9, 2001	EN
AU 200134420 A	August 14, 2001	EN
EP 1255735 A2	November 13, 2002	EN
US 20040006114 A1	January 8, 2004	EN
US 6800651 B2	October 5, 2004	EN

**DESIGNATED-STATES:** AE AG AL AM AT AU AZ BA BB BG BR BY BZ  
 CA CH CN CR CU CZ DE DK DM DZ EE ES FI  
 GB GD GE GH GM HR HU ID IL IN IS JP KE KG  
 KP KR KZ LC LK LR LS LT LU LV MA MD MG  
 MK MN MW MX MZ NO NZ PL PT RO RU SD  
 SE SG SI SK SL T J TM TR TT TZ UA UG US  
 UZ VN YU ZA ZW AT BE CH CY DE DK EA ES  
 FI FR GB GH GM GR IE IT KE LS LU MC MW  
 MZ NL OA PT SD SE SL SZ TR TZ UG ZW AL  
 AT BE CH CY DE DK ES FI FR GB GR IE IT LI  
 LT LU LV MC MK NL PT RO SE SI TR

**APPLICATION-DATA:**

<b>PUB-NO</b>	<b>APPL-DESCRIPTOR</b>	<b>APPL-NO</b>	<b>APPL-DATE</b>
WO2001056990A2	N/A	2001WO- US00643	January 22, 2001
AU 200134420A	N/A	2001AU- 034420	January 22, 2001
EP 1255735A2	N/A	2001EP- 906521	January 22, 2001
EP 1255735A2	N/A	2001WO- US00643	January 22, 2001
US20040006114A1	N/A	2001WO- US00643	January 22, 2001
US 6800651B2	N/A	2001WO- US00643	January 22, 2001
US20040006114A1	N/A	2002US- 182961	November 20, 2002
US 6800651B2	Based on	2002US- 182961	November 20, 2002

**INT-CL-CURRENT:**

<b>TYPE</b>	<b>IPC DATE</b>
CIPS	A61P25/06 20060101
CIPS	A61P25/08 20060101

CIPS	A61P25/18 20060101
CIPS	C07D213/30 20060101
CIPS	C07D213/38 20060101
CIPS	C07D213/40 20060101
CIPS	C07D213/42 20060101
CIPS	C07D213/61 20060101
CIPS	C07D213/643 20060101
CIPS	C07D213/76 20060101
CIPS	C07D213/89 20060101
CIPS	C07D405/12 20060101
CIPS	C07D409/12 20060101

**ABSTRACTED-PUB-NO:** WO 0156990 A2

**BASIC-ABSTRACT:**

NOVELTY - Potentiators of metabotropic glutamate receptors (mGluR) (I) are new.

DESCRIPTION - Potentiators of metabotropic glutamate receptors of formula (I) are new:

$R_1 = C(O)R_3, C(O)OR_4$  or  $SO_2R_5$ ;

$R_2 = 1-3$  substituents selected from H, OH, trisubstituted silyloxy, (substituted)alkyl, (substituted)alkenyl, (substituted)alkoxy, (substituted) cycloalkyl, halo, cyano, nitro, (substituted)phenyl, pyridyloxy, (substituted) thiophenoxy, (substituted)phenylsulfinyl, (substituted)phenylsulfonyl, (substituted)benzoyl or (substituted)phenoxy; or

two  $R_2$  taken together on adjacent positions = fused cycloalkyl or methylenedioxy ring;

$R_3, R_4 =$  alkyl or cycloalkyl;

$R_5 =$  alkyl, cycloalkyl or fluorinated alkyl;

R6 = 1-2 substituents selected from H, alkyl, alkoxy, trifluoromethyl, halo, (substituted)phenoxy;

X = CH<sub>2</sub>, CHR<sub>7</sub> or CH<sub>2</sub>CH<sub>2</sub>;

Y = CH<sub>2</sub>, CHR<sub>8</sub>, CH<sub>2</sub>CH<sub>2</sub>, CHR<sub>9</sub>CH<sub>2</sub> or CH<sub>2</sub>CHR<sub>9</sub>; and

R<sub>7</sub>, R<sub>8</sub>, R<sub>9</sub> = lower alkyl.

AN INDEPENDENT CLAIM is also included for a compound comprising (I) and a diluent.

MGluR potentiator receptors.

Test are described but no results are given.

USE - (I) is used as a medicament for treating neurological and psychiatric disorders associated with glutamate dysfunction. (I) is particularly used to treat anxiety, migraine, schizophrenia or epilepsy (all claimed).

ADVANTAGE - None given.

## **EQUIVALENT-ABSTRACTS:**

### ORGANIC CHEMISTRY

Preparation: (I) may be prepared by reacting a compound of formula (a) with an amine of formula (b) and a reducing agent such as sodium cyanoborohydride to give a compound of formula (c). Acylation of compound (c) gives (I).

Preferred Potentiator: The mGluR potentiator is either a MGluR<sub>2</sub> or MGluR<sub>3</sub> potentiator.

Administration is oral, subcutaneous, intramuscular, intravenous, transdermal, intranasal, rectal, ocular, topical, sublingual or buccal. No dosage is given.

### SPECIFIC COMPOUNDS

Several specific compounds are claimed e.g. N-(3-(2-methoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-yl ethylamine of formula (Ia).

A solution of N-(3-(2-methoxyphenoxy)phenyl)pyrid-3-ylmethylamine was dissolved in 2:1 1,2-dichloroethane/pyridine (6 ml) under nitrogen, cooled to 5 degreesC and treated with 2,2,2-trifluoroethylsulfonyl chloride (300 mg). After one hour at 5 degreesC the mixture was allowed to warm to room temperature and stirred for 18 hours. The mixture was treated with anhydrous potassium carbonate (500 mg), stirred for 30 minutes, filtered and the filtrate concentrated in vacuo. The residue was chromatographed on silica gel (eluted with 2:1 dichloromethane/ethylacetate) to afford N-(3-(2-methoxyphenoxy)phenyl)-N-(2,2,2-trifluoroethylsulfonyl)pyrid-3-yl ethylamine (226 mg, 50%) as a viscous pale amber oil.

**TITLE-TERMS:** POTENTIATE GLUTAMATE RECEPTOR USEFUL TREAT  
NEUROLOGICAL PSYCHIATRIC DISORDER  
ASSOCIATE DYSFUNCTION ANXIETY MIGRAINE

**DERWENT-CLASS:** B03 P14

**CPI-CODES:** B07-D04C; B14-F02C; B14-J01B3; B14-J01B4; B14-J07; B14-L01;

**CHEMICAL-CODES:** Chemical Indexing M2 \*01\* Fragmentation Code  
C316 F013 F431 G011 G012 G100 H5 H542 H6  
H685 H8 K0 K3 K353 M1 M121 M141 M210 M211  
M272 M281 M311 M312 M321 M332 M342 M344  
M362 M373 M391 M413 M510 M521 M532 M540  
M710 P411 P442 P446 P448 P527 Specific  
Compounds RA5BCJ Registry Numbers 456757

Chemical Indexing M2 \*02\* Fragmentation Code  
C316 F013 F431 G010 G011 G012 G100 H5 H542  
H6 H685 H8 K0 K3 K353 M1 M121 M141 M280  
M311 M312 M321 M322 M332 M342 M344 M362  
M373 M391 M392 M413 M510 M521 M533 M540  
M710 P411 P442 P446 P448 P527 Specific  
Compounds RA5BCR Registry Numbers 456772

Chemical Indexing M2 \*03\* Fragmentation Code  
C316 F013 F431 G010 G013 G100 H5 H541 H8 K0



K3 K353 M1 M121 M141 M210 M212 M271 M281  
M312 M321 M332 M342 M373 M391 M413 M510  
M521 M532 M540 M710 P411 P442 P446 P448  
P527 Specific Compounds RA5BD3 Registry  
Numbers 456785

Chemical Indexing M2 \*04\* Fragmentation Code  
C316 F013 F431 G010 G013 G100 H5 H541 H8 K0  
K3 K353 M1 M121 M141 M210 M212 M271 M281  
M311 M312 M321 M332 M342 M373 M392 M413  
M510 M521 M532 M540 M710 P411 P442 P446  
P448 P527 Specific Compounds RA5BD4 Registry  
Numbers 456786

Chemical Indexing M2 \*05\* Fragmentation Code  
C316 F013 F431 G010 G013 G100 H5 H541 H8 K0  
K3 K353 M1 M121 M141 M210 M212 M271 M281  
M311 M322 M342 M373 M392 M413 M510 M521  
M532 M540 M710 P411 P442 P446 P448 P527  
Specific Compounds RA5BD5 Registry Numbers  
456787

Chemical Indexing M2 \*06\* Fragmentation Code  
C316 F013 F431 G010 G013 G100 H5 H541 H8 K0  
K3 K353 M1 M121 M141 M210 M213 M232 M271  
M281 M311 M321 M342 M373 M391 M413 M510  
M521 M532 M540 M710 P411 P442 P446 P448  
P527 Specific Compounds RA5BD6 Registry  
Numbers 456788

Chemical Indexing M2 \*07\* Fragmentation Code  
C316 F013 F431 G010 G011 G100 H5 H541 H6  
H685 H8 K0 K3 K353 M1 M121 M141 M280 M311  
M312 M321 M332 M342 M344 M362 M373 M391  
M413 M510 M521 M532 M540 M710 P411 P442  
P446 P448 P527 Specific Compounds RA5BD8  
Registry Numbers 456790

Chemical Indexing M2 \*08\* Fragmentation Code  
C316 F013 F431 G010 G013 G100 H5 H541 H8 K0  
K3 K353 M1 M121 M141 M210 M212 M271 M281

M311 M321 M342 M373 M391 M413 M510 M521  
M532 M540 M710 P411 P442 P446 P448 P527  
Specific Compounds RA5BD9 Registry Numbers  
456791

Chemical Indexing M2 \*09\* Fragmentation Code H5  
H542 H6 H602 H641 H685 H8 K0 K3 K353 M1 M121  
M141 M210 M211 M272 M281 M311 M312 M321  
M332 M342 M344 M362 M373 M391 M413 M510  
M521 M532 M540 M710 P411 P442 P446 P448  
P527 Specific Compounds RA5BDJ Registry  
Numbers 456801

Chemical Indexing M2 \*10\* Fragmentation Code  
C316 F013 F431 G014 G015 G100 H5 H543 H6  
H602 H641 H685 H8 K0 K3 K353 M1 M121 M141  
M210 M211 M272 M282 M311 M312 M321 M332  
M342 M344 M362 M373 M391 M413 M510 M521  
M532 M540 M710 P411 P442 P446 P448 P527  
Specific Compounds RA5BDL Registry Numbers  
456803

Chemical Indexing M2 \*11\* Fragmentation Code  
C316 F013 F431 G010 G011 G013 G100 H5 H542  
H6 H685 H8 K0 K3 K353 M1 M121 M141 M280  
M311 M312 M321 M322 M332 M342 M344 M362  
M373 M391 M392 M413 M510 M521 M533 M540  
M710 P411 P442 P446 P448 P527 Specific  
Compounds RA5BDM Registry Numbers 456804

Chemical Indexing M2 \*12\* Fragmentation Code  
C316 F013 F431 G011 G013 G100 H5 H542 H8 K0  
K3 K353 M1 M121 M141 M210 M211 M212 M271  
M272 M281 M311 M321 M342 M373 M391 M413  
M510 M521 M532 M540 M710 P411 P442 P446  
P448 P527 Specific Compounds RA5BDO Registry  
Numbers 456806

Chemical Indexing M2 \*13\* Fragmentation Code  
C316 F013 F431 G011 G013 G030 G112 G530 H5  
H542 H8 K0 K3 K353 M1 M121 M123 M141 M147

M210 M211 M272 M281 M311 M321 M342 M373  
M391 M413 M510 M521 M532 M541 M710 P411  
P442 P446 P448 P527 Specific Compounds  
RA5BDQ Registry Numbers 456808

Chemical Indexing M2 \*14\* Fragmentation Code  
C316 F013 F431 G011 G015 G100 H5 H542 H6  
H602 H641 H8 K0 K3 K353 M1 M121 M141 M210  
M211 M212 M271 M272 M281 M311 M321 M342  
M373 M391 M413 M510 M521 M532 M540 M710  
P411 P442 P446 P448 P527 Specific Compounds  
RA5BDU Registry Numbers 456812

Chemical Indexing M2 \*15\* Fragmentation Code  
C316 F013 F431 G011 G015 G030 G112 G530 H5  
H542 H6 H602 H641 H8 K0 K3 K353 M1 M121 M123  
M141 M147 M210 M211 M272 M281 M311 M321  
M342 M373 M391 M413 M510 M521 M532 M541  
M710 P411 P442 P446 P448 P527 Specific  
Compounds RA5BDY Registry Numbers 456816

Chemical Indexing M2 \*16\* Fragmentation Code  
C316 F013 F431 G011 G013 G100 H5 H542 H6  
H685 H8 K0 K3 K353 M1 M121 M141 M210 M211  
M272 M281 M311 M312 M321 M332 M342 M344  
M362 M373 M391 M413 M510 M521 M532 M540  
M710 P411 P442 P446 P448 P527 Specific  
Compounds RA5BE3 Registry Numbers 456821

Chemical Indexing M2 \*17\* Fragmentation Code  
C316 F013 F431 G011 G013 G100 H5 H542 H6  
H685 H8 K0 K3 K353 M1 M121 M141 M210 M212  
M272 M281 M311 M312 M321 M332 M342 M344  
M362 M373 M391 M413 M510 M521 M532 M540  
M710 P411 P442 P446 P448 P527 Specific  
Compounds RA5BE6 Registry Numbers 456824

Chemical Indexing M2 \*18\* Fragmentation Code  
C316 F013 F431 G011 G013 G100 H5 H542 H8 K0  
K3 K353 M1 M121 M141 M210 M212 M271 M272  
M281 M311 M321 M342 M373 M391 M413 M510

M521 M532 M540 M710 P411 P442 P446 P448  
P527 Specific Compounds RA5BEB Registry  
Numbers 456829

Chemical Indexing M2 \*19\* Fragmentation Code  
C316 F013 F431 G011 G013 G030 G112 G530 H5  
H542 H8 K0 K3 K353 M1 M121 M123 M141 M147  
M210 M212 M272 M281 M311 M321 M342 M373  
M391 M413 M510 M521 M532 M541 M710 P411  
P442 P446 P448 P527 Specific Compounds  
RA5BEK Registry Numbers 456838

Chemical Indexing M2 \*20\* Fragmentation Code  
C316 F013 F431 G011 G012 G030 G112 G530 H5  
H542 H8 K0 K3 K353 M1 M121 M123 M141 M147  
M210 M211 M272 M281 M311 M321 M342 M373  
M391 M413 M510 M521 M532 M541 M710 P411  
P442 P446 P448 P527 Specific Compounds  
RA5BEL Registry Numbers 456839

Chemical Indexing M2 \*21\* Fragmentation Code  
C316 F013 F431 G011 G012 G100 H5 H542 H8 K0  
K3 K353 M1 M121 M141 M210 M211 M212 M271  
M272 M281 M311 M321 M342 M373 M391 M413  
M510 M521 M532 M540 M710 P411 P442 P446  
P448 P527 Specific Compounds RA5BEW Registry  
Numbers 456850

Chemical Indexing M2 \*22\* Fragmentation Code  
C316 F013 F431 G010 G013 G100 H5 H541 H6  
H685 H8 K0 K3 K353 M1 M121 M141 M280 M311  
M312 M321 M332 M342 M344 M362 M373 M391  
M413 M510 M521 M532 M540 M710 P411 P442  
P446 P448 P527 Specific Compounds RA5BF3  
Registry Numbers 456857

Chemical Indexing M2 \*23\* Fragmentation Code  
C316 F013 F431 G011 G012 G100 H4 H401 H441  
H5 H541 H6 H685 H8 K0 K3 K353 M1 M121 M141  
M280 M311 M312 M321 M332 M342 M344 M362  
M373 M391 M413 M510 M521 M532 M540 M710

P411 P442 P446 P448 P527 Specific Compounds  
RA5BF5 Registry Numbers 456859

Chemical Indexing M2 \*24\* Fragmentation Code  
C316 F013 F431 G011 G013 G100 H4 H401 H441  
H5 H541 H6 H685 H8 K0 K3 K353 M1 M121 M141  
M280 M311 M312 M321 M332 M342 M344 M362  
M373 M391 M413 M510 M521 M532 M540 M710  
P411 P442 P446 P448 P527 Specific Compounds  
RA5BF6 Registry Numbers 456860

Chemical Indexing M2 \*25\* Fragmentation Code  
M413 M510 M521 M533 M540 M710 P411 P442  
P446 P448 P527 Specific Compounds RA5BF7  
Registry Numbers 456861

Chemical Indexing M2 \*26\* Fragmentation Code  
C316 F013 F431 G013 G015 G100 H5 H541 H6  
H601 H608 H642 H685 H8 K0 K3 K353 M1 M121  
M141 M280 M311 M312 M321 M332 M342 M344  
M362 M373 M391 M413 M510 M521 M532 M540  
M710 P411 P442 P446 P448 P527 Specific  
Compounds RA5BF8 Registry Numbers 456862

Chemical Indexing M2 \*27\* Fragmentation Code  
M344 M362 M373 M391 M413 M510 M521 M532  
M541 M710 P411 P442 P446 P448 P527 Specific  
Compounds RA5BFB Registry Numbers 456865

Chemical Indexing M2 \*28\* Fragmentation Code  
C316 F013 F431 G010 G013 G019 G100 H5 H541  
H6 H685 H8 K0 K3 K353 M1 M121 M129 M132  
M141 M150 M280 M311 M312 M321 M322 M332  
M342 M344 M362 M373 M391 M413 M510 M521  
M533 M540 M710 P411 P442 P446 P448 P527  
Specific Compounds RA5BGJ Registry Numbers  
456910

Chemical Indexing M2 \*29\* Fragmentation Code  
C316 F013 F019 F211 F431 G013 G019 G100 H5  
H521 H541 H6 H685 H8 K0 K3 K353 M1 M121 M123

M141 M149 M280 M311 M312 M321 M332 M342  
M344 M362 M373 M391 M413 M510 M522 M532  
M540 M710 P411 P442 P446 P448 P527 Specific  
Compounds RA5BGK Registry Numbers 456911

Chemical Indexing M2 \*30\* Fragmentation Code  
C316 F013 F019 F211 F431 G013 G019 G100 H5  
H521 H541 H6 H685 H8 K0 K3 K353 M1 M121 M123  
M141 M149 M280 M311 M322 M342 M344 M362  
M373 M391 M413 M510 M522 M532 M540 M710  
P411 P442 P446 P448 P527 Specific Compounds  
RA5BGL Registry Numbers 456912

Chemical Indexing M2 \*31\* Fragmentation Code  
C316 F013 F431 G013 G019 G030 G112 G563 H5  
H541 H6 H685 H8 K0 K3 K353 M1 M113 M121  
M141 M280 M311 M322 M342 M344 M362 M373  
M391 M413 M510 M521 M532 M541 M710 P411  
P442 P446 P448 P527 Specific Compounds  
RA5BGN Registry Numbers 456914

Chemical Indexing M2 \*32\* Fragmentation Code  
C316 F013 F431 G010 G013 G019 G100 H5 H541  
H6 H685 H8 K0 K3 K353 M1 M121 M129 M132  
M141 M150 M280 M311 M323 M342 M344 M362  
M373 M391 M413 M510 M521 M533 M540 M710  
P411 P442 P446 P448 P527 Specific Compounds  
RA5BGO Registry Numbers 456915

Chemical Indexing M2 \*33\* Fragmentation Code  
C316 F013 F431 G010 G013 G019 G100 H5 H542  
H6 H685 H8 K0 K3 K353 M1 M121 M141 M280  
M311 M323 M342 M344 M362 M373 M391 M392  
M413 M510 M521 M533 M540 M710 P411 P442  
P446 P448 P527 Specific Compounds RA5BGP  
Registry Numbers 456916

Chemical Indexing M2 \*34\* Fragmentation Code  
C316 F013 F431 G010 G013 G019 G100 H5 H542  
H8 K0 K3 K353 M1 M121 M141 M210 M213 M231  
M271 M281 M311 M322 M342 M373 M392 M413

M510 M521 M533 M540 M710 P411 P442 P446  
P448 P527 Specific Compounds RA5BGQ Registry  
Numbers 456917

Chemical Indexing M2 \*35\* Fragmentation Code  
C316 F013 F431 G013 G019 G030 G112 G563 H5  
H541 H8 K0 K3 K353 M1 M113 M121 M141 M210  
M213 M231 M271 M281 M311 M321 M342 M373  
M391 M413 M510 M521 M532 M541 M710 P411  
P442 P446 P448 P527 Specific Compounds  
RA5BGR Registry Numbers 456918

Chemical Indexing M2 \*36\* Fragmentation Code  
C316 F013 F431 G010 G013 G019 G100 H5 H541  
H8 K0 K3 K353 M1 M121 M129 M132 M141 M150  
M210 M213 M231 M271 M281 M311 M322 M342  
M373 M391 M413 M510 M521 M533 M540 M710  
P411 P442 P446 P448 P527 Specific Compounds  
RA5BGS Registry Numbers 456919

Chemical Indexing M2 \*37\* Fragmentation Code  
C316 F012 F431 G013 G019 G030 G112 G563 H5  
H541 H8 K0 K3 K353 M1 M113 M121 M141 M210  
M212 M271 M281 M311 M321 M342 M373 M391  
M413 M510 M521 M532 M541 M710 P411 P442  
P446 P448 P527 Specific Compounds RA5BGY  
Registry Numbers 456925

Chemical Indexing M2 \*38\* Fragmentation Code M1  
M121 M141 M210 M212 M271 M281 M311 M322  
M342 M373 M392 M413 M510 M521 M533 M540  
M710 P411 P442 P446 P448 P527 Specific  
Compounds RA5BH0 Registry Numbers 456927

Chemical Indexing M2 \*39\* Fragmentation Code  
C316 F012 F431 G010 G013 G019 G100 H5 H541  
H8 K0 K3 K353 M1 M121 M129 M132 M141 M150  
M210 M212 M271 M281 M311 M322 M342 M373  
M391 M413 M510 M521 M533 M540 M710 P411  
P442 P446 P448 P527 Specific Compounds  
RA5BH2 Registry Numbers 456929

Chemical Indexing M2 \*40\* Fragmentation Code  
C316 F012 F013 F211 F431 G013 G019 G100 H5  
H521 H541 H8 K0 K3 K353 M1 M121 M123 M141  
M149 M210 M212 M271 M281 M311 M321 M342  
M373 M391 M413 M510 M522 M532 M540 M710  
P411 P442 P446 P448 P527 Specific Compounds  
RA5BH3 Registry Numbers 456930

Chemical Indexing M2 \*41\* Fragmentation Code  
C316 F012 F431 G010 G013 G019 G100 H5 H542  
H6 H685 H8 K0 K3 K353 M1 M121 M141 M280  
M311 M312 M321 M322 M332 M342 M344 M362  
M373 M391 M392 M413 M510 M521 M533 M540  
M710 P411 P442 P446 P448 P527 Specific  
Compounds RA5BH4 Registry Numbers 456931

Chemical Indexing M2 \*42\* Fragmentation Code  
C316 F012 F431 G013 G019 G030 G112 G563 H5  
H541 H6 H685 H8 K0 K3 K353 M1 M113 M121  
M141 M280 M311 M312 M321 M332 M342 M344  
M362 M373 M391 M413 M510 M521 M532 M541  
M710 P411 P442 P446 P448 P527 Specific  
Compounds RA5BH5 Registry Numbers 456932

Chemical Indexing M2 \*43\* Fragmentation Code  
C316 F012 F431 G010 G013 G019 G100 H5 H541  
H6 H685 H8 K0 K3 K353 M1 M121 M129 M132  
M141 M150 M280 M311 M312 M321 M322 M332  
M342 M344 M362 M373 M391 M413 M510 M521  
M533 M540 M710 P411 P442 P446 P448 P527  
Specific Compounds RA5BH6 Registry Numbers  
456933

Chemical Indexing M2 \*44\* Fragmentation Code  
C316 F012 F013 F211 F431 G013 G019 G100 H5  
H521 H541 H6 H685 H8 K0 K3 K353 M1 M121 M123  
M141 M149 M280 M311 M312 M321 M332 M342  
M344 M362 M373 M391 M413 M510 M522 M532  
M540 M710 P411 P442 P446 P448 P527 Specific  
Compounds RA5BHA Registry Numbers 456937



Chemical Indexing M2 \*45\* Fragmentation Code  
 F013 F431 G013 G030 G111 G530 H5 H541 H8 J0  
 J011 J3 J341 M1 M123 M136 M210 M215 M231  
 M272 M281 M311 M321 M342 M373 M391 M413  
 M510 M521 M531 M541 M710 P411 P442 P446  
 P448 P527 Specific Compounds RA5BHD Registry  
 Numbers 456940

Chemical Indexing M2 \*46\* Fragmentation Code  
 D022 D140 F013 F431 G013 G030 G111 G530 H5  
 H541 H8 J0 J011 J3 J341 M1 M122 M123 M136  
 M141 M280 M311 M321 M342 M373 M391 M412  
 M511 M521 M531 M541 M710 P411 P442 P446  
 P448 P527 Specific Compounds RA5BHF Registry  
 Numbers 456942

Chemical Indexing M2 \*47\* Fragmentation Code  
 F013 F431 G013 G030 G111 G530 H5 H541 H8 J0  
 J011 J3 J341 M1 M123 M136 M210 M211 M272  
 M281 M311 M321 M342 M373 M391 M413 M510  
 M521 M531 M541 M710 P411 P442 P446 P448  
 P527 Specific Compounds RA5BHI Registry  
 Numbers 456945

Chemical Indexing M2 \*48\* Fragmentation Code  
 F013 F431 G013 G021 G030 G111 G223 G530 H5  
 H541 H8 J0 J011 J3 J341 M1 M122 M123 M136  
 M141 M280 M311 M321 M342 M373 M391 M413  
 M510 M521 M532 M541 M710 P411 P442 P446  
 P448 P527 Specific Compounds RA5BHJ Registry  
 Numbers 456946

Chemical Indexing M2 \*49\* Fragmentation Code  
 F013 F431 G013 G030 G039 G111 G530 G563 H5  
 H541 H8 J0 J011 J3 J341 M1 M123 M129 M136  
 M141 M280 M311 M321 M342 M373 M391 M413  
 M510 M521 M531 M542 M710 P411 P442 P446  
 P448 P527 Specific Compounds RA5BJL Registry  
 Numbers 457019

Chemical Indexing M2 \*50\* Fragmentation Code  
 F012 F013 F431 F499 G013 G019 G030 G112 G530  
 H5 H521 H541 H8 J0 J011 J3 J341 M1 M121 M123  
 M129 M136 M141 M149 M280 M311 M321 M342  
 M373 M391 M413 M510 M522 M532 M541 M710  
 P411 P442 P446 P448 P527 Specific Compounds  
 RA5BJS Registry Numbers 457026

Chemical Indexing M2 \*51\* Fragmentation Code  
 C316 F013 F431 G013 G100 H5 H541 H8 K0 K3  
 K353 M210 M211 M215 M231 M271 M272 M281  
 M311 M321 M342 M373 M391 M413 M510 M521  
 M531 M540 M710 P411 P442 P446 P448 P527  
 Specific Compounds RA5BJV Registry Numbers  
 457029

Chemical Indexing M2 \*52\* Fragmentation Code  
 C316 D022 D140 F013 F431 G013 G100 H5 H541  
 H8 K0 K3 K353 M1 M122 M141 M210 M211 M271  
 M281 M311 M321 M342 M373 M391 M412 M511  
 M521 M531 M540 M710 P411 P442 P446 P448  
 P527 Specific Compounds RA5BJX Registry  
 Numbers 457031

Chemical Indexing M2 \*53\* Fragmentation Code  
 C316 F013 F431 G013 G100 H5 H541 H8 K0 K3  
 K353 M210 M211 M271 M272 M281 M311 M321  
 M342 M373 M391 M413 M510 M521 M531 M540  
 M710 P411 P442 P446 P448 P527 Specific  
 Compounds RA5BJZ Registry Numbers 457033

Chemical Indexing M2 \*54\* Fragmentation Code  
 C316 F013 F431 G013 G021 G111 G223 H5 H541  
 H8 K0 K3 K353 M1 M122 M141 M210 M211 M271  
 M281 M311 M321 M342 M373 M391 M413 M510  
 M521 M532 M540 M710 P411 P442 P446 P448  
 P527 Specific Compounds RA5BK2 Registry  
 Numbers 457036

Chemical Indexing M2 \*55\* Fragmentation Code  
 C316 F013 F431 G013 G030 G111 G563 H5 H541

H8 K0 K3 K353 M1 M123 M141 M210 M211 M271  
M281 M311 M321 M342 M373 M391 M413 M510  
M521 M531 M541 M710 P411 P442 P446 P448  
P527 Specific Compounds RA5BK4 Registry  
Numbers 457038

Chemical Indexing M2 \*56\* Fragmentation Code  
C316 F012 F013 F431 F499 G013 G019 G100 H5  
H521 H541 H8 K0 K3 K353 M1 M121 M123 M141  
M149 M210 M211 M271 M281 M311 M321 M342  
M373 M391 M413 M510 M522 M532 M540 M710  
P411 P442 P446 P448 P527 Specific Compounds  
RA5BK6 Registry Numbers 457040

Chemical Indexing M2 \*57\* Fragmentation Code  
C316 F013 F431 G013 G100 H5 H541 H8 K0 K3  
K353 M210 M212 M215 M231 M271 M272 M281  
M311 M321 M342 M373 M391 M413 M510 M521  
M531 M540 M710 P411 P442 P446 P448 P527  
Specific Compounds RA5BK8 Registry Numbers  
457042

Chemical Indexing M2 \*58\* Fragmentation Code  
C316 D022 D140 F013 F431 G013 G100 H5 H541  
H8 K0 K3 K353 M1 M122 M141 M210 M212 M271  
M281 M311 M321 M342 M373 M391 M412 M511  
M521 M531 M540 M710 P411 P442 P446 P448  
P527 Specific Compounds RA5BKA Registry  
Numbers 457044

Chemical Indexing M2 \*59\* Fragmentation Code  
C316 F013 F431 G013 G100 H5 H541 H8 K0 K3  
K353 M210 M211 M212 M271 M272 M281 M311  
M321 M342 M373 M391 M413 M510 M521 M531  
M540 M710 P411 P442 P446 P448 P527 Specific  
Compounds RA5BKD Registry Numbers 457047

Chemical Indexing M2 \*60\* Fragmentation Code  
C316 F013 F431 G013 G021 G111 G223 H5 H541  
H8 K0 K3 K353 M1 M122 M141 M210 M212 M271  
M281 M311 M321 M342 M373 M391 M413 M510

M521 M532 M540 M710 P411 P442 P446 P448  
P527 Specific Compounds RA5BL6 Registry  
Numbers 457076

Chemical Indexing M2 \*61\* Fragmentation Code  
C316 F013 F431 G013 G030 G111 G563 H5 H541  
H8 K0 K3 K353 M1 M123 M141 M210 M212 M271  
M281 M311 M321 M342 M373 M391 M413 M510  
M521 M531 M541 M710 P411 P442 P446 P448  
P527 Specific Compounds RA5BL8 Registry  
Numbers 457078

Chemical Indexing M2 \*62\* Fragmentation Code  
C316 F012 F013 F431 F499 G013 G019 G100 H5  
H521 H541 H8 K0 K3 K353 M1 M121 M123 M141  
M149 M210 M212 M271 M281 M311 M321 M342  
M373 M391 M413 M510 M522 M532 M540 M710  
P411 P442 P446 P448 P527 Specific Compounds  
RA5BLB Registry Numbers 457081

Chemical Indexing M2 \*63\* Fragmentation Code  
C316 F013 F431 G013 G100 H5 H541 H6 H685 H8  
K0 K3 K353 M210 M215 M231 M272 M281 M311  
M312 M321 M332 M342 M344 M362 M373 M391  
M413 M510 M521 M531 M540 M710 P411 P442  
P446 P448 P527 Specific Compounds RA5BLC  
Registry Numbers 457082

Chemical Indexing M2 \*64\* Fragmentation Code  
C316 D022 D140 F013 F431 G013 G100 H5 H541  
H6 H685 H8 K0 K3 K353 M1 M122 M141 M280  
M311 M312 M321 M332 M342 M344 M362 M373  
M391 M412 M511 M521 M531 M540 M710 P411  
P442 P446 P448 P527 Specific Compounds  
RA5BLE Registry Numbers 457084

Chemical Indexing M2 \*65\* Fragmentation Code  
C316 F013 F431 G013 G100 H5 H541 H6 H685 H8  
K0 K3 K353 M210 M211 M272 M281 M311 M312  
M321 M332 M342 M344 M362 M373 M391 M413  
M510 M521 M531 M540 M710 P411 P442 P446

# P448 P527 Specific Compounds RA5BLH Registry Numbers 457087

Chemical Indexing M2 \*66\* Fragmentation Code  
C316 F013 F431 G013 G021 G111 G223 H5 H541  
H6 H685 H8 K0 K3 K353 M1 M122 M141 M280  
M311 M312 M321 M332 M342 M344 M362 M373  
M391 M413 M510 M521 M532 M540 M710 P411  
P442 P446 P448 P527 Specific Compounds  
RA5BLP Registry Numbers 457095

Chemical Indexing M2 \*67\* Fragmentation Code  
C316 F013 F431 G013 G030 G111 G563 H5 H541  
H6 H685 H8 K0 K3 K353 M1 M123 M141 M280  
M311 M312 M321 M332 M342 M344 M362 M373  
M391 M413 M510 M521 M531 M541 M710 P411  
P442 P446 P448 P527 Specific Compounds  
RA5BLR Registry Numbers 457097

Chemical Indexing M2 \*68\* Fragmentation Code  
C316 F012 F013 F431 F499 G013 G019 G100 H5  
H521 H541 H6 H685 H8 K0 K3 K353 M1 M121 M123  
M141 M149 M280 M311 M312 M321 M332 M342  
M344 M362 M373 M391 M413 M510 M522 M532  
M540 M710 P411 P442 P446 P448 P527 Specific  
Compounds RA5BLT Registry Numbers 457099

Chemical Indexing M2 \*69\* Fragmentation Code  
C316 F013 F431 G013 G100 H5 H541 H6 H685 H8  
K0 K3 K353 M210 M215 M231 M272 M281 M311  
M322 M342 M344 M362 M373 M391 M413 M510  
M521 M531 M540 M710 P411 P442 P446 P448  
P527 Specific Compounds RA5BLW Registry  
Numbers 457102

Chemical Indexing M2 \*70\* Fragmentation Code  
C316 D022 D140 F013 F431 G013 G100 H5 H541  
H6 H685 H8 K0 K3 K353 M1 M122 M141 M280  
M311 M322 M342 M344 M362 M373 M391 M412  
M511 M521 M531 M540 M710 P411 P442 P446  
P448 P527 Specific Compounds RA5BRM Registry

## Numbers 457307

Chemical Indexing M2 \*71\* Fragmentation Code  
C316 F013 F431 G013 G100 H5 H541 H6 H685 H8  
K0 K3 K353 M210 M211 M272 M281 M311 M322  
M342 M344 M362 M373 M391 M413 M510 M521  
M531 M540 M710 P411 P442 P446 P448 P527  
Specific Compounds RA5BRN Registry Numbers  
457308

Chemical Indexing M2 \*72\* Fragmentation Code  
H541 H6 H685 H8 K0 K3 K353 M1 M122 M141  
M280 M311 M322 M342 M344 M362 M373 M391  
M413 M510 M521 M532 M540 M710 P411 P442  
P446 P448 P527 Specific Compounds RA5BRO  
Registry Numbers 457309

Chemical Indexing M2 \*73\* Fragmentation Code  
C316 F013 F431 G013 G030 G111 G563 H5 H541  
H6 H685 H8 K0 K3 K353 M1 M123 M141 M280  
M311 M322 M342 M344 M362 M373 M391 M413  
M510 M521 M531 M541 M710 P411 P442 P446  
P448 P527 Specific Compounds RA5BRP Registry  
Numbers 457310

Chemical Indexing M2 \*74\* Fragmentation Code  
M141 M149 M280 M311 M322 M342 M344 M362  
M373 M391 M413 M510 M522 M532 M540 M710  
P411 P442 P446 P448 P527 Specific Compounds  
RA5BSE Registry Numbers 457338

Chemical Indexing M2 \*75\* Fragmentation Code  
C316 F013 F431 G013 G100 H5 H541 H8 K0 K3  
K353 M210 M213 M215 M231 M271 M272 M281  
M311 M321 M342 M373 M391 M413 M510 M521  
M531 M540 M710 P411 P442 P446 P448 P527  
Specific Compounds RA5BSF Registry Numbers  
457339

Chemical Indexing M2 \*76\* Fragmentation Code  
C316 D022 D140 F013 F431 G013 G100 H5 H541

H8 K0 K3 K353 M1 M122 M141 M210 M213 M231  
M271 M281 M311 M321 M342 M373 M391 M412  
M511 M521 M531 M540 M710 P411 P442 P446  
P448 P527 Specific Compounds RA5BSG Registry  
Numbers 457340

Chemical Indexing M2 \*77\* Fragmentation Code  
C316 F013 F431 G013 G021 G111 G223 H5 H541  
H8 K0 K3 K353 M1 M122 M141 M210 M213 M231  
M271 M281 M311 M321 M342 M373 M391 M413  
M510 M521 M532 M540 M710 P411 P442 P446  
P448 P527 Specific Compounds RA5BSQ Registry  
Numbers 457350

Chemical Indexing M2 \*78\* Fragmentation Code  
C316 F013 F431 G013 G030 G111 G563 H5 H541  
H8 K0 K3 K353 M1 M123 M141 M210 M213 M231  
M271 M281 M311 M321 M342 M373 M391 M413  
M510 M521 M531 M541 M710 P411 P442 P446  
P448 P527 Specific Compounds RA5BST Registry  
Numbers 457354

Chemical Indexing M2 \*79\* Fragmentation Code  
C316 F012 F013 F431 F499 G013 G019 G100 H5  
H521 H541 H8 K0 K3 K353 M1 M121 M123 M141  
M149 M210 M213 M231 M271 M281 M311 M321  
M342 M373 M391 M413 M510 M522 M532 M540  
M710 P411 P442 P446 P448 P527 Specific  
Compounds RA5BSU Registry Numbers 457355

Chemical Indexing M2 \*80\* Fragmentation Code  
C316 F012 F431 G013 G100 H5 H541 H8 K0 K3  
K353 M210 M212 M215 M231 M271 M272 M281  
M311 M321 M342 M373 M391 M413 M510 M521  
M531 M540 M710 P411 P442 P446 P448 P527  
Specific Compounds RA5BSW Registry Numbers  
457357

Chemical Indexing M2 \*81\* Fragmentation Code  
C316 D022 D140 F012 F431 G013 G100 H5 H541  
H8 K0 K3 K353 M1 M122 M141 M210 M212 M271

M281 M311 M321 M342 M373 M391 M412 M511  
M521 M531 M540 M710 P411 P442 P446 P448  
P527 Specific Compounds RA5BSX Registry  
Numbers 457359

Chemical Indexing M2 \*82\* Fragmentation Code  
C316 F012 F431 G013 G100 H5 H541 H8 K0 K3  
K353 M210 M211 M212 M271 M272 M281 M311  
M321 M342 M373 M391 M413 M510 M521 M531  
M540 M710 P411 P442 P446 P448 P527 Specific  
Compounds RA5BSZ Registry Numbers 457361

Chemical Indexing M2 \*83\* Fragmentation Code  
C316 F012 F431 G013 G021 G111 G223 H5 H541  
H8 K0 K3 K353 M1 M122 M210 M212 M271 M281  
M311 M321 M342 M373 M391 M413 M510 M521  
M532 M540 M710 P411 P442 P446 P448 P527  
Specific Compounds RA5BT0 Registry Numbers  
457363

Chemical Indexing M2 \*84\* Fragmentation Code  
C316 F012 F431 G013 G030 G111 G563 H5 H541  
H8 K0 K3 K353 M1 M123 M141 M210 M212 M271  
M281 M311 M321 M342 M373 M391 M413 M510  
M521 M531 M541 M710 P411 P442 P446 P448  
P527 Specific Compounds RA5BT5 Registry  
Numbers 457368

Chemical Indexing M2 \*85\* Fragmentation Code  
C316 F012 F019 F431 F499 G013 G019 G100 H5  
H521 H541 H8 K0 K3 K353 M1 M121 M123 M141  
M149 M210 M212 M271 M281 M311 M321 M342  
M373 M391 M413 M510 M522 M532 M540 M710  
P411 P442 P446 P448 P527 Specific Compounds  
RA5BT6 Registry Numbers 457369

Chemical Indexing M2 \*86\* Fragmentation Code  
C316 F012 F431 G013 G100 H5 H541 H6 H685 H8  
K0 K3 K353 M210 M215 M231 M272 M281 M311  
M312 M321 M332 M342 M344 M362 M373 M391  
M413 M510 M521 M531 M540 M710 P411 P442



P446 P448 P527 Specific Compounds RA5BT8  
Registry Numbers 457371

Chemical Indexing M2 \*87\* Fragmentation Code  
C316 D022 D140 F012 F431 G013 G100 H5 H541  
H6 H685 H8 K0 K3 K353 M1 M122 M141 M280  
M311 M312 M321 M332 M342 M344 M362 M373  
M391 M412 M511 M521 M531 M540 M710 P411  
P442 P446 P448 P527 Specific Compounds RA5BT9  
Registry Numbers 457372

Chemical Indexing M2 \*88\* Fragmentation Code  
C316 F012 F431 G013 G100 H5 H541 H6 H685 H8  
K0 K3 K353 M210 M211 M272 M281 M311 M312  
M321 M332 M342 M344 M362 M373 M391 M413  
M510 M521 M531 M540 M710 P411 P442 P446  
P448 P527 Specific Compounds RA5BTA Registry  
Numbers 457373

Chemical Indexing M2 \*89\* Fragmentation Code  
C316 F012 F431 G013 G021 G111 G223 H5 H541  
H6 H685 H8 K0 K3 K353 M1 M122 M141 M280  
M311 M312 M321 M332 M342 M344 M362 M373  
M391 M413 M510 M521 M532 M540 M710 P411  
P442 P446 P448 P527 Specific Compounds  
RA5BTB Registry Numbers 457374

Chemical Indexing M2 \*90\* Fragmentation Code  
C316 F012 F431 G013 G030 G111 G563 H5 H541  
H6 H685 H8 K0 K3 K353 M1 M123 M141 M280  
M311 M312 M321 M332 M342 M344 M362 M373  
M391 M413 M510 M521 M531 M541 M710 P411  
P442 P446 P448 P527 Specific Compounds  
RA5BTC Registry Numbers 457375

Chemical Indexing M2 \*91\* Fragmentation Code  
C316 F012 F019 F431 F499 G013 G019 G100 H5  
H521 H541 H6 H685 H8 K0 K3 K353 M1 M121 M123  
M141 M149 M280 M311 M312 M321 M332 M342  
M344 M362 M373 M391 M413 M510 M522 M532  
M540 M710 P411 P442 P446 P448 P527 Specific

## Compounds RA5BTD Registry Numbers 457376

Chemical Indexing M2 \*92\* Fragmentation Code  
F013 F431 G010 G012 G030 G112 G530 H5 H541  
H8 J0 J011 J3 J341 M1 M121 M123 M136 M141  
M280 M311 M321 M342 M373 M391 M413 M510  
M521 M532 M541 M710 P411 P442 P446 P448  
P527 Specific Compounds RA5BTE Registry  
Numbers 457377

Chemical Indexing M2 \*93\* Fragmentation Code  
F013 F431 G010 G011 G030 G112 G530 H5 H541  
H8 J0 J011 J3 J341 M1 M121 M123 M136 M141  
M280 M311 M321 M342 M373 M391 M413 M510  
M521 M532 M541 M710 P411 P442 P446 P448  
P527 Specific Compounds RA5BTF Registry  
Numbers 457378

Chemical Indexing M2 \*94\* Fragmentation Code  
F013 F431 G013 G019 G030 G112 G530 H5 H541  
H6 H602 H641 H8 J0 J011 J3 J341 M1 M121 M123  
M136 M141 M280 M311 M321 M342 M373 M391  
M413 M510 M521 M532 M541 M710 P411 P442  
P446 P448 P527 Specific Compounds RA5BTG  
Registry Numbers 457379

Chemical Indexing M2 \*95\* Fragmentation Code  
F013 F431 G012 G013 G030 G112 G530 H5 H541  
H6 H685 H8 J0 J011 J3 J341 M1 M121 M123 M136  
M141 M280 M311 M322 M342 M344 M353 M373  
M391 M413 M510 M521 M532 M541 M710 P411  
P442 P446 P448 P527 Specific Compounds  
RA5BTH Registry Numbers 457380

Chemical Indexing M2 \*96\* Fragmentation Code  
C316 F013 F019 F211 F431 G013 G019 G100 H5  
H521 H541 H8 K0 K3 K353 M1 M121 M123 M141  
M149 M210 M213 M231 M271 M281 M311 M321  
M342 M373 M391 M413 M510 M522 M532 M540  
M710 P411 P442 P446 P448 P527 Specific  
Compounds RA5BTM Registry Numbers 457385

## Chemical Indexing M2 \*97\* Fragmentation Code

B614 B711 B712 B713 B720 B831 B832 B833 C216  
C316 F010 F012 F013 F014 F015 F016 F019 F021  
F029 F211 F299 F431 F499 G003 G010 G011 G012  
G013 G014 G015 G016 G017 G019 G030 G039  
G050 G100 G111 G112 G113 G553 G563 H341  
H342 H343 H401 H402 H403 H441 H442 H443 H521  
H522 H523 H541 H542 H543 H600 H601 H608 H609  
H621 H622 H641 H642 H643 H681 H682 H683 H685  
H689 H715 H721 H722 H723 J011 J321 J341 J351  
J371 J581 J582 J583 K353 K442 K499 L143 L199  
L462 L463 L930 L941 L943 L999 M111 M113 M119  
M121 M123 M126 M129 M131 M136 M137 M139  
M141 M142 M147 M149 M150 M210 M211 M212  
M213 M214 M215 M216 M220 M221 M222 M223  
M224 M225 M226 M231 M232 M233 M240 M262  
M271 M272 M280 M281 M282 M283 M311 M312  
M313 M314 M315 M316 M321 M322 M323 M331  
M332 M333 M340 M342 M344 M353 M362 M373  
M391 M392 M411 M413 M510 M521 M522 M523  
M531 M532 M533 M540 M541 M542 M543 M630  
M640 M650 M710 P411 P442 P446 P448 P527  
Markush Compounds 004819701

## Chemical Indexing M2 \*98\* Fragmentation Code

B614 B711 B720 B831 C216 C316 D021 D022 D023  
D140 F010 F012 F013 F014 F015 F016 F019 F021  
F211 F431 F499 G003 G010 G011 G012 G013  
G019 G030 G031 G032 G039 G050 G051 G100  
G111 G112 G113 G553 G563 H341 H361 H401  
H441 H461 H521 H522 H523 H541 H561 H600 H601  
H608 H609 H621 H622 H641 H661 H681 H682 H683  
H685 H689 H715 H721 J011 J321 J341 J351 J361  
J371 J581 K353 K442 L143 L144 L462 L463 L930  
L941 L943 L999 M112 M113 M115 M116 M122  
M123 M125 M126 M129 M131 M136 M137 M141  
M142 M147 M149 M150 M210 M211 M212 M213  
M214 M215 M216 M220 M221 M222 M223 M224  
M225 M226 M231 M232 M233 M240 M262 M271  
M272 M280 M281 M282 M283 M311 M312 M313

M314 M315 M316 M321 M322 M323 M331 M332  
M333 M340 M342 M344 M353 M362 M373 M391  
M392 M411 M412 M413 M510 M511 M521 M522  
M530 M531 M532 M533 M540 M541 M542 M543  
M630 M640 M650 M710 P411 P442 P446 P448  
P527 Markush Compounds 004819702

**SECONDARY-ACC-NO:**

**CPI Secondary Accession Numbers:** 2001-169537